

Preservation and detection of biomarkers in mineralized communities and its potential link to the variations in ORP

L. Garcia-Descalzo¹ and. F. Gomez², and the MASE team: C.S. Cockell (UK), P. Schwendner (UK), F. Westall, F. Gaboyer, P. Rettberg, K. Beblo-Vranesevic, M. Bohmeier, E. Rabbow, M. Moissl-Eichinger (A), A. Perras (A), R. Amils, (SP), P. Ehrenfreund, E. Monaghan (NL), V. Marteinson, P. Vannier (IS), M. Malki (SP) N. Walter (F), P. Cabezas (F). ^{1,2}Extremophiles Laboratory. Center for Astrobiology (INTA-CSIC) Ctra. Ajalvir Km4. 28850 Torrejon de Ardoz. Madrid. Spain. ¹garciadl@cab.inta-csic.es. ²gomezfg@cab.inta-csic.es.

Abstract

The search for traces of life is one of the principal aims of Mars exploration. Within MASE project (Mars Analogues for Space Exploration) we work to improve approaches and methods for biomarker detection and extraction from Mars analogue sites. One promising strategy to study the preservation of biosignatures in Mars conditions consists of the combined study of biomarkers detection and monitoring physicochemical parameters in mineralized samples. We have observed that it seems to be a potential correlation between biomarker detection and changes in oxi-reduction potential (ORP) over mineralization process. In addition, these methods and the study of samples from MASE sites have enabled us to develop an antibody microarray for competitive sandwich immunoassays as a potential tool for the detection of biomarkers in salty and anaerobic conditions.

1. Introduction

Physico-chemical processes of living organisms leave tell-tale signals in the environment. Its search entails one of the main goals for Astrobiology and improving and optimizing its detection regarding Mars conditions is part of the MASE project objectives. Besides, the traces of some kinds of microorganisms can be well preserved, provided that they are rapidly mineralized and that the sediments in which they occur are rapidly cemented [1].

A developed antibody multiarray competitive immunoassay (MACIA) for the simultaneous detection of compounds of a wide range of molecular sizes or whole spores and cells [2] [3] has revealed as suitable option for biomarker detection in samples with low biomass from Mars analogue sites as well as with biomineralized microorganism communities.

Moreover, biomineralization is often the first step of fossilization and produces particular chemical, structural and morphological features that can be preserved in fossil biominerals or microfossils [4] and some parameters as ORP or pH vary over the process.

2. Methods and objectives

Samples from the three MASE campaigns in Iceland (Graenvatun Lake), United Kingdom (Boulby Mines) and Germany (Regensburg) were used to obtain enrichments and isolates as well as to extract and detect biomarkers presence in them. Some of the enrichments were led to mineralization to study, among others, the preservation of biosignatures by the assessment of antigen-antibody binding at different times. Simultaneously, the evolution of ORP through this process was monitored by two modules system (DTIVA: automated tools for microbial life detection) where ORP variations in those communities were followed through continuous measurements of nanosensors in closed chambers.

An additional objective for MASE project has been to develop a specific microarray with antibodies performed from natural samples and isolates from MASE sample sites.

3. Summary and Conclusions

The presence of traces from some microorganism metabolic groups were detected in the mineralized communities at three different times over the fossilization process. It was undertaken by using a 168 antibody microarray for the immunoassay. There were observed variations in the resulting immunoprofiles. There seems to be a probably correlation between these changes and those in ORP through time. We consider that the simultaneous use of both approaches arises a promising tool to broaden

the knowledge and consequent improvement in the search for traces of past and present life.

Acknowledgements

MASE is supported by European Community's Seventh Framework Programme (FP7/2007-2013) under Grant Agreement n° 607297.

References

[1] Westall F, Foucher F, Bost N, Bertrand M, Loizeau D, Vago JL, Kminek G, Gaboyer F, Campbell KA, Bréhéret JG, Gautret P, Cockell CS: Biosignatures on Mars: What, Where, and How? Implications for the Search for Martian Life *Astrobiology*. 15(11):998-1029. 2015.

[2] Fernández-Calvo, P., Näge, C., Rivas, L. A., García-Villadangos, M., Gómez-Elvira, J., & Parro, V: A multi-array competitive immunoassay for the detection of broad-range molecular size organic compounds relevant for astrobiology *Planetary and Space Science*, 54, 1612-1621. 2006.

[3] Parro V, Fernández-Calvo P, Rodríguez Manfredi JA, Moreno-Paz M, Rivas LA, García-Villadangos M, Bonaccorsi R, González-Pastor JE, Prieto-Ballesteros O, Schuerger AC, Davidson M, Gómez-Elvira J, Stoker CR: SOLID2: an antibody array-based life-detector instrument in a Mars Drilling Simulation Experiment (MARTE). *Astrobiology*. 8(5):987-99. 2008.

[4] Jinhua Lia, Karim Benzeraraa, Sylvain Bernardb, Olivier Beyssaca: The link between biomineralization and fossilization of bacteria: Insights from field and experimental studies *Chemical Geology*. 359:49-69. 2013.