

[Sign on](#)[SAO/NASA ADS](#) [Astronomy Abstract Service](#)

- [Find Similar Abstracts](#) (with [default settings below](#))
- [Also-Read Articles](#) ([Reads History](#))
-
- [Translate This Page](#)

Title: Ceres Evolution: From Thermodynamic Modeling and Telescope to Dawn Orbital Observations

Authors: [McCord, Thomas B.](#) ; [Combe, Jean Philippe](#) ; [Castillo-Rogez, Julie C.](#) ;
[Raymond, Carol A.](#) ; [De Sanctis, Maria Cristina](#) ; [Jaumann, Ralf](#) ;
[Ammannito, Eleonora](#) ; [Russell, Christopher](#)

Affiliation: AA(The Bear Fight Institute), AB(The Bear Fight Institute), AC(Jet Propulsion Laboratory), AD(Jet Propulsion Laboratory), AE(1 Istituto di Astrofisica e Planetologia Spaziali – INAF), AF(DLR, Institute of Planetary Research), AG(University of California at Los Angeles), AH(University of California at Los Angeles)

Publication: American Astronomical Society, DPS meeting #47, id.#212.11

Publication Date: 11/2015

Origin: [AAS](#)

Abstract Copyright: (c) 2015: American Astronomical Society

Bibliographic Code: [2015DPS....4721211M](#)

Abstract

Thermodynamic modeling indicated that Ceres has experienced planetary processes, including extensive melting of its ~25% water and differentiation, (McCord and Sotin, JGR, 2005; Castillo and McCord, Icarus, 2009). Early telescopic studies showed Ceres' surface to be spectrally similar to carbaceous-chondrite-like material, i.e., aqueously altered silicates darkened by carbon, with a water-OH-related absorption near 3.06 μm . Later observations improved the spectra and suggested more specific interpretations: Structural water in clay minerals, phyllosilicates, perhaps ammoniated, iron-rich clays, carbonates, brucite, all

implying extensive aqueous alteration, perhaps in the presence of CO₂. Telescopic observations and thermodynamic models predicted Dawn would find a very different body compared to Vesta (e.g. McCord et al., SSR, 2011), as current Dawn observations are confirming. Ceres' original water ice should have melted early in its evolution, with the resulting differentiation and mineralization strongly affecting Ceres' composition, size and shape over time. The ocean should have become very salty and perhaps may still be liquid in places. The surface composition from telescopes seems to reflect this complex history. The mineralization with repeated mixing of the crust with the early liquid interior and with in-fall from space would create a complex surface that will present an interpretation challenge for Dawn. The Dawn spacecraft is currently collecting observations of Ceres' landforms, elemental and mineralogical/molecular composition and gravity field from orbit. Early results suggest a heavily cratered but distorted and lumpy body with features and composition consistent with internal activity, perhaps recent or current, associated with water and perhaps other volatiles. We will present and interpret the latest Dawn Ceres findings and how they affect our earlier understanding of Ceres evolution from modeling and telescope observations.

[Bibtex entry for this abstract](#) [Preferred format for this abstract](#) (see [Preferences](#))

Add this article to private library

Remove from private library

Submit corrections to this record

View record in ADSLabs **NEW!**

Find Similar Abstracts:

Use:

- ☐ Authors
- ☒ Title
- ☒ Abstract Text

Return:

- ☒ Query Results Return **100**
- ☐ Query Form

items starting with number **1**

Database:

- ☒ Astronomy
- ☐ Physics
- ☐ arXiv e-prints

Send Query

Reset
