

NF- κ B ACTIVATION AND EXPRESSION OF ITS TARGET GENES AFTER HEAVY IONS EXPOSURE

Chishti AA ^{1,2}, Koch K ¹, Spitta LF ¹, Henschenmacher B ¹, Diegeler S ¹, Konda B ¹,
Baumstark-Khan C ¹, Feles S ¹, Schmitz C ¹, and Hellweg CE ¹

1) German Aerospace Centre (DLR), Institute of Aerospace Medicine, Radiation Biology Department, Linder Höhe, D-51147 Köln, Germany

2) Present address: The Karachi Institute of Biotechnology and Genetic Engineering (KIBGE), University of Karachi, 75270 Karachi, Pakistan

Introduction: As a prerequisite for developing appropriate countermeasures to mitigate acute effects and late radiation risks for the astronaut and thereby enabling long-term human space flight, the cellular radiation response to densely ionizing radiation needs to be better understood. The biological effectiveness of accelerated heavy ions (which constitute the biologically most important radiation type in space) with high linear energy transfer (LET) for effecting DNA damage response pathways as a gateway to cell death or survival is of major concern not only for space missions but also for new regimes of tumor radiotherapy.

Methods: The contribution of NF- κ B to the cellular response to space-relevant radiation qualities was determined by a NF- κ B reporter cell line (HEK-pNF- κ B-d2EGFP/Neo L2). The NF- κ B dependent reporter gene expression (d2EGFP) after ionizing radiation (X-rays and heavy ions in broad range of LET: 0.3 - 9674 keV/ μ m) exposure was evaluated by flow cytometry. Survival was determined by the colony forming ability test. The biological effectiveness (RBE) of NF- κ B activation and reduction of cellular survival were calculated for each radiation quality. Furthermore, the effect of LET on NF- κ B target gene expression was analysed by real time reverse transcriptase quantitative PCR (RT-qPCR).

Results: NF- κ B activation and NF- κ B dependent gene expression comprises occurred as an early step in the cellular radiation response. NF- κ B activation and NF- κ B-dependent gene expression by heavy ions were highest in the LET range of 50-300 keV/ μ m. The expression of several chemokines and cytokines (CXCL1, CXCL2, CXCL10, IL-8 and TNF) was up-regulated and the extent of upregulation depended on LET.

Conclusion: The upregulated chemokines and cytokines might be important for cell-cell communication among hit as well as unhit cells (bystander effect). Their role in the cellular and tissue response to ionizing radiation needs to be further examined as they might induce proinflammatory effects.

Acknowledgement: AAC, KK, BH, SD, BK were supported by the Helmholtz Space Life Sciences Research School (SpaceLife).

GBS Annual Meeting 2016, 26. – 28. September 2016, in Erlangen

Poster

<http://www.strahlenklinik.uk-erlangen.de/fort-und-weiterbildung/gbs-2016/>