

CONTROL ID: 2568059

TITLE: VIRTIS-M observations of 67P/Churyumov-Gerasimenko at low phase angle

ABSTRACT BODY:

Abstract (2,250 Maximum Characters): On 9-10 of April 2016, the Rosetta orbiter completed a close flyby around the nucleus of 67P/Churyumov-Gerasimenko (CG), when the comet was at heliocentric distance of 2.76 AU, along the outbound leg of its orbit around the Sun. This allowed the VIRTIS imaging spectrometer to observe the surface of CG at visible wavelengths (0.2-1 μm) in the 0.93° - 89.7° phase angle range with a spatial resolution ranging from 7 to 46 m/pix, resulting in a total of 105 hyperspectral images. Previous observations of the comet at low phase angle were acquired by VIRTIS during the approach phase on July 2014 (3.7 AU), with pixel resolution varying from 450 to 3200 m/pix, preventing disk-resolved imaging. The April 2016 observations fill this gap, allowing us to constrain the spectrophotometric properties of the surface at higher spatial resolution in the opposition surge geometry, therefore investigating both the 'shadow hiding' and 'coherent backscattering' opposition effect. Extrapolation to 0° phase angle of the reflectance measured during the flyby at latitude between -10° and 30° indicates a surface normal albedo of 0.06 at 0.55 μm . This result is close to the previous average value derived by Ciarniello et al. (2015) from a full-disk analysis giving 0.062 ± 0.002 (3.7 AU, inbound orbit). Subsequent measurements by Filacchione et al. (2016) and Ciarniello et al. (in preparation), derived from pre-perihelion observations at lower heliocentric distances, revealed a progressive enrichment of water ice abundance on the nucleus, in northern hemisphere regions, with a consequent brightening of the surface. The measured normal albedo derived from the 9-10 April 2016 dataset seems to indicate that during the post-perihelion phase the northern hemisphere has returned to albedo values compatible to the ones measured during the inbound part of the orbit, when the comet was at 3.7 AU from the Sun. This could be an indication that the northern hemisphere has been either covered by dust emitted by the southern hemisphere during the very active perihelion phase, or that dehydration of the upper surface layers has taken place.

CURRENT * CATEGORY: Comet Physical Characteristics: Surfaces

CURRENT : None

AUTHORS (FIRST NAME, LAST NAME): Mauro Ciarniello¹, Gianrico Filacchione¹, Fabrizio Capaccioni¹, Andrea Raponi¹, Maria Cristina De Sanctis¹, Federico Tosi¹, Giuseppe Piccioni¹, Alessandra Migliorini¹, Priscilla Cerroni¹, Maria Teresa Capria¹, Stéphane Érard², Dominique Bockelee-Morvan², Cedric Leyrat², Gabriele Arnold³, Maria A. Barucci²,

Bernard Schmitt⁴, Eric Quirico⁴, Fredric W. Taylor⁵, David Kappel³,
Andrea Longobardo¹

INSTITUTIONS (ALL): 1. INAF-IAPS, Rome, Rome, Lazio, Italy, Italy.
2. LESIA, Observatoire de Paris, Paris, France.
3. DLR, Institute of Planetary Research, Berlin, Germany.
4. IPAG, Grenoble, France.
5. University of Oxford, Oxford, United Kingdom.

Contributing Teams: VIRTIS Team