



Evaluation of gas-particle partitioning in a regional air quality model for organic pollutants

Christos I. Efstathiou¹, Jana Matejovičová^{1,2}, Johannes Bieser^{3,4}, and Gerhard Lammel^{1,5}

¹Masaryk University, Research Centre for Toxic Compounds in the Environment, Kamenice 5, 62500 Brno, Czech Republic

²Slovak Hydrometeorological Institute, Jeséniova 17, 83315 Bratislava, Slovak Republic

³Institute of Coastal Research, Helmholtz-Zentrum Geesthacht, Max-Planck-Str. 1, 21502 Geesthacht, Germany

⁴German Aerospace Center (DLR), Oberpfaffenhofen, 82234 Weßling, Germany

⁵Max Planck Institute for Chemistry, Multiphase Chemistry Department, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

Correspondence to: Christos I. Efstathiou (efstathiou@recetox.muni.cz)

Received: 22 January 2016 – Published in Atmos. Chem. Phys. Discuss.: 29 February 2016

Revised: 23 September 2016 – Accepted: 9 November 2016 – Published: 9 December 2016

Abstract. Persistent organic pollutants (POPs) are of considerable concern due to their well-recognized toxicity and their potential to bioaccumulate and engage in long-range transport. These compounds are semi-volatile and, therefore, create a partition between vapour and condensed phases in the atmosphere, while both phases can undergo chemical reactions. This work describes the extension of the Community Multiscale Air Quality (CMAQ) modelling system to POPs with a focus on establishing an adaptable framework that accounts for gaseous chemistry, heterogeneous reactions, and gas-particle partitioning (GPP). The effect of GPP is assessed by implementing a set of independent parameterizations within the CMAQ aerosol module, including the Junge–Pankow (JP) adsorption model, the Harner–Bidleman (HB) organic matter (OM) absorption model, and the dual Dachs–Eisenreich (DE) black carbon (BC) adsorption and OM absorption model. Use of these descriptors in a modified version of CMAQ for benzo[a]pyrene (BaP) results in different fate and transport patterns as demonstrated by regional-scale simulations performed for a European domain during 2006. The dual DE model predicted 24.1 % higher average domain concentrations compared to the HB model, which was in turn predicting 119.2 % higher levels compared to the baseline JP model. Evaluation with measurements from the European Monitoring and Evaluation Programme (EMEP) reveals the capability of the more extensive DE model to better capture the ambient levels and seasonal behaviour of BaP. It is found that the heterogeneous reaction of BaP with O₃ may decrease its atmospheric lifetime by 25.2 % (domain and

annual average) and near-ground concentrations by 18.8 %. Marginally better model performance was found for one of the six EMEP stations (Košetice) when heterogeneous BaP reactivity was included. Further analysis shows that, for the rest of the EMEP locations, the model continues to underestimate BaP levels, an observation that can be attributed to low emission estimates for such remote areas. These findings suggest that, when modelling the fate and transport of organic pollutants on large spatio-temporal scales, the selection and parameterization of GPP can be as important as degradation (reactivity).

1 Introduction

Polycyclic aromatic hydrocarbons (PAHs) are a group of lipophilic organic compounds with demonstrated carcinogenicity and potential to bioaccumulate (Diamond and Hodge, 2007; Finlayson-Pitts and Pitts, 1999; Pedersen et al., 2004, 2005; WHO, 2003). Typically emitted to the atmosphere as a mixture of semi- to low-volatile congeners with variable carcinogenic potency, their fate and transport in the environment is difficult to assess. Current limitations involve inadequate knowledge of photochemistry (Keyte et al., 2013), air–surface exchange (Galarneau et al., 2014; Keyte et al., 2013; Lammel et al., 2009), and combustion sources (Bieser et al., 2012; Lammel et al., 2013). In addition, aluminium and steel production, along with petrogenic sources, are significant in some environments (Wenborn et al., 1999).

This relationship with anthropogenic activity and population density raised awareness of their potential as health stressors, with hotspots identified in urban industrial settings around the world (Šrám et al., 2013; Zhu et al., 2015). On the other hand, observations of PAHs at remote sites and global circulation model studies indicate long-range transport (Keyte et al., 2013; Sehili and Lammel, 2007). Therefore, and due to resistance to degradation, PAHs are classified as persistent organic pollutants (POPs) by the United Nations Economic Commission for Europe (UNECE) Convention on Long-range Transboundary Air Pollution (CLRTAP) and are listed in the Protection of the Marine Environment of the North-East Atlantic (OSPAR).

Much of the scientific literature is focussed on the 16 US EPA priority PAHs, i.e. acenaphthene (Ace), acenaphthylene (Acy), fluorene (Flu), naphthalene (Nap), anthracene (Ant), fluoranthene (Flt), phenanthrene (Phe), benzo[a]anthracene (BaA), benzo[b]fluoranthene (BbF), benzo[k]fluoranthene (BkF), chrysene (Chr), pyrene (Pyr), benzo[ghi]perylene (BgP), benzo[a]pyrene (BaP), dibenzo[a,h]anthracene (DBahA), and indeno[1,2,3-cd]-pyrene (IPy), albeit without justification (Andersson and Achten, 2015). Used as a biomarker for the carcinogenic risk of PAHs, BaP is a criteria pollutant in many countries, sometimes together with a few other bioaccumulative members. While in North America no federal guidelines are currently in effect, the European Union indicative limit value is set at 1 ng m^{-3} of BaP (EC, 2004), and certain countries have established stricter air quality standards (i.e. 0.25 ng m^{-3} of BaP in the United Kingdom) (DETR, 2009).

A number of studies, which aimed at exploring the atmospheric fate and transport of POPs on various scales, have been presented in the last two decades. These include relatively simple mass balance models developed under the Lagrangian framework, some of which have been extended to multimedia applications for studying the accumulation and exchange of POPs between various compartments (Halsall et al., 2001; Lang et al., 2008, 2007; Liu et al., 2007; Prevedouros et al., 2004, 2008; Van Jaarsveld et al., 1997). Eulerian grid models add yet another level of complexity regarding process detail that is accompanied by additional computational demands. Global-scale chemical transport models (Friedman and Selin, 2012; Lammel et al., 2009; Sehili and Lammel, 2007), as well as hemispheric models (Gusev et al., 2005; Hansen et al., 2006; Shatalov et al., 2005), have been applied to study the long-range transport of POPs. Moreover, coupled meteorological and air quality modelling frameworks have presented efforts to further investigate the treatment of physical and chemical atmospheric processes related to POPs in higher resolution (Aulinger et al., 2007; Bieser et al., 2012; Cooter and Hutzell, 2002; Galarneau et al., 2014; Inomata et al., 2012; Matthias et al., 2009; Meng et al., 2007; San José et al., 2013; Silibello et al., 2012; Zhang et al., 2009). Table S1 in the Supplement provides a summary of the most relevant processes covered in the respective regional

model implementations and subsequent evolution tracked by the authors. While such studies represent the initial efforts to accurately model POPs on regional scale, important processes such as gas-particle partitioning, heterogeneous reactivity, and a multicompartamental approach for the relevant compounds are not always studied. The Weather Research Forecast along with the Community Multiscale Air Quality for BaP (WRF-CMAQ-BaP) model presented in this article is considered to be the first effort to examine the full spectrum of gas-particle partitioning models relevant to POPs in the same regional-scale modelling application. The model attempts to capture POP aerosol chemistry and dynamics within the CMAQ framework and can be extended to account for mass exchange between adjacent compartments for compounds for which such processes are deemed critical (i.e. soil volatilization).

2 Model development

The model development framework is based on the CMAQ version 4.7.1 modelling system (Byun and Schere, 2006; Byun and Ching, 1999) with the CB05 gas-phase chemistry mechanism (Yarwood et al., 2005) and the AERO4 version of the aerosol module (Binkowski, 2003). In addition, CMAQ contains modules representing advection, eddy diffusion, and in-cloud and below-cloud scavenging with precipitation. Advection and diffusion satisfy mass conservation and include removal by dry deposition, while in-cloud and precipitation processes simulate aqueous chemistry and wet deposition by cloud droplets (Byun, 1999a,b). The aerosol module (Binkowski, 2003) determines concentrations of trimodal size-distributed particulate material with diameters less than $10 \mu\text{m}$. According to this model, each mode can be described as an internal mixture of several compounds; the mixing state on the single-particle level is not addressed, but is considered equivalent to internal mixing. Aitken and accumulation modes include particles with diameters $\sim 2.5 \mu\text{m}$ or less ($\text{PM}_{2.5}$), which are either emitted or produced by gas-to-particle conversion processes. A thermodynamic mechanism, based on temperature and relative humidity, determines air concentrations of water, sulfate, ammonium, and nitrate aerosols in the fine modes (Nenes et al., 1998; Zhang et al., 2000). Organic and elemental carbon species are also included in these modes. The third mode represents coarse material having diameters between 2.5 and $10 \mu\text{m}$ and includes dry inorganic material emitted by natural and anthropogenic sources. Other processes like coagulation, particle growth, aerosol–cloud interactions, and new particle formation are included in the treatment.

CMAQ adaptations are required to fully address the behaviour of semi-volatile organic compounds (SOCs) in the atmosphere and include additional homogeneous gas-phase reactions, a modular algorithm to describe the mass exchange of SOCs between the gas and particulate phases, and a de-

scription of deposition for each phase state of the compound of interest. For the case of benzo[a]pyrene (BaP), one gas-phase species and three particulate-sorbed species (one for each aerosol mode) are introduced. It is assumed that only the gaseous molecule undergoes chemical degradation through reaction with the hydroxyl (OH) radical, leading to an unspecified product that does not undergo further chemical reaction. A summary of the physico-chemical data used in this study is presented in Table 1. Furthermore, a module that accounts for heterogeneous chemistry was included for compounds for which such reactions represent a significant loss according to experimental studies (e.g. BaP reaction with O₃).

2.1 Modular implementation of gas-particle partitioning (GPP) in CMAQ

The distribution of POPs between the gas and the particulate phases is widely recognized as the most important parameter in describing their fate and transport in the atmosphere. PAHs like benzo[a]pyrene can be adsorbed to mineral surfaces or elemental carbon, absorbed by organic aerosol, or absorbed by aerosol water (Lohmann and Lammel, 2004). Therefore, the aerosol module of CMAQ was enhanced with an array of models that assume different processes to determine gas-particle partitioning, i.e. an adsorption model (Junge–Pankow, Junge, 1977; Pankow, 1987), an organic matter (OM) absorption model (K_{oa} , Harner and Bidleman, 1998), and a dual OM absorption and elemental or black carbon (EC/BC) adsorption model (Dachs and Eisenreich, 2000; Lohmann and Lammel, 2004). In addition, a method for calculating the absorption in particulate-phase water was included in the module, employed for all aerosol modes in a manner similar to the GPP processes (Aulinger et al., 2007; Cooter and Hutzell, 2002).

The new GPP algorithm for CMAQ brings together existing implementations for pesticides (Cooter and Hutzell, 2002), PAHs (Aulinger et al., 2007; San José et al., 2013), polychlorinated biphenyls (PCBs) (Meng et al., 2007), dibenzo-*p*-dioxins, and dibenzofurans (PCDD/Fs) (Zhang et al., 2009) in a unified module that allows for flexible GPP model combinations, making it applicable to a wide range of SOCs. The following assumptions were taken into consideration: (1) instantaneous relaxation to phase equilibrium (no mass transport kinetic limitations), (2) the compound does not irreversibly transform in the particulate phase, and (3) sorption processes do not interact. By extending the gas-phase and modal aerosol treatments in CMAQ for SOCs, these assumptions allow the use of simple ratios to determine the concentration of each form, following

$$a_i = \phi_i (a_i + g), \quad (1)$$

where a_i is the particulate concentration in each mode ($i = 1 = \text{Aitken}, 2 = \text{accumulation}, 3 = \text{coarse}$), g is the gaseous concentration, and ϕ_i is the particulate fraction of the com-

ound in mode, with a general formulation (Aulinger et al., 2007; Cooter and Hutzell, 2002).

$$\phi_i = \phi_i^{\text{ad}} + \phi_i^{\text{ab}} + \phi_i^{\text{aq}} \quad (2)$$

This way, all partitioning processes, adsorption (ϕ_i^{ad}), absorption in OM (ϕ_i^{ab}), and absorption (solution) in aerosol water (ϕ_i^{aq}), are simultaneously treated. Following the rule of mass conservation for the compound in both phases, the total concentration c_{tot} can be introduced:

$$g = c_{\text{tot}} - a_1 - a_2 - a_3, \quad (3)$$

leading to a system of three linear equations.

$$\begin{bmatrix} 1 & \phi_1 & \phi_1 \\ \phi_2 & 1 & \phi_2 \\ \phi_3 & \phi_3 & 1 \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = c_{\text{tot}} \times \begin{bmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{bmatrix}, \quad (4)$$

which yield the particulate concentrations a_i in each mode. Subsequently, using the previous equation, the gaseous concentration g can be calculated.

2.1.1 Junge–Pankow adsorption model

The first GPP scheme is based on the model of exchangeable SOC adsorption to aerosols presented by Junge (1977) and later critically reviewed by Pankow (1987).

$$\phi_i^{\text{ad}} = c_J S_i / (p_L^\circ + c_J S_i) \quad (5)$$

The equation above relates the mass fraction of chemical adsorbed particles in each mode i (ϕ_i^{ad}) to the subcooled liquid vapour pressure of the compound (p_L° , Pa) and the aerosol surface density (S_i , m² of particle surface per m³ of air). The parameter c_J (unit, Pa m) depends on the chemical nature of both the adsorbent and the adsorbate and cannot be expected to be a constant, as confirmed by field experiments (Lammel et al., 2010). Nevertheless, a value of $c_J = 0.172$ Pa m has been suggested (Pankow, 1987) and is often used in modelling applications. The surface area parameter, S_i , which is a characteristic of the aerosol required for the Junge–Pankow (JP) model, is calculated at each time step for each mode within the aerosol module of CMAQ (Binkowski, 2003).

The empiric JP model, despite assuming adsorption to be the only relevant partitioning process, may eventually not be free from absorptive contributions (to c_J), but these should be very low. The fitting in the empiric parameterization was done for all possible vapours, many of these not soluble in octanol. No c_J for BaP derived from ambient measurements has ever been published. However, we recently found in background aerosol of central Europe (Košetice 2012–2013, $n = 162$ aerosol samples, c_J derived from experimental S/V data; Shahpoury et al., 2016) that c_J for BaP is very low, at 0.01–5.46 Pa cm (median 0.44 Pa cm), supporting the perception that a possible implicit absorptive contribution to c_J must be very low.

Table 1. Physicochemical parameters and reaction rate coefficients used in the WRF-CMAQ-BaP model.

Inverse Henry's law constant ¹	$\log H'$	-4.7
Subcooled liquid vapour pressure ^{1,2}	$\log p_L$	-5.2 (Pa)
Octanol–water partition coefficient ¹	$\log K_{ow}$	5.9
Octanol–air partition coefficient ^{1,3}	$\log K_{oa(dry)}$	11.1 ¹ ($b = 12.04$, $m = 5382$) ³
Water-saturated octanol–air partition coefficient ¹	$\log K_{oa(wet)}$	10.6
Diesel soot–water partition coefficient ⁴	$\log K_{soot-water}$	8.4
Diesel soot–air partition coefficient ^{4,8}	$\log K_{soot-air}$	13.0 ⁴ , 11.59 ⁸
Gas-phase OH reaction rate constant	k_{OH}	50×10^{-12} (cm ³ mol ⁻¹ s ⁻¹)
Ozone gas–surface equilibrium constant ⁶	K_{O_3}	0.028×10^{-13} (cm ³)
Maximum rate coefficient ⁶	k_{max}	0.060 (s ⁻¹)

¹ Beyer et al. (2000). ² de Maagd et al. (1998). Offenber and Baker (1999). ³ Odabasi et al. (2006) $\log K_{oa} = m T^{-1}$ (K) + b .

⁴ Bärning et al. (2002), $\log K_{soot-water} = 1.39 \log K_{ow} + 0.1$. ⁵ $\log K_{soot-air} = \log K_{soot-water} - \log H'$. ⁶ Estimated values (EPIWIN;USEPA). ⁷ Kwamena et al. (2004). ⁸ van Noort (2003).

2.1.2 K_{oa} absorption model

The second GPP scheme follows the work of Pankow (1994, 2007), with refinements from Harner and Bidleman (1998) that described how the particulate fraction of a compound could be derived from the aerosol–air partitioning coefficient K_p . Following this approach for each mode i , the predicted particulate fraction ϕ_i^{ab} is linked to the total mass concentration of suspended particles c_{TSP_i} with the following equation:

$$\phi_i^{ab} = K_{p,i}^{OM} c_{TSP_i} / (K_{p,i}^{OM} c_{TSP_i} + 1). \quad (6)$$

Pankow (1987, 1998) demonstrated that the gas-particle partitioning coefficient, K_p , and the use of p_L° as its descriptor are valid both for adsorption and absorption of SOCs. In the case of absorptive partitioning, Pankow (1998) derived

$$K_{p,i}^{OM} (m^3 \mu g^{-1}) = \frac{f_{OM_i} 760RT}{10^6 M_{OM} \gamma_{OM} p_L^\circ}, \quad (7)$$

where f_{OM_i} is the mass fraction of OM in the particle that can absorb gaseous SOCs, R is the ideal gas constant ($= 8.314 \text{ Pa m}^3 \text{ mol}^{-1} \text{ K}^{-1}$), T (K) is absolute temperature, M_{OM} (g mol^{-1}) is the mean molecular weight of the OM phase, γ_{OM} is the activity coefficient of the selected SOC in the OM on a mole fraction basis, and p_L° (Pa) is the vapour pressure of the pure SOC (subcooled liquid in the case of solids). Partitioning coefficients normalized to the OM content were introduced to further support the absorption hypothesis (Finizio et al., 1997). Pankow (2007) and Harner and Bidleman (1998) replaced p_L° by the octanol–air partitioning coefficient, K_{oa} .

$$K_{p,i}^{OM} = 10^{-12} \frac{f_{OM_i} M_{oct} \gamma_{oct} K_{oa}}{M_{OM} \gamma_{OM} \rho_{oct}}, \quad (8)$$

where M_{oct} , M_{OM} (g mol^{-1}) are the molecular weights of octanol ($= 130 \text{ g mol}^{-1}$) and the OM phase, γ_{oct} , γ_{OM} are the activity coefficients of the SOC in octanol and OM, respectively, and ρ_{oct} is the density of octanol (0.82 kg L^{-1}). Assuming that octanol imitates organic matter in PM, Harner

and Bidleman (1998) suggested that the ratio of γ_{oct}/γ_{OM} and M_{oct}/M_{OM} can be assumed to be 1. However, it was later suggested that M_{OM} could be much higher, particularly in secondary organic aerosols containing polymeric structures (Kalberer et al., 2004); a mean value of 500 g mol^{-1} was later suggested by Götz et al. (2007), which results in M_{oct}/M_{OM} of 0.26. Under the assumption that γ_{oct}/γ_{OM} and $M_{oct}/M_{OM} = 1$, the equation above can be simplified to the following:

$$\log K_{p,i}^{OM} = \log K_{oa} + \log f_{OM} - 11.91. \quad (9)$$

2.1.3 Dual OM absorption and black carbon adsorption model

Finally, the third GPP scheme extends the previous OM absorption model to include adsorption onto particulate soot, according to evidence showing the measured K_p to be exceeding the predicted K_p based solely on absorption in OM (Dachs and Eisenreich, 2000; Fernandez et al., 2002; Lohmann and Lammel, 2004; Ngabe and Poissant, 2003). Considering the contribution of those two additive processes, a modified partition coefficient can be employed.

$$K_{p,i}^{dual} = 10^{-12} \left(\frac{f_{OM_i} M_{oct} \gamma_{oct} K_{oa}}{M_{OM} \gamma_{OM} \rho_{oct}} + \frac{f_{BC_i} a_{atm-BC} K_{SA}}{a_{soot} \rho_{BC}} \right), \quad (10)$$

where f_{BC_i} is the fraction of black carbon in the particulate matter, a_{atm-BC} and a_{soot} are the available surfaces of atmospheric black carbon (BC) and diesel soot, respectively, K_{SA} is the partitioning coefficient between diesel soot and air. A density value of 1 kg L^{-1} has often been assumed for BC; however values in the range of $1.7\text{--}1.9 \text{ kg L}^{-1}$ have been measured and recommended by Bond and Bergstrom (2006). Thus, the equation can be simplified.

$$K_{p,i}^{dual} = 10^{-12} (0.32 \times f_{OM_i} K_{oa} + 0.55 \times f_{BC_i} K_{SA}). \quad (11)$$

This method is subject to uncertainties (Goss, 2004), but is accepted and suitable (Dachs et al., 2004 and references

therein). Soot–air partition coefficients (K_{SA}) were calculated as the ratio of soot–water adsorption constants K_{SW} and the inverse Henry’s law constant (H'), with K_{SW} values adopted from Barring et al. (2002). As a step to address uncertainties contributed by different methods in calculating K_{SA} , a thermodynamic estimation model suggested by van Noort (2003) was also tested:

$$\log K_{SA} = -0.85 \log p_L^\circ + 8.94 - \log(998/a_{\text{soot}}). \quad (12)$$

The soot-specific surface area was set to $18.21 \text{ m}^2 \text{ g}^{-1}$, derived as the geometric mean of surface areas for traffic, wood, coal, and diesel soot (i.e. 59.4, 3.6, 8.2, and $62.7 \text{ m}^2 \text{ g}^{-1}$, respectively) (Jonker and Koelmans, 2002). Additionally, in order to estimate uncertainties arising from K_{oa} and its temperature dependence, two different parameterizations were tested, the first based on the work of Beyer et al. (2000) and the second following the temperature-dependent parameterization suggested by Odabasi et al. (2006).

2.2 Dissolution into aerosol water

An assumption of the CMAQ aerosol module is that organics influence neither the water content nor the ionic strength of the system. While this assumption may not be valid for many atmospheric aerosols, sufficient basic data are not available to treat the system in a more complete way. In a similar manner, the dissolution of PAHs in the aqueous electrolyte was calculated using the inverse Henry’s law constant (Aulinger et al., 2007; Cooter and Hutzell, 2002; Lohmann and Lamme, 2004). Lohmann and Lamme (2004) derived the partitioning coefficient $K_{p,i}^{\text{aq}}$, although they conclude that as long as a solid–air interface exists, the contribution of dissolved PAHs to the gas-particle partitioning will always be negligible. To account for this, a “wet aerosol” switch was introduced and associated with the relevant partitioning coefficient.

$$\log K_{p,i}^{\text{aq}} = \log K_{wa} + \log f_{wa,i} - 12.0, \quad (13)$$

where $f_{wa,i}$ is the mass fraction of aerosol water in each mode i and K_{wa} is the water–air partitioning coefficient, which is equal to H' . The aerosol water content is computed in CMAQ based on a variation of the Zdanovskii, Stokes, and Robinson (ZSR) method (Byun and Ching, 1999; Kim et al., 1993; Stokes and Robinson, 1966). As suggested by Aulinger et al. (2007) for CMAQ, only if the ratio of ammonium sulfate to aerosol water drops below the solubility of ammonium sulfate, the aerosol is treated as wet and absorption into aerosol water takes effect, while adsorption to inorganic material is switched off.

2.3 Gas-phase and heterogeneous reactions

Reactions with the hydroxyl radical (OH), ozone (O_3), and the nitrate radical (NO_3) may determine the atmospheric fate and long-range transport potential of SOCs. Gas-phase reactions with OH are believed to be the dominant loss pathway for semi-volatile PAHs (3–4 rings) like anthracene and pyrene (Atkinson and Arey, 1994; Finlayson-Pitts and Pitts, 1999; Keyte et al., 2013). On the other hand, in the case of non-volatile PAHs (5 and more rings, like BaP), loss due to heterogeneous reaction with ozone is believed to dominate (Finlayson-Pitts and Pitts, 1999). Experimental studies of the kinetics of PAHs sorbed to particles indicate strong influences of mixing state (single-particle anisotropy), morphology, and phase state of the particles (Kwamena et al., 2004; Perraudin et al., 2007; Pöschl et al., 2001; Shiraiwa et al., 2009; Zhou et al., 2012). Yet, detailed heterogeneous chemistry models would require corresponding additional parameters and a more sophisticated surface layer model (Kaiser et al., 2011; Shiraiwa et al., 2010; Springmann et al., 2009). However, in the studies by Kahan et al. (2006) and Kwamena et al. (2004, 2007), the reactivity could be well described by a Langmuir–Hinshelwood mechanism (i.e. one species [BaP] is adsorbed to the surface while the second species, ozone, is in phase equilibrium). This approach is followed in the CMAQ aerosol module by introducing the degradation rate coefficient k in the following form:

$$k = \frac{k_{\text{max}} K_{\text{O}_3} c_{\text{O}_3}}{1 + K_{\text{O}_3} c_{\text{O}_3}}, \quad (14)$$

where k_{max} is the maximum rate coefficient ($0.060 \pm 0.018 \text{ s}^{-1}$), K_{O_3} is the ozone gas to surface equilibrium constant ($0.028 \pm 0.014 \times 10^{-13} \text{ cm}^3$), and c_{O_3} is the concentration of ozone (mol cm^{-3}). As this reaction neglects the substance fraction not accessible for gaseous O_3 molecules (so-called burial effect; Zhou et al., 2012), an upper estimate for BaP reactivity is simulated.

2.4 Dry and wet deposition

SOCs and their degradation products are removed from the atmosphere through dry and wet deposition. Dry deposition is modelled in CMAQ as a one-way flux out of the lowest layer of the atmosphere. Particulate forms have dry deposition velocities determined by Brownian diffusion, turbulence, and sedimentation. Mass and diameter of the mode are used to estimate the sedimentation velocity (Binkowski, 2003; Byun and Ching, 1999). The deposition velocity of gaseous molecules is based on an electrical resistance analogue that includes aerodynamic (R_a), quasi-laminar boundary layer (R_b), and surface canopy (R_c) resistance (Pleim et al., 1999). R_a is computed using similarity theory with heat flux. R_b depends on land-use-specific friction velocity and molecular characteristics of gases. R_c consists of several components, including bulk stomatal, dry cuticle, wet cuti-

cle, ground, and in-canopy aerodynamic resistance (Pleim et al., 1999). Cuticle and ground resistances are based on ozone and sulfur dioxide observations and neglect absorption into organic plant or soil material. Values for time-dependent R_c parameters are determined via land use and mesoscale meteorological models, which provide fractional vegetation cover, leaf area index, fractional leaf area wetness, and leaf stress associated with radiation, root zone soil moisture, temperature, and humidity (Pleim and Xiu, 1995).

Wet deposition of gaseous molecules occurs through in-cloud and below-cloud scavenging. Both processes are assumed to have the same scavenging efficiency and dependency on Henry's law, water content of the cloud, and precipitation rate (Chang et al., 1986). Wet scavenging parameterization of the particulate-phase molecules follows the method of Roselle and Binkowski (Byun and Ching, 1999) and depends on the aerosol mode (size) and time required to remove all liquid water from a cloud volume. No ice-phase scavenging is included in the fourth version of the aerosol module in CMAQ.

3 Model implementation and study design

The model domain included the whole of Europe and neighbouring countries as shown in Fig. 1. The domain is within the area covered by the UNECE-CLRTAP (and its POP protocol, covering PAHs) and was specifically chosen to exploit recently prepared high-resolution emission data (Bieser et al., 2011) and make use of the only monitoring network in the region (EMEP; see below). Computational simulations were performed on a 36×36 km Lambert Conformal Conic grid using offline hourly meteorological fields, generated by the Weather Research Forecasting (WRF) model version 3.4.1 using GFS meteorological reanalysis data (spatial resolution $0.5^\circ \times 0.5^\circ$) as input. The parameterization of WRF included the following schemes: Milbrandt–Yau double-moment 7-class microphysics (Milbrandt and Yau, 2005a, b), rapid radiative transfer model (RRTMG) long-wave and shortwave scheme (Iacono et al., 2008), asymmetric convective model of the planetary boundary layer (PBL) (Pleim, 2007), Pleim–Xiu surface layer model (Pleim and Xiu, 2003; Xiu and Pleim, 2001), and the improved version of Grell–Devenyi (Grell, 1993; Grell and Devenyi, 2002) ensemble scheme for cumulus parameterization. On the vertical dimension, the domain was resolved in 36 layers, following the eta coordinate system. Subsequently, CMAQ-ready meteorological input fields for 2006 were prepared using the Meteorology–Chemistry Interface Processor (MCIP) (Otte and Pleim, 2010). Computational simulations were performed in a distributed computing infrastructure operated and managed by the Czech National Grid Organization Meta-Centrum NGI.




Figure 1. Study domain and 2006 annual grid cell mean BaP emission flux (g s^{-1}) at the surface.

3.1 Emissions

Capturing the spatio-temporal pattern of emissions is crucial to determine the fate and transport of PAHs. Several attempts have been carried out during the past decades, with varying spatial and temporal resolutions in historical, current, and future projection terms for Europe (Bieser et al., 2012; Pacyna et al., 2003). Currently, four database sources with gridded BaP emissions exist for Europe and Russia: the TNO database (Denier van der Gon et al., 2006), the EMEP database (Mantseva et al., 2004), a dataset for POPCYCLING-Baltic project (Pacyna et al., 2003), and a global emission inventory for the year 2008. The SMOKE for Europe (SMOKE-EU) model (Bieser et al., 2011) output used in this study follows a methodology that is based on the TNO database and has been evaluated in a number of subsequent CMAQ applications (Aulinger et al., 2011; Bewersdorff et al., 2009; Matthias et al., 2008). In order to spatially disaggregate BaP emissions in a more realistic way, SMOKE-EU improves on using a linear dependency on the population density as surrogate factor by introducing different relevant relationships based on source classification and meteorological parameters. For example, the relationship to wood availability and the level of urban agglomeration have been introduced in an effort to include the effect of burning biomass. The resulting spatial distribution and annual average emission flux of BaP for our domain during the year 2006 is shown in Fig. 1. In addition, BaP emissions are shaped temporally using diurnal and weekly functions. As residential combustion is associated with the majority of BaP emissions, a strong dependence on the season and spatial variability is expected. To account for this dependence, heating




Figure 2. Annual average surface-level BaP concentrations [$\log c_{\text{BaP}} (\text{ng m}^{-3})$] during 2006 plotted in logarithmic scale for (a) the Junge–Pankow (JP) scenario 1 and (b) for the fully expanded GPP scenario 5, which includes a heterogeneous reaction with ozone.

degree days have been introduced as a proxy (Aulinger et al., 2011). A major strength of the SMOKE-EU inventory is the provision of CMAQ-ready emissions for all species involved in the desired chemical mechanism (e.g. CB5) and aerosol module (e.g. aero4). Following the trimodal representation of aerosol in CMAQ, 0.1 % of the emissions of BaP was allocated to the Aitken mode, while the rest was emitted in the accumulation mode, as this is the default treatment with all primary organic aerosol species in the model.

3.2 Evaluation data and sensitivity simulations

The spatio-temporal coverage of BaP in air and deposition in Europe and Russia is limited, in many regions sporadic or not available at all. For 2006, concentration in air is available from eight stations ($n = 173$), while deposition information is available from six stations of the EMEP monitoring network (Aas and Breivik, 2012). An overview of the data, including site codes, geographical information, and the sampling strategies, is provided in Table S2. It is evident that during 2006, only 4 of the total 10 sites provide information on concentration and deposition simultaneously. Additionally, the sampling strategy varies widely among the stations, leading to a non-homogeneous pool of data. For example, at the Košetice (CZ0003R) and Niembro (ES0008R) stations, BaP is measured for 24 h once a week, while in Aspveten (SE0012R) and in Pallas (FI0036R) samples cover one week and are taken once every month. On the other hand, the Latvian sites (LV0010R, LV0016R) are sampled on a monthly basis, while for the High Muffles location (GB0014R) a 3-month sample is obtained four times in 2006. Finally, the

sites located in Germany (DE0001R, DE0009R) provided information only in terms of wet deposition and on a monthly basis during 2006.

Based on the set of GPP mechanisms described before, the effect of each parameterization was evaluated with the EMEP measurements by incrementally testing the models involved, while keeping the same meteorological and emission inputs. Table 2 summarizes the five annual scenarios that were developed for 2006: (1) Junge–Pankow, (2) scenario 1 with water dissolution, (3) scenario 2 with the Harner-Bidleman (HB) K_{oa} absorption model, (4) scenario 2 with dual OM absorption and black carbon adsorption model Dachs-Eisenreich (DE) scheme, and (5) scenario 4 supplemented by a heterogeneous reactivity module. A spin-up time of 7 days was allowed for each simulation in order to maintain a low influence of the initial conditions. Comparisons with measurements of BaP in ambient air were produced using the statistical computing framework of the R language, and the model performance was evaluated, building on a wide range of statistics calculated using the Models-3 and openair packages (Carslaw and Ropkins, 2012).

4 Results and discussion

4.1 Spatio-temporal distributions of BaP over Europe

The pattern of normalized annual mean BaP concentrations over the domain using the baseline scenario 1 (JP) is presented in Fig. 2a. Elevated levels were found in regions with strong emission sources in central Europe (Po Valley, Rhine-Ruhr, Poland, and Ukraine areas), as well as in the vicinity

Table 2. Summary of the simulated scenarios designed to test gas-particle partitioning and heterogeneous reactivity of BaP.

Description and references	Scenario				
	1	2	3	4 ⁵	5
Gas-phase reaction with OH	Yes	Yes	Yes	Yes	Yes
Adsorption model (JP) ¹	JP	JP	JP	JP	JP
Aerosol water dissolution (W)	–	W	W	W	W
Absorption to OM (HB) ²	–	–	HB	–	–
Absorption to EC/BC (DE) ³	–	–	–	DE	DE
Heterogeneous reaction with O ₃ (KW) ⁴	–	–	–	–	KW

¹ Junge (1977) and Pankow (1987). ² Harner and Bidleman (1998). ³ Dachs and Eisenreich (2000).
⁴ Kwamena et al. (2004). ⁵ Additional simulations with temperature-dependent octanol–air partition coefficient calculated based on Odabasi et al. (2006) and a thermodynamic estimation model suggested by van Noort (2003).

of large cities (Moscow, Paris, Vienna, Madrid, and Istanbul, among others). In addition, annual mean BaP concentrations according to the fully expanded scenario 5 (KW) are depicted in Fig. 2b. Comparing the output from the aforementioned simulations revealed higher BaP levels with scenario 5, which can be associated with the geographic distribution of emissions. This is the first indication that the chosen partitioning approach has a significant impact on the model results for total BaP concentrations. However, it is also evident that in southern Europe and the Mediterranean region, the BaP losses associated with the addition of an upper estimate of heterogeneous reactivity with O₃ (scenario 5, Fig. 2b) may have a limiting effect on BaP long-range transport (LRT) potential. This can be supported by the prominent BaP gradients calculated over the Mediterranean Sea, acting as a sink that is driven by low BaP emissions and elevated O₃ levels. The overall model patterns show a good agreement with recent regression methods mapping BaP levels that provide with a partial coverage of our EU domain. With respect to previous studies involving simulations of BaP over Europe, we find lower BaP concentrations, but within the same order of magnitude as those estimated by initial CMAQ applications (Aulinger et al., 2007; Bewersdorff et al., 2009; Matthias et al., 2009). This supports the perception that BaP emissions over Europe have decreased substantially during the recent decades, a feature improved in subsequent evolution of the emissions model which led to better agreement with the more recent CMAQ applications (Bieser et al., 2012). A closer look over Italy and the Iberian Peninsula reveals a good agreement with previous studies depicting BaP distributions for these regions (San José et al., 2013; Silibello et al., 2012). Finally, discrepancies over eastern Europe indicate that a more elaborate emissions inventory (i.e. domestic heating adjustments, fire activities) would be beneficial to address the elevated BaP levels predicted by recent hybrid modelling approaches (Guerreiro et al., 2016a).

4.1.1 Effect of GPP parameterizations and heterogeneous reactivity with O₃

Figure 3 and Table 3 illustrate the differences resulting from the incrementally tested scenarios (Table 2), expressed as annual mean BaP concentrations over the entire domain of interest and the individual 28 EU member states. As expected, the effect of dissolution into aerosol water (Fig. 3a) is limited throughout most of the domain, with coastal urban areas being the most influenced (i.e. Istanbul, North Sea). According to scenario 2, BaP dissolution in aerosol liquid water led to a reduction of less than 0.36 % of ground-level BaP in Europe compared to the baseline JP scheme (Table 3). Introducing the HB absorption scheme (scenario 3) resulted in ~ 119.2 % higher average BaP concentrations (Fig. 3b) with extremes observed in areas associated with emission sources in central Europe (i.e. Po Valley) and large cities (i.e. Moscow, Paris, Vienna, and Istanbul). Expressed in terms of domain-wide atmospheric lifetimes (defined as $\tau = b_{\text{tot}}/F_{\text{em}}$, where b_{tot} denotes the total atmospheric burden and F_{em} the total emissions), scenario 3 leads to a 210 % increase for BaP over scenario 2. Similarly, shifting to the DE scheme (scenario 4) appears to have an additive effect on near-ground BaP concentrations with total lifetimes reaching a domain-average increase of 13 %. Compared to scenario 3, scenario 4 affects a larger area of the domain (Fig. 3c), even though the magnitude of the effect is less pronounced due to its association with the availability of black carbon. The domain mean of modelled elemental carbon is $0.0212 \mu\text{g m}^{-3}$ and the spatial distribution matches its emission pattern (Fig. S3). These levels are significantly lower than in the field studies in the urban air of south-western Europe (EEA, 2013; Milford et al., 2016) and may subsequently lead to a hindered performance of the DE scheme.

Finally, the introduction of heterogeneous chemistry (scenario 5) had, as expected, a substantially negative effect on the BaP concentrations within our domain, culminating in major conurbations (i.e. Po Valley, Moscow, and Istanbul;




Figure 3. Differences in annual average surface-level BaP concentrations (ng m^{-3}) during 2006 plotted for (a) water dissolution effect: scenario 1 to scenario 2, (b) K_{oa} effect: scenario 3 to scenario 2, (c) dual model effect: scenario 4 to scenario 3, and (d) O_3 degradation effect: scenario 4 to scenario 5.

see Fig. 3d). Heterogeneous degradation due to the reaction with O_3 accounts for an approximate 18.8 % reduction of the mean ground-level BaP and a 25 % reduction of its atmospheric lifetime in the European domain (Table 3). As mentioned in Sect. 4.1, the BaP concentration gradients reveal a strong spatial link with emission sources in the south, as well as high O_3 concentrations typically found along the coasts of the Mediterranean Sea (Fig. S4). CMAQ calculates ozone patterns that are in accordance to the findings of recent analyses, exhibiting a similar performance to measurements at the European level (de Smet et al., 2009).

4.1.2 Seasonality and relevant parameters affecting BaP fate and transport

Seasonally disaggregated distributions of BaP, as simulated in scenario 4, are presented in Fig. 4. Strong emissions during wintertime and autumn have an apparent effect on the average ground-level concentrations. In addition, a normalized effect of long-range transport for BaP can be observed in the vicinity of Istanbul and other coastal metropolitan areas, over parts of the large water bodies of the Mediterranean and the Black Sea. This scenario, however, does not include the treatment of the heterogeneous reaction with ozone and, therefore, provides an upper-bound estimate of the LRT potential

Table 3. Average ground-level BaP concentrations, c_{BaP} (pg m^{-3}) over the entire domain and in individual 28 EU member states for all simulated scenarios and concentration changes Δc_{BaP} (%) between subsequent scenarios.

	1		2		3		4		5	
	c_{BaP} Conc. pg m^{-3}	c_{BaP} pg m^{-3}	Δc_{BaP} %							
Domain average	10.4	10.4	−0.4	22.8	119.2	28.3	24.1	23.0	−18.8	
Austria	58.4	58.4	0.0	89.0	52.4	103.4	16.1	86.5	−16.3	
Belgium	36.2	36.1	−0.1	65.8	82.1	75.8	15.2	62.5	−17.6	
Bulgaria	14.7	14.7	−0.1	28.9	97.1	37.7	30.4	26.7	−29.3	
Croatia	15.2	15.2	−0.1	35.1	131.0	45.5	29.9	32.2	−29.4	
Cyprus	11.2	11.2	0.0	13.2	17.8	24.7	87.2	10.1	−59.1	
Czech Republic	38.6	38.6	0.0	70.1	81.5	83.5	19.1	67.9	−18.7	
Denmark	7.3	7.2	−1.3	22.4	213.0	26.0	16.0	24.5	−5.7	
Estonia	9.5	9.5	−0.9	28.9	205.3	34.6	19.7	32.0	−7.4	
Finland	6.0	5.8	−2.9	16.1	176.0	21.9	36.2	17.7	−19.5	
France	27.1	27.1	0.0	48.0	76.9	56.6	18.0	44.7	−21.0	
Germany	30.3	30.3	−0.1	57.6	90.2	69.2	20.2	55.6	−19.6	
Greece	7.4	7.4	−0.1	14.7	97.7	22.3	52.2	12.4	−44.6	
Hungary	29.3	29.3	−0.1	61.3	109.5	75.1	22.5	59.7	−20.5	
Ireland	3.8	3.8	−0.1	6.8	80.0	8.7	27.5	6.0	−31.7	
Italy	36.9	36.9	0.0	62.8	70.1	76.3	21.6	57.9	−24.1	
Lithuania	10.0	10.0	−0.5	32.7	228.4	38.4	17.5	36.3	−5.5	
Luxembourg	37.1	37.1	0.0	62.0	67.2	71.7	15.8	57.9	−19.3	
Latvia	6.0	5.9	−1.3	25.7	335.0	30.2	17.7	29.8	−1.4	
Malta	0.5	0.4	−0.5	0.5	20.6	0.7	37.1	0.5	−36.6	
Netherlands	24.0	23.9	−0.3	52.5	119.5	63.1	20.2	51.5	−18.4	
Poland	25.6	25.6	−0.2	56.7	121.7	67.4	18.8	57.4	−14.8	
Portugal	15.0	15.0	0.0	20.9	39.1	28.4	36.3	18.1	−36.5	
Romania	12.7	12.7	0.0	27.4	115.8	36.0	31.4	25.7	−28.6	
Spain	8.4	8.4	0.0	11.9	41.8	16.8	41.4	9.3	−44.8	
Sweden	3.7	3.6	−2.4	11.3	211.7	14.3	26.7	13.2	−7.7	
Slovakia	27.0	27.0	−0.1	53.5	98.1	65.2	21.9	52.5	−19.5	
Slovenia	28.5	28.5	0.0	60.3	111.3	71.3	18.3	57.0	−20.1	
United Kingdom	3.3	3.3	−0.4	8.9	170.6	11.4	28.4	8.2	−27.6	

of BaP. On the other hand, Fig. 5 demonstrates the seasonal effect of the heterogeneous reaction of BaP with ground-level ozone on BaP concentrations. Distributions of BaP concentration differences reveal a stronger reduction during winter that is influenced by central and southern European emission sources and midlatitude ozone availability. Interestingly, the differences between winter and summertime appear to be comparable in terms of magnitude, despite the substantially weaker sources during summer that shift the process to the south.

An important factor for the performance of the BaP model is the mass concentration and composition of the aerosols in the three different modes and their relationship with the temperature of the ambient air. Figure 6 illustrates the aerosol distributions as simulated at model cells representing three different settings (urban: the city of Vienna, suburban: CZ0003R, remote: FI0036R) during a typical winter and summer day. As we move further away from the vicinity of the sources, we notice not only a decrease of the over-

all BaP levels but also a change in the distribution influenced by each individual simulated scenario. While the basic Junge–Pankow implementation results in higher coarse-mode BaP concentrations, the HB and DE schemes accentuate the strength of the accumulation mode where most of the BaP aerosol mass is emitted. The disproportionately higher coarse-BaP fractions calculated at the remote grid cells during the summertime can be explained by conditions favouring long-range transport.

Regarding the parameterizations of the HB and DE schemes, the calculation of the parameter K_{oa} has a direct effect on the relevant model results (scenarios 3 and 4). Use of the regression parameters for the semi-logarithmic temperature-dependent form proposed by Odabasi et al., (2006) results in a mixed effect over its predecessor scheme, which is seasonally disaggregated and presented in Fig. S5. These figures suggest that the partitioning based on each K_{oa} approach is sufficiently similar for most of the domain during the winter. However, differences between




Figure 4. Seasonally disaggregated average surface-level BaP concentration [$\log c_{\text{BaP}}$ (ng m^{-3})] maps predicted by scenario 4 during 2006.

the two expressions with respect to simulating summertime BaP concentrations over central Europe were noted. In addition, simulations that address the uncertainties contributed by the method for estimating K_{SA} were performed for the winter period of 2006 using the DE scheme. Figure S6 reveals similar results, with BaP levels slightly higher in cells with strong emissions sources. However, with the exception of Moscow, the average difference BaP concentration in urban cells did not exceed 0.02 ng m^{-3} when K_{SA} was calculated based on the thermodynamic estimation model suggested by van Noort (2003). Similarly to K_{oa} , results suggest that the partitioning based on each K_{SA} approach is sufficiently similar.

4.2 Model evaluation with EMEP measurements

The EMEP network database and model output from all five scenarios were imported in the R language framework to create measurement–model pairs for further analysis. Figure 7 illustrates the resulting time series plots for selected EMEP sites against CMAQ output extracted from scenario 4. This figure also demonstrates differences in terms of sampling protocols among individual sites, with the extreme of the High Muffles station ($4 \text{ samples year}^{-1}$ – Table S2). Aggregate temporal profiles of the modelled and measured BaP


concentrations across all sites depicted in Fig. 8 reveal a strong seasonal pattern that is captured well by the model. However, BaP concentrations appear to be underestimated during the colder months. This effect can be attributed to the strength of emissions, which is probably related to inadequate coverage of residential heating in current inventories (wood/biofuel burning). Figure S7 illustrates this by comparing annual average BaP emission fluxes between model cells enclosing EMEP monitoring site locations and selected metropolitan areas of Europe during 2006. While the deficiency of PAH inventories is known, it may limit the usefulness of background measurements (such as those in the EMEP network) in assessing the overall model performance under different GPP scenarios.

It is evident from the metrics obtained across all sites (Table 4 and Taylor diagram summary Fig. S8) that the increased complexity in GPP formulation results in better agreement with the EMEP measurements. For reasons mentioned in Sects. 2.2–2.3 and 4.1.1, deviations from this observation are the simulated scenarios that introduce the dissolution to aerosol water (scenario 2) and degradation due to the heterogeneous reaction with ozone (scenario 5). Overall, the dual model (DE) employed in scenario 4 showed the best agreement with the measurements. The DE parameterization also performed best when compared with the JP and HB schemes

Table 4. Comparison between modelled and measured BaP concentrations for the entire 2006 EMEP dataset ($n = 173$) along with performance metrics and ranking across all sites.

Rank	Scenario	FAC2	MB	MGE	NMB	NMGE	RMSE	r	COE	IOA
1	4	0.29480	-0.05622	0.07145	-0.65264	0.82950	0.14729	0.35193	0.22443	0.61222
2	3	0.25434	-0.06279	0.07280	-0.72889	0.84516	0.15081	0.33423	0.20979	0.60490
3	5	0.21387	-0.07021	0.07345	-0.81511	0.85272	0.15179	0.47598	0.20273	0.60136
4	1	0.18497	-0.07640	0.07887	-0.88699	0.91559	0.16069	0.31088	0.14395	0.57197
5	2	0.17919	-0.07649	0.07894	-0.88798	0.91642	0.16076	0.30793	0.14317	0.57158

FAC: fraction of predictions within a factor of two, MB: mean bias, MGE: mean gross error, NMB: normalized mean bias, NMGE: normalized mean gross error, RMSE: root mean squared error, r : correlation coefficient, COE: coefficient of efficiency, IOA: index of agreement.

**Figure 5.** Average difference of surface-level BaP concentrations (ng m^{-3}) between scenario 4 and the fully expanded GPP scenario 5, which includes the heterogeneous reaction with ozone during the winter and summer months of 2006 (scenario 4–scenario 5).

on global-scale applications (Lammel et al., 2009). However, when looking at site-specific performance metrics (Table S9), the Košetice location (CZ0003R) reveals a better agreement with the fully expanded scenario 5, which treats

degradation by ozone. The Košetice site is the closest to conurbations among the EMEP sites available during 2006. This could indicate that the heterogeneous chemistry, which is implemented as an upper estimate in this model, tends to underestimate BaP levels further away from the sources. The best model performance was observed at the Pallas station (FI0036R), where scenario 4 was capable of matching 50 % of the measurements within a factor of 2. For the remaining remote sites in Finland, the simulation under scenario 5 calculated underestimated BaP levels. Based on the index of agreement (IOA) metric, a ranking of the overall model performance similar to the Pallas site was observed for the rest of the monitoring sites. Having compared site-specific performance to previous modelling efforts over Europe (Aulinger et al., 2007; Matthias et al., 2009), we reach the same conclusion as pointed out with the spatial distributions; i.e. we attribute the lower-performance metrics presented here to the substantially weaker emission sources in recent years, hence the better agreement with the more recent SMOKE-EU/CMAQ application (Bieser et al., 2012). With respect to other annual simulations, we observed lower average BaP levels over the Piedmont region compared to the FARM model, which were higher than the EMEP (MSC-E) output used in the same study (Silibello et al., 2012). Finally, a qualitative analysis of the deposition measurements for the few stations that measured BaP showed that CMAQ calculated plausible patterns of dry and wet deposition (see Fig. S10).

5 Conclusions

This study presented the development of a new modelling system for investigating the dynamics of transport and gas-particle partitioning schemes for POPs at the regional scale over Europe. The implementation of this model is based on the WRF-CMAQ model framework with additional algorithms for GPP schemes and a module that accounts for heterogeneous reactivity of BaP with ozone. Predictions from the WRF-CMAQ-BaP model were compared to measurements obtained from the EMEP monitoring network. Model evaluation has revealed satisfactory agreement to the mea-




Figure 6. Distribution of BaP mass [$\log c_{\text{BaP}}$ (ng m^{-3})] in aerosols of three different modes in three different cells [urban (Vienna), suburban (CZ0003R), and remote (FI0036R)] for each scenario as calculated during a winter (20 January) and a summer (20 July) day.

measurements and performance metrics similar to those of previous studies with significantly higher measurement availability. The results presented in this work suggest that the new expanded model is able to simulate the ambient levels of BaP fairly well. It is found that BaP distributions in Europe are very sensitive to the choice of GPP models. While this statement has been also indicated at the global scale (Lammel et al., 2009), GPP has not been adequately explored by previous regional-scale studies (see Table S1 and references therein). We conclude that the dual OM absorption and black carbon adsorption (DE) parameterization (scenario 4–5) offer a better performance, as BaP predictions tend to be closer to the measurements. Introducing an independent upper estimate of heterogeneous BaP reactivity is found to significantly reduce BaP levels throughout the domain, with a re-

duction pattern that follows the spatial distribution of ozone and emission sources. Despite the limited BaP monitoring network, a better agreement was observed when complementing WRF-CMAQ-BaP with the upper estimate for heterogeneous reactivity (scenario 5) at certain EMEP sites. Uncertainties and limitations of the current emission inventory estimates of BaP, OM, and BC, along with insufficient laboratory data, are of major concern, hampering efforts to fully evaluate the long-range transport potential, the effect of photochemistry, and interactions of such compounds. In addition, certain simplifications in the treatment of coarse PM and OM within the aerosol module of CMAQ (Binkowski, 2003; Mathur et al., 2008) are posing limitations on fully exploiting the advanced empirical gas-particle partitioning models (K_{oa} , Dachs-Eisenreich) in this specific mode. An implicit




Figure 7. Model predicted and observed concentration (ng m^{-3}) time series at selected EMEP monitoring sites during 2006 for the dual GPP model scenario 4.

overlap of the empiric adsorption model (JP) with the K_{oa} (or another absorption) model (see Sect. 2.1.1) may artificially enhance partitioning, something that has not been addressed by previous modelling efforts. However, such discrepancy is expected to be negligible for BaP, especially when compared to existing uncertainties of output aerosol parameters propagating from emission estimates and from the current design and state of development of the CMAQ aerosol module (AERO).

The modelling approach presented here allows a simultaneous estimation of organic pollutants, including semivolatiles (such as most POPs) within the CMAQ framework, and can be used as a supplementary component of a population exposure modelling system. Although similar systems are currently explored, they rely heavily on observation data while using a very basic air quality model structure (Guerreiro et al., 2016a, b). At this time, annual exposure estimates are more useful due to the low accuracy accompanying highly time-resolved models and observations. As previ-

ous efforts to evaluate the use of mesoscale models for BaP dispersion studies also conclude, changes in emission patterns and their strengths are of particular importance. While there is undeniable effort to shift from fossil fuels to renewables which has led to updated emission inventories, domestic heating across Europe has also a complex relationship to economic factors and human activities that varies tremendously between countries and fuel types (Lohmann et al., 2006; Saffari et al., 2013). Besides the need for updated emission inventories for the eastern Europe, extended simulations are suggested for years with higher measurement availability and density. Following modelling, studies should focus on quantifying the long-range transport potential and examining the hypothesis that secondary organic aerosol (SOA) may facilitate PAH transport by utilizing more sophisticated aerosol and heterogeneous chemistry parameterizations or submodels (i.e. accounting also for the burial effect; see Sect. 2.2). More specifically, recent evidence suggests that, in cold and dry air, accessibility of PAHs in OM is reduced (due to low




Figure 8. Model predicted and observed monthly BaP concentrations (ng m^{-3}) across all EMEP sites in 2006 for the dual GPP model scenario 4.

diffusivity), which might explain the apparent inconsistency of high LRT potential (BaP levels in the Arctic) on one hand and the relative high heterogeneous reactivity measured in the laboratory on the other (Friedman et al., 2014; Zelenyuk et al., 2012; Zhou et al., 2012, 2013). To understand the atmospheric lifetime of PAH, interaction between PAH and SOA, and progression of the chemical composition of ambient OM, a new need for additional studies quantifying GPP and LRT potential under a wide range of atmospheric conditions is emerging. In view of such findings, the WRF-CMAQ-BaP modelling system should be extended to study a wide range of additional organic pollutants and processes (e.g. multicompartamental cycling, biodegradation, heterogeneous chemistry).

6 Data availability

The WRF-CMAQ-BaP output data used in this study are archived in MetaCentrum and are available upon request from the authors (efstathiou@recetox.muni.cz).

The Supplement related to this article is available online at doi:10.5194/acp-16-15327-2016-supplement.

Acknowledgements. The authors thank Frank Binkowski, Sarav Arunachalam, and Zac Adelman at the University of North Carolina (UNC) for their guidance and support during the onset of this project. EMEP is acknowledged for the monitoring data. Access to computing and storage facilities owned by parties and projects contributing to the National Grid Infrastructure MetaCentrum provided under the programme “Projects of Large Infrastructure for Research, Development, and Innovations” (LM2010005) is also greatly appreciated. This research was supported by the Granting Agency of the Czech Republic (project no. 312334), the National Sustainability Programme of the Czech Ministry of Education, Youth and Sports (LO1214), and the RECETOX research infrastructure (LM2015051).

Edited by: I. Riipinen

Reviewed by: three anonymous referees

References

- Aas, W. and Breivik, K.: Heavy metals and POP measurements, 2010, EMEP/CCC, Kjeller, 2012.
- Andersson, J. T. and Achten, C.: Time to say goodbye to the 16 EPA PAHs? Toward an up-to-date use of PACs for environmental purposes, *Polycycl. Aromat. Comp.*, 35, 330–354, doi:10.1080/10406638.2014.991042, 2015.

- Atkinson, R. and Arey, J.: Atmospheric chemistry of gas-phase polycyclic aromatic hydrocarbons: formation of atmospheric mutagens, *Environ. Health Persp.*, 102, 117–126, 1994.
- Aulinger, A., Matthias, V., and Quante, M.: Introducing a partitioning mechanism for PAHs into the Community Multiscale Air Quality Modeling System and its application to simulating the transport of benzo(a)pyrene over Europe, *J. Appl. Meteor. Climatol.*, 46, 1718–1730, doi:10.1175/2007JAMC1395.1, 2007.
- Aulinger, A., Matthias, V., and Quante, M.: An approach to temporally disaggregate benzo(a)pyrene emissions and their application to a 3D Eulerian atmospheric chemistry transport model, *Water Air Soil Pollut.*, 216, 1–13, doi:10.1007/s11270-010-0559-x, 2011.
- Bärning, H., Bucheli, T. D., Broman, D., and Gustafsson, Ö.: Soot–water distribution coefficients for polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans and polybrominated diphenylethers determined with the soot cosolvency-column method, *Chemosphere*, 49, 515–523, doi:10.1016/s0045-6535(02)00389-2, 2002.
- Bewersdorff, I., Aulinger, A., Matthias, V., and Quante, M.: The effect of temporal resolution of PAH emission data on transport and deposition patterns simulated with the Community Multiscale Air Quality modelling system (CMAQ), *Meteorol. Z.*, 18, 41–53, doi:10.1127/0941-2948/2009/351, 2009.
- Beyer, A., Mackay, D., Matthies, M., Wania, F., and Webster, E.: Assessing long-range transport potential of persistent organic pollutants, *Environ. Sci. Technol.*, 34, 699–703, doi:10.1021/es990207w, 2000.
- Bieser, J., Aulinger, A., Matthias, V., Quante, M., and Bultjes, P.: SMOKE for Europe – adaptation, modification and evaluation of a comprehensive emission model for Europe, *Geosci. Model. Dev.*, 4, 47–68, doi:10.5194/gmd-4-47-2011, 2011.
- Bieser, J., Aulinger, A., Matthias, V., and Quante, M.: Impact of emission reductions between 1980 and 2020 on atmospheric benzo[a]pyrene concentrations over Europe, *Water Air Soil Pollut.*, 223, 1393–1414, doi:10.1007/s11270-011-0953-z, 2012.
- Binkowski, F. S.: Models-3 Community Multiscale Air Quality (CMAQ) model aerosol component, 1. Model description, *J. Geophys. Res.*, 108, 4183, doi:10.1029/2001JD001409, 2003.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol. Sci. Tech.*, 40, 27–67, doi:10.1080/02786820500421521, 2006.
- Byun, D. and Schere, K. L.: Review of the governing equations, computational algorithms, and other components of the Models-3 Community Multiscale Air Quality (CMAQ) modeling system, *Appl. Mech. Rev.*, 59, 51–77, 2006.
- Byun, D. W.: Dynamically consistent formulations in meteorological and air quality models for multiscale atmospheric studies, Part II: Mass conservation issues, *J. Atmos. Sci.*, 56, 3789–3807, doi:10.1175/1520-0469(1999)056<3789:DCFIMA>2.0.CO;2, 1999a.
- Byun, D. W.: Dynamically consistent formulations in meteorological and air quality models for multiscale atmospheric studies, Part II: Mass conservation issues, *J. Atmos. Sci.*, 56, 3808–3820, doi:10.1175/1520-0469(1999)056<3808:DCFIMA>2.0.CO;2, 1999b.
- Byun, D. W. and Ching, J.: Science algorithms of the EPA Models-3 community multiscale air quality (CMAQ) modeling system, US Environmental Protection Agency, Office of Research and Development Washington, DC, USA, Washington, DC 20460., 1999.
- Carslaw, D. C. and Ropkins, K.: openair – An R package for air quality data analysis, *Environ. Model. Softw.*, 27/28, 52–61, doi:10.1016/j.envsoft.2011.09.008, 2012.
- Chang, J., Middleton, P., Stockwell, W., Pleim, J. E., Lansford, H., Binkowski, F. S., Seaman, N., Stauder, D., and Madronic, S.: The regional acid deposition model and engineering model, in: *Acid Deposition State of Science and Technology*, 1986.
- Cooter, E. J. and Hutzell, W. T.: A regional atmospheric fate and transport model for atrazine, 1. Development and implementation, *Environ. Sci. Technol.*, 36, 4091–4098, doi:10.1021/es011371y, 2002.
- Dachs, J. and Eisenreich, S. J.: Adsorption onto aerosol soot carbon dominates gas-particle partitioning of polycyclic aromatic hydrocarbons, *Environ. Sci. Technol.*, 34, 3690–3697, doi:10.1021/es991201, 2000.
- Dachs, J., Ribes, S., van Drooge, B., Grimalt, J., Eisenreich, S. J., and Gustafsson, Ö.: Response to the comment on “Influence of soot carbon on the soil-air partitioning of polycyclic aromatic hydrocarbons”, *Environ. Sci. Technol.*, 38, 1624–1625, doi:10.1021/es0307118, 2004.
- de Maagd, P., Hulscher, Ten, D., Van den Heuvel, H., Opperhuizen, A., and Sijm, D.: Physicochemical properties of polycyclic aromatic hydrocarbons: Aqueous solubilities, n-octanol/water partition coefficients, and Henry’s law constants, *Environ. Toxicol. Chem.*, 17, 251–257, 1998.
- Denier van der Gon, H. A. C., Visschedijk, A. J. H., and Bolscher, M. V. H.: Study to the effectiveness of the UNECE Persistent Organic Pollutants (POP), Protocol and cost of additional measures, Phase II: Estimated emission reduction and cost of options for a possible revision of the POP Protocol, TNO, 2006.
- de Smet, P., Horálek, J., Coňková, M., Kurfürst, P., de Leeuw, F., and Denby, B.: European air quality maps of ozone and PM₁₀ for 2006 and their uncertainty analysis, ETAC/ACC Technical Paper 2008/8, 2009.
- DETR: Polycyclic Aromatic Hydrocarbons, Department of the Environment, Transport and the Regions Expert Panel on Air Quality Standards, Stationery Office, London, 2009.
- Diamond, M. L. and Hodge, E.: Urban Contaminant Dynamics: From Source to Effect, *Environ. Sci. Technol.*, 41, 3796–3800, doi:10.1021/es072542n, 2007.
- EC: Directive 2004/107/EC of the European Parliament and of the Council, Official Journal of the European Union, 2004.
- EEA: Status of black carbon monitoring in ambient air in Europe, European Environment Agency, European Environment Agency, 2013.
- Fernandez, P., Grimalt, J. O., and Vilanova, R. M.: Atmospheric gas-particle partitioning of polycyclic aromatic hydrocarbons in high mountain regions of Europe, *Environ. Sci. Technol.*, 36, 1162–1168, doi:10.1021/es010190t, 2002.
- Finizio, A., Mackay, D., Bidleman, T., and Harner, T.: Octanol-air partition coefficient as a predictor of partitioning of semi-volatile organic chemicals to aerosols, *Atmos. Environ.*, 31, 2289–2296, doi:10.1016/S1352-2310(97)00013-7, 1997.
- Finlayson-Pitts, B. J. and Pitts Jr., J. N.: *Chemistry of the Upper and Lower Atmosphere*, Academic Press, 1999.
- Friedman, C. L. and Selin, N. E.: Long-range atmospheric transport of polycyclic aromatic hydrocarbons: a global 3-D model analy-

- sis including evaluation of arctic sources, *Environ. Sci. Technol.*, 46, 9501–9510, doi:10.1021/es301904d, 2012.
- Friedman, C. L., Pierce, J. R., and Selin, N. E.: Assessing the influence of secondary organic versus primary carbonaceous aerosols on long-range atmospheric polycyclic aromatic hydrocarbon transport, *Environ. Sci. Technol.*, 48, 3293–3302, doi:10.1021/es405219r, 2014.
- Galarneau, E., Makar, P. A., Zheng, Q., Narayan, J., Zhang, J., Moran, M. D., Bari, M. A., Pathela, S., Chen, A., and Chlumsky, R.: PAH concentrations simulated with the AURAMS-PAH chemical transport model over Canada and the USA, *Atmos. Chem. Phys.*, 14, 4065–4077, doi:10.5194/acp-14-4065-2014, 2014.
- Goss, K.-U.: Comment on “Influence of soot carbon on the soil-air partitioning of polycyclic aromatic hydrocarbon”, *Environ. Sci. Technol.*, 38, 1622–1623, doi:10.1021/es0301370, 2004.
- Götz, C. W., Scheringer, M., MacLeod, M., Roth, C. M., and Hungerbühler, K.: Alternative approaches for modeling gas-particle partitioning of semivolatile organic chemicals: model development and comparison, *Environ. Sci. Technol.*, 41, 1272–1278, doi:10.1021/es060583y, 2007.
- Grell, G. A.: Prognostic Evaluation of Assumptions Used by Cumulus Parameterizations, *Mon. Weather Rev.*, 121, 764–787, doi:10.1175/1520-0493(1993)121<0764:PEOAUB>2.0.CO;2, 1993.
- Grell, G. A. and Devenyi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation techniques, *Geophys. Res. Lett.*, 29, 1693, doi:10.1029/2002GL015311, 2002.
- Guerreiro, C., Horálek, J., and De Leeuw, F.: Benzo(a)pyrene in Europe: Ambient air concentrations, population exposure and health effects, *Environ. Pollut.*, 214, 657–667, doi:10.1016/j.envpol.2016.04.081, 2016a.
- Guerreiro, C., Horálek, J., de Leeuw, F., and Couvidat, F.: Mapping ambient concentrations of benzo(a)pyrene in Europe, European Topic Centre on Air Pollution and Climate Change Mitigation, Technical Paper 2014/6, 2016b.
- Gusev, A., Mantseva, E., and Shatalov, V.: Regional multicompartiment model MSCE-POP, MSC-E Technical Report, 2005.
- Halsall, C. J., Sweetman, A. J., Barrie, L. A., and Jones, K. C.: Modelling the behaviour of PAHs during atmospheric transport from the UK to the Arctic, *Atmos. Environ.*, 35, 255–267, doi:10.1016/S1352-2310(00)00195-3, 2001.
- Hansen, K. M., Halsall, C. J., and Christensen, J. H.: A dynamic model to study the exchange of gas-phase persistent organic pollutants between air and a seasonal snowpack, *Environ. Sci. Technol.*, 40, 2644–2652, doi:10.1021/es051685b, 2006.
- Harner, T. and Bidleman, T. F.: Octanol-air partition coefficient for describing particle/gas partitioning of aromatic compounds in urban air, *Environ. Sci. Technol.*, 32, 1494–1502, doi:10.1021/es970890r, 1998.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.-Atmos.*, 113, D13103, doi:10.1029/2008JD009944, 2008.
- Inomata, Y., Kajino, M., Sato, K., Ohara, T., Kurokawa, J.-I., Ueda, H., Tang, N., Hayakawa, K., Ohizumi, T., and Akimoto, H.: Emission and atmospheric transport of particulate PAHs in Northeast Asia, *Environ. Sci. Technol.*, 46, 4941–4949, doi:10.1021/es300391w, 2012.
- Jonker, M. T. O. and Koelmans, A. A.: Sorption of polycyclic aromatic hydrocarbons and polychlorinated biphenyls to soot and soot-like materials in the aqueous environment: Mechanistic considerations, *Environ. Sci. Technol.*, 36, 3725–3734, doi:10.1021/es020019x, 2002.
- Junge, C. E.: Fate of pollutants in the air and water environment, edited by: Suffet, I. H., Wiley, New York, 1977.
- Kahan, T. F. and Donaldson, D. J.: Heterogeneous ozonation kinetics of polycyclic aromatic hydrocarbons on organic films, *Atmos. Environ.*, 40, 3448–3459, doi:10.1016/j.atmosenv.2006.02.004, 2006.
- Kaiser, J. C., Riemer, N., and Knopf, D. A.: Detailed heterogeneous oxidation of soot surfaces in a particle-resolved aerosol model, *Atmos. Chem. Phys.*, 11, 4505–4520, doi:10.5194/acp-11-4505-2011, 2011.
- Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, doi:10.1126/science.1092185, 2004.
- Keyte, I. J., Harrison, R. M., and Lammel, G.: Chemical reactivity and long-range transport potential of polycyclic aromatic hydrocarbons – a review, *Chem. Soc. Rev.*, 42, 9333–9391, doi:10.1039/C3CS60147A, 2013.
- Kim, Y. P., Seinfeld, J. H., and Saxena, P.: Atmospheric gas-aerosol equilibrium, I. Thermodynamic model, *Aerosol Sci. Tech.*, 19, 157–181, doi:10.1080/02786829308959628, 1993.
- Kwamena, N.-O. A., Thornton, J. A., and Abbatt, J. P. D.: Kinetics of surface-bound benzo[a]pyrene and ozone on solid organic and salt aerosols, *J. Phys. Chem. A*, 108, 11626–11634, doi:10.1021/jp046161x, 2004.
- Kwamena, N.-O. A., Clarke, J. P., Kahan, T. F., Diamond, M. L., and Donaldson, D. J.: Assessing the importance of heterogeneous reactions of polycyclic aromatic hydrocarbons in the urban atmosphere using the Multimedia Urban Model, *Atmos. Environ.*, 41, 37–50, doi:10.1016/j.atmosenv.2006.08.016, 2007.
- Lammel, G., Sehili, A. M., Bond, T. C., Feichter, J., and Grassl, H.: Gas/particle partitioning and global distribution of polycyclic aromatic hydrocarbons – A modelling approach, *Chemosphere*, 76, 98–106, doi:10.1016/j.chemosphere.2009.02.017, 2009.
- Lammel, G., Klánová, J., Ilić, P., Kohoutek, J., Gasić, B., Kovacic, I., and Škrdlíková, L.: Polycyclic aromatic hydrocarbons in air on small spatial and temporal scales – II. Mass size distributions and gas-particle partitioning, *Atmos. Environ.*, 44, 5022–5027, doi:10.1016/j.atmosenv.2010.08.001, 2010.
- Lammel, G., Heil, A., Stemmler, I., Dvorská, A., and Klánová, J.: On the contribution of biomass burning to POPs (PAHs and PCDDs) in air in Africa, *Environ. Sci. Technol.*, 47, 11616–11624, doi:10.1021/es401499q, 2013.
- Lang, C., Tao, S., Zhang, G., Fu, J., and Simonich, S.: Outflow of polycyclic aromatic hydrocarbons from Guangdong, Southern China, *Environ. Sci. Technol.*, 41, 8370–8375, doi:10.1021/es071853v, 2007.
- Lang, C., Tao, S., Liu, W., Zhang, Y., and Simonich, S.: Atmospheric transport and outflow of polycyclic aromatic hydrocarbons from China, *Environ. Sci. Technol.*, 42, 5196–5201, doi:10.1021/es800453n, 2008.

- Liu, S., Tao, S., Liu, W., Liu, Y., Dou, H., Zhao, J., Wang, L., Wang, J., Tian, Z., and Gao, Y.: Atmospheric polycyclic aromatic hydrocarbons in North China: A winter-time study, *Environ. Sci. Technol.*, 41, 8256–8261, doi:10.1021/es0716249, 2007.
- Lohmann, R. and Lammel, G.: Adsorptive and absorptive contributions to the gas-particle partitioning of polycyclic aromatic hydrocarbons: state of knowledge and recommended parametrization for modeling, *Environ. Sci. Technol.*, 38, 3793–3803, doi:10.1021/es035337q, 2004.
- Lohmann, R., Lee, R. G. M., Abbott, J., Coleman, P., and Jones, K. C.: Verifying emission factors and national POPs emission inventories for the UK using measurements and modelling at two rural locations, *J. Environ. Monit.*, 8, 79–88, doi:10.1039/B511399G, 2006.
- Mantseva, E., Dutchak, S., Rozovskaya, O., and Shatalov, V.: EMEP contribution to the preparatory work for the review of the CLR-TAP Protocol on Persistent Organic Pollutants, EMEP MSC-E Information Note, 2004.
- Mathur, R., Yu, S., Kang, D., and Schere, K. L.: Assessment of the wintertime performance of developmental particulate matter forecasts with the Eta-Community Multiscale Air Quality modeling system, *J. Geophys. Res.*, 113, D0230315, doi:10.1029/2007JD008580, 2008.
- Matthias, V., Aulinger, A., and Quante, M.: Adapting CMAQ to investigate air pollution in north sea coastal regions, *Environ. Model. Softw.*, 23, 356–368, doi:10.1016/j.envsoft.2007.04.010, 2008.
- Matthias, V., Aulinger, A., and Quante, M.: CMAQ simulations of the benzo(a)pyrene distribution over Europe for 2000 and 2001, *Atmos. Environ.*, 43, 4078–4086, doi:10.1016/j.atmosenv.2009.04.058, 2009.
- Meng, F., Zhang, B., Gbor, P., Wen, D., Yang, F., Shi, C., Aronson, J., and Sloan, J.: Models for gas/particle partitioning, transformation and air/water surface exchange of PCBs and PCDD/Fs in CMAQ, *Atmos. Environ.*, 41, 9111–9127, doi:10.1016/j.atmosenv.2007.08.009, 2007.
- Milbrandt, J. A. and Yau, M. K.: A multimoment bulk microphysics parameterization, Part I: Analysis of the role of the spectral shape parameter, *J. Atmos. Sci.*, 62, 3051–3064, doi:10.1175/JAS3535.1, 2005a.
- Milbrandt, J. A. and Yau, M. K.: A multimoment bulk microphysics parameterization, Part II: A proposed three-moment closure and scheme description, *J. Atmos. Sci.*, 62, 3065–3081, doi:10.1175/JAS3535.1, 2005b.
- Milford, C., Fernández-Camacho, R., la Campa, de, A. M. S., Rodríguez, S., Castell, N., Marrero, C., Bustos, J. J., la Rosa, de, J. D., and Stein, A. F.: Black Carbon aerosol measurements and simulation in two cities in south-west Spain, *Atmos. Environ.*, 126, 55–65, doi:10.1016/j.atmosenv.2015.11.026, 2016.
- Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multi-component inorganic aerosols, *Aquat. Geochem.*, 4, 123–152, doi:10.1023/A:1009604003981, 1998.
- Ngabe, B. and Poissant, L.: Polycyclic aromatic hydrocarbons in the air in the St. Lawrence Basin (Quebec), *Environ. Sci. Technol.*, 37, 2094–2099, doi:10.1021/es0210345, 2003.
- Odabasi, M., Cetin, E., and Sofuoglu, A.: Determination of octanol-air partition coefficients and supercooled liquid vapor pressures of PAHs as a function of temperature: Application to gas-particle partitioning in an urban atmosphere, *Atmos. Environ.*, 40, 6615–6625, doi:10.1016/j.atmosenv.2006.05.051, 2006.
- Offenberg, J. H. and Baker, J. E.: Aerosol size distributions of polycyclic aromatic hydrocarbons in urban and over-water atmospheres, *Environ. Sci. Technol.*, 33, 3324–3331, doi:10.1021/es990089c, 1999.
- Otte, T. L. and Pleim, J. E.: The Meteorology-Chemistry Interface Processor (MCIP) for the CMAQ modeling system: updates through MCIPv3.4.1, *Geosci. Model Dev.*, 3, 243–256, doi:10.5194/gmd-3-243-2010, 2010.
- Pacyna, J. M., Breivik, K., Münch, J., and Fudala, J.: European atmospheric emissions of selected persistent organic pollutants, 1970–1995, *Atmos. Environ.*, 37, 119–131, doi:10.1016/S1352-2310(03)00240-1, 2003.
- Pankow, J. F.: Review and comparative-analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere, *Atmos. Environ.*, 21, 2275–2283, doi:10.1016/0004-6981(87)90363-5, 1987.
- Pankow, J. F.: An absorption model of gas/particle partitioning of organic compounds in the atmosphere, *Atmos. Environ.*, 28, 185–188, doi:10.1016/1352-2310(94)90093-0, 1994.
- Pankow, J. F.: Further discussion of the octanol/air partition coefficient K_{oa} as a correlating parameter for gas/particle partitioning coefficients, *Atmos. Environ.*, 32, 1493–1497, doi:10.1016/S1352-2310(97)00383-X, 1998.
- Pankow, J. F.: An absorption model of the gas/aerosol partitioning involved in the formation of secondary organic aerosol, *Atmos. Environ.*, 41, 75–79, doi:10.1016/j.atmosenv.2007.10.060, 2007.
- Pedersen, D. U., Durant, J. L., Penman, B. W., Crespi, C. L., Hemond, H. F., Lafleur, A. L., and Cass, G. R.: Human cell mutagens in respirable airborne particles from the Northeastern United States, 1. Mutagenicity of fractionated samples, *Environ. Sci. Technol.*, 38, 682–689, doi:10.1021/es0347282, 2004.
- Pedersen, D. U., Durant, J. L., Taghizadeh, K., Hemond, H. F., Lafleur, A. L., and Cass, G. R.: Human cell mutagens in respirable airborne particles from the Northeastern United States, 2. Quantification of mutagens and other organic compounds, *Environ. Sci. Technol.*, 39, 9547–9560, doi:10.1021/es050886c, 2005.
- Perraudin, E., Budzinski, H., and Villenave, E.: Kinetic study of the reactions of ozone with polycyclic aromatic hydrocarbons adsorbed on atmospheric model particles, *J. Atmos. Chem.*, 56, 57–82, doi:10.1007/s10874-006-9042-x, 2007.
- Pleim, J. E.: A combined local and nonlocal closure model for the atmospheric boundary layer, Part I: Model description and testing, *J. Appl. Meteor. Climatol.*, 46, 1383–1395, doi:10.1175/JAM2539.1, 2007.
- Pleim, J. E. and Xiu, A.: Development and testing of a surface flux and planetary boundary layer model for application in mesoscale models, *J. Appl. Meteor.*, 34, 16–32, doi:10.1175/1520-0450(1995)034<0034:DATOAS>2.0.CO;2, 1995.
- Pleim, J. E. and Xiu, A. J.: Development of a land surface model, Part II: Data assimilation, *J. Appl. Meteor.*, 42, 1811–1822, doi:10.1175/1520-0450(2003)042<1811:DOALSM>2.0.CO;2, 2003.
- Pleim, J. E., Finkelstein, P. L., Clarke, J. F., and Ellestad, T. G.: A technique for estimating dry deposition velocities based on similarity with latent heat flux, *Atmos. Environ.*, 33, 2257–2268, doi:10.1016/S1352-2310(98)00162-9, 1999.

- Pöschl, U., Letzel, T., Schauer, C., and Niessner, R.: Interaction of ozone and water vapor with spark discharge soot aerosol particles coated with benzo[a]pyrene: O₃ and H₂O adsorption, benzo[a]pyrene degradation, and atmospheric implications, *J. Phys. Chem. A*, 105, 4029–4041, doi:10.1021/jp004137n, 2001.
- Prevedouros, K., Jones, K. C., and Sweetman, A. J.: Modelling the atmospheric fate and seasonality of polycyclic aromatic hydrocarbons in the UK, *Chemosphere*, 56, 195–208, doi:10.1016/j.chemosphere.2004.02.032, 2004.
- Prevedouros, K., Palm-Cousins, A., Gustafsson, Ö., and Cousins, I. T.: Development of a black carbon-inclusive multi-media model: Application for PAHs in Stockholm, *Chemosphere*, 70, 607–615, doi:10.1016/j.chemosphere.2007.07.002, 2008.
- Saffari, A., Daher, N., Samara, C., Voutsas, D., Kouras, A., Manoli, E., Karagkiozidou, O., Vlachokostas, C., Moussiopoulos, N., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Increased biomass burning due to the economic crisis in Greece and its adverse impact on wintertime air quality in Thessaloniki, *Environ. Sci. Technol.*, 47, 13313–13320, doi:10.1021/es403847h, 2013.
- San José, R., Pérez, J. L., Callén, M. S., López, J. M., and Masstral, A.: BaP (PAH) air quality modelling exercise over Zaragoza (Spain) using an adapted version of WRF-CMAQ model, *Environ. Pollut.*, 183, 151–158, doi:10.1016/j.envpol.2013.02.025, 2013.
- Sehili, A. M. and Lammel, G.: Global fate and distribution of polycyclic aromatic hydrocarbons emitted from Europe and Russia, *Atmos. Environ.*, 41, 8301–8315, doi:10.1016/j.atmosenv.2007.06.050, 2007.
- Shatalov, V., Gusev, A., Dutchak, S., Holoubek, I., Mantseva, E., Rozovskaya, O., Sweetman, A., Strukov, B., and Vulykh, N.: Modelling of POP contamination in European region: evaluation of the model performance, Moscow, 2005.
- Shiraiwa, M., Garland, R. M., and Pöschl, U.: Kinetic double-layer model of aerosol surface chemistry and gas-particle interactions (K2-SURF): Degradation of polycyclic aromatic hydrocarbons exposed to O₃, NO₂, H₂O, OH and NO₃, *Atmos. Chem. Phys.*, 9, 9571–9586, doi:10.5194/acp-9-9571-2009, 2009.
- Shiraiwa, M., Pfrang, C., and Pöschl, U.: Kinetic multi-layer model of aerosol surface and bulk chemistry (KM-SUB): the influence of interfacial transport and bulk diffusion on the oxidation of oleic acid by ozone, *Atmos. Chem. Phys.*, 10, 3673–3691, doi:10.5194/acp-10-3673-2010, 2010.
- Silibello, C., Calori, G., Costa, M. P., Dirodi, M. G., Mircea, M., Radice, P., Vitali, L., and Zanini, G.: Benzo[a]pyrene modelling over Italy: comparison with experimental data and source apportionment, *Atmos. Pollut. Res.*, 3, 399–407, doi:10.5094/APR.2012.046, 2012.
- Springmann, M., Knopf, D. A., and Riemer, N.: Detailed heterogeneous chemistry in an urban plume box model: reversible co-adsorption of O₃, NO₂, and H₂O on soot coated with benzo[a]pyrene, *Atmos. Chem. Phys.*, 9, 7461–7479, doi:10.5194/acp-9-7461-2009, 2009.
- Šrám, R., Dostal, M., Libalova, H., Rossner, P., Rossnerova, A., Svecova, V., Topinka, J., and Bartonova, A.: The European hot spot of B[a]P and PM_{2.5} exposure – The Ostrava region, Czech Republic: Health research results, *ISRN Public Health*, 57, 12, doi:10.3109/19396368.2010.500440, 2013.
- Stokes, R. H. and Robinson, R. A.: Interactions in aqueous nonelectrolyte solutions, I. solute-solvent equilibria, *J. Phys. Chem.*, 70, 2126–2131, doi:10.1021/j100879a010, 1966.
- Van Jaarsveld, J., Van Pul, W., and De Leeuw, F.: Modelling transport and deposition of persistent organic pollutants in the European region, *Atmos. Environ.*, 31, 1011–1024, doi:10.1016/S1352-2310(96)00251-8, 1997.
- van Noort, P.: A thermodynamics-based estimation model for adsorption of organic compounds by carbonaceous materials in environmental sorbents, *Environ. Toxicol. Chem.*, 22, 1179–1188, doi:10.1002/etc.5620220601, 2003.
- Wenborn, M. J., Coleman, P. J., Passant, N. R., Lymberidi, E., Sully, J., and Weir, A.: Speciated PAH inventory for the United Kingdom, Report AEAT 3512, available at: https://uk-air.defra.gov.uk/assets/documents/reports/cat08/0512011419_REPFIN_all_nov.pdf (last access: 6 December 2016), 1999.
- WHO: World Health Organization: Health risks of persistent organic pollutants from long-range transboundary air pollution, WHO Regional Office for Europe, Copenhagen, 2003.
- Xiu, A. J. and Pleim, J. E.: Development of a land surface model, Part I: Application in a mesoscale meteorological model, *J. Appl. Meteor.*, 40, 192–209, doi:10.1175/1520-0450(2001)040<0192:DOALSM>2.0.CO;2, 2001.
- Yarwood, G., Rao, S., Yocke, M., and Whitten, G.: Updates to the carbon bond chemical mechanism: CB05, Final report to the US EPA, RT-0400675, available at <http://www.camx.com>. (last access: 6 December 2016), 2005.
- Zelenyuk, A., Imre, D., Beránek, J., Abramson, E., Wilson, J., and Shrivastava, M.: Synergy between secondary organic aerosols and long-range transport of polycyclic aromatic hydrocarbons, *Environ. Sci. Technol.*, 46, 12459–12466, doi:10.1021/es302743z, 2012.
- Zhang, B., Meng, F., Shi, C., Yang, F., Wen, D., Aronsson, J., Gbor, P. K., and Sloan, J. J.: Modeling the atmospheric transport and deposition of polychlorinated dibenzo-p-dioxins and dibenzofurans in North America, *Atmos. Environ.*, 43, 2204–2212, doi:10.1016/j.atmosenv.2009.01.004, 2009.
- Zhang, Y., Seigneur, C., Seinfeld, J. H., Jacobson, M., Clegg, S. L., and Binkowski, F. S.: A comparative review of inorganic aerosol thermodynamic equilibrium modules: similarities, differences, and their likely causes, *Atmos. Environ.*, 34, 117–137, doi:10.1016/S1352-2310(99)00236-8, 2000.
- Zhou, S., Lee, A. K. Y., McWhinney, R. D., and Abbatt, J. P. D.: Burial effects of organic coatings on the heterogeneous reactivity of particle-borne benzo[a]pyrene (BaP) toward ozone, *J. Phys. Chem. A*, 116, 7050–7056, doi:10.1021/jp3030705, 2012.
- Zhou, S., Shiraiwa, M., McWhinney, R. D., Pöschl, U., and Abbatt, J. P. D.: Kinetic limitations in gas-particle reactions arising from slow diffusion in secondary organic aerosol, *Faraday Discuss.*, 165, 391–406, doi:10.1039/c3fd00030c, 2013.
- Zhu, Y., Tao, S., Price, O. R., Shen, H., Jones, K. C., and Sweetman, A. J.: Environmental distributions of benzo[a]pyrene in China: Current and future emission reduction scenarios explored using a spatially explicit multimedia fate model, *Environ. Sci. Technol.*, 49, 13868–13877, doi:10.1021/acs.est.5b00474, 2015.