

Kerr-Newman black hole in the formalism of isolated horizonsM. Scholtz^{*} and A. Flandera[†]*Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University,
V Holešovičkách 2, 182 00 Prague, Czech Republic*Norman Gürlebeck[‡]*ZARM, University of Bremen, Am Fallturm, 28359 Bremen, Germany and DLR Institute for Space Systems,
Linzer Strasse 1, 28359 Bremen, Germany*
(Received 26 May 2017; published 15 September 2017)

The near horizon geometry of general black holes in equilibrium can be conveniently characterized in the formalism of weakly isolated horizons in the form of the Bondi-like expansions (Krishnan B, *Classical Quantum Gravity* **29**, 205006, 2012). While the intrinsic geometry of the Kerr-Newman black hole has been extensively investigated in the weakly isolated horizon framework, the off-horizon description in the Bondi-like system employed by Krishnan has not been studied. We extend Krishnan's work by explicit, nonperturbative construction of the Bondi-like tetrad in the full Kerr-Newman spacetime. Namely, we construct the Bondi-like tetrad which is parallelly propagated along a nontwisting null geodesic congruence transversal to the horizon and provide all Newman-Penrose scalars associated with this tetrad. This work completes the description of the Kerr-Newman spacetime in the formalism of weakly isolated horizons and is a starting point for the investigation of deformed black holes.

DOI: 10.1103/PhysRevD.96.064024

I. INTRODUCTION

The formalism of weakly isolated horizons (WIHs) provides a powerful framework for the analysis of black holes in equilibrium which are a final state of gravitational collapse [1–4]. One of the earliest applications of the new formalism was the calculation of statistical mechanical entropy of a black hole in the framework of loop quantum gravity [5]. But WIHs have a plethora of applications also in classical general relativity (see [6] for a review) and they exhibit a number of properties which make them more realistic in astrophysical settings as compared to standard stationary axisymmetric black holes. For example, WIHs can be embedded in otherwise dynamical spacetimes, they admit the presence of radiation or matter in the neighborhood of the black hole, and they do not require asymptotic flatness. Despite this more general context, WIHs satisfy the usual laws of thermodynamics [7]. Moreover, there is a well defined notion of multipole moments for axially symmetric WIHs [8]. These moments are intrinsic to the horizon and therefore allow to define the mass or angular momentum of a black hole quasilocally without the reference to spatial infinity, which is necessary for the definition of Geroch-Hansen multipole moments [9,10]. However, for the higher moments both definitions do not agree in general [11,12] and not even in the case of a Kerr black hole [8].

WIHs represent a class of black holes much wider than the standard Kerr-Newman family of solutions [13] describing isolated axially symmetric and stationary charged black holes. Nonetheless, the Kerr-Newman metric is the prototypical example of a WIH with well-understood geometry and physical interpretation. One also expects that the geometry of isolated black holes distorted by, e.g., accreting matter or electromagnetic fields outside the black hole, will deviate only slightly from the Kerr solution; even such small deviations might be measurable in the future experiments [14,15]. From the mathematical point of view, one can generate a large class of solutions representing black holes whose intrinsic geometry coincides with the geometry of Kerr-Newman black holes but is (even strongly) distorted in a neighborhood of the horizon by the appropriate choice of the initial data [16].

The near horizon geometry of a general WIH has been investigated in [17] in the Newman-Penrose formalism. In the neighborhood of the horizon it is possible to introduce coordinates similar to those used by Bondi [18] in the neighborhood of null infinity and to find a solution of the field equations near the horizon in a form resembling asymptotic expansions of Newman-Penrose and Newman-Unti near null infinity [19,20]. An important feature of the Newman-Penrose formalism is that the field equations naturally split into constraint and evolution equations. Thanks to the properties of WIHs, all constraints can be solved explicitly. Using the evolution part of the equations one can construct an expansion of the solution near the horizon to arbitrary order. However, in this approach, it is not evident how to choose the initial data in order to

^{*}scholtz@utf.mff.cuni.cz[†]flandera.ales@utf.mff.cuni.cz[‡]norman.guerlebeck@zarm.uni-bremen.de

reproduce the standard Kerr-Newman solution, nor what the explicit form of the Bondi-like tetrad used in [17] is and what the corresponding Newman-Penrose scalars are.

The formalism of [17] has been recently employed in [21] where we discussed the Meissner effect, i.e., the expulsion of electromagnetic fields from extremal horizons, in the language of WIHs. We have shown that the Meissner effect is an inherent property of extremal, stationary and axisymmetric horizon and takes place also in the strong field regime and independently of the deformations of the black hole. In order to judge, if specific instances of the deformations are physically viable, a description of the full Kerr-Newman spacetime in the formalism of WIHs explicit expressions for Newman-Penrose quantities in the Bondi-like tetrad is essential.

Whether a given WIH coincides with the horizon of the Kerr metric can be decided using the conditions found in [22]. This result however includes only the intrinsic geometry of the horizon or the near horizon geometry up to the first order. Bondi-like coordinates in the Kerr spacetime adapted to the null infinity have been introduced in [23], but the corresponding tetrad has not been discussed. In [21] we considered near horizon geometry of the Kerr spacetime up to the first order in Bondi-like coordinates and the Bondi-like tetrad of [17].

In this paper we complete the description of the Kerr-Newman black hole in the formalism of WIHs. We perform the construction of [17] explicitly for the Kerr-Newman spacetime. In particular, we construct a null tetrad adapted to the horizon which is covariantly constant along the nontwisting null congruence transversal to the horizon. We compute all Newman-Penrose scalars with respect to this tetrad and infer the initial data which has to be given on the horizon and on the transversal null hypersurface in order to reproduce the Kerr-Newman spacetime. As explained above, this is a starting point for the analysis of physically reasonable deformations of the Kerr black hole.

In Sec. II, we summarize the description of the Kerr-Newman spacetime in standard horizon-penetrating coordinates and Kinnersley null tetrad. The definition of a

WIH and the basic properties of the Bondi-like tetrad employed in [17] are reviewed in Sec. III. The explicit construction is carried out in Sec. IV. First, we find the parametrization of nontwisting null geodesic congruences transversal to the horizon. Then, we find the desired tetrad by a sequence of Lorentz transformations of the Kinnersley one. In order to obtain an explicit solution up to integration, we employ the coordinate transformation of [23] but adapted to the horizon rather than to null infinity. In Sec. V, we summarize the results and extract the initial data reproducing the Kerr-Newman spacetime. Definitions of the Newman-Penrose formalism and, in particular, the transformation properties of the Newman-Penrose scalars are summarized in Appendix A. Finally, we visualize and compare the geodesic congruences induced by the Kinnersley tetrad and by the Bondi-like tetrad in Appendix B.

II. KERR-NEWMAN METRIC

The main purpose of this section is to set up the notation and conventions used in this paper and to provide equations for later references. Typically, we employ the abstract index notation [24] and denote the abstract indices by Latin letters from the beginning of the alphabet, a, b, \dots . Greek indices will take values from 0 to 3 and they will denote components of a tensor with respect to particular coordinates. Indices I, J, \dots will take values 2,3. We use the signature $(+ - - -)$ so that the Newman-Penrose (NP) null tetrad [19,25] is normalized by the conditions $\ell^a n_a = 1, m^a \bar{m}_a = -1$. For a review of the relevant definitions and relations of the NP formalism, see Appendix A. The Riemann tensor is defined by $2\nabla_{[c}\nabla_{d]}X^a = -R^a_{bcd}X^b$, where square brackets denote the total antisymmetrization. In the NP formalism, Einstein's equations for electro-vacuum read $\Phi_{mn} = \phi_m \bar{\phi}_n, \Lambda = 0$.

We start with the Kerr-Newman metric describing a black hole of mass M , spin a , and charge Q in the ingoing null coordinates $x^\mu = (v, r, \theta, \varphi)$, in which the line element takes the form [13,26]

$$\begin{aligned} ds^2 &= \left(1 - \frac{2Mr - Q^2}{|\rho|^2}\right)dv^2 - 2dv\,dr + \frac{2a}{|\rho|^2}(2Mr - Q^2)\sin^2\theta dv\,d\varphi \\ &\quad + 2a\sin^2\theta dr\,d\varphi - |\rho|^2 d\theta^2 + \frac{\sin^2\theta}{|\rho|^2}(\tilde{\Delta}a^2\sin^2\theta - (a^2 + r^2)^2)d\varphi^2, \end{aligned} \quad (1)$$

where the functions ρ and $\tilde{\Delta}$ are given by relations

$$\rho = r + ia\cos\theta, \quad \tilde{\Delta} = a^2 + r^2 - 2Mr + Q^2. \quad (2)$$

The outer and inner horizons are located at

$$r_\pm = M \pm \sqrt{M^2 - a^2 - Q^2} \quad (3)$$

respectively. We will write $X \dot{=} Y$ when the two quantities are equal on the outer horizon r_+ , for example, $\tilde{\Delta} \dot{=} 0$.

In order to analyze the Kerr-Newman metric in the NP formalism, we first introduce the standard Kinnersley null tetrad [26,27] adapted to the principal null directions of the Kerr-Newman metric,

$$\begin{aligned}\ell_K &= \partial_v + \frac{\tilde{\Delta}}{2(a^2 + r^2)} \partial_r + \frac{a}{a^2 + r^2} \partial_\varphi, \\ n_K &= -\frac{a^2 + r^2}{|\rho|^2} \partial_r, \\ m_K &= \frac{1}{\sqrt{2}\rho} \left(ia \sin \theta \partial_v + \partial_\theta + \frac{i}{\sin \theta} \partial_\varphi \right),\end{aligned}\quad (4)$$

where the fourth vector of the tetrad \bar{m}_K^a is the complex conjugate of m_K^a . Notice also that the triad $(\ell_K^a, m_K^a, \bar{m}_K^a)$ is tangent to the horizon. The spin coefficients associated with the tetrad (4) are

$$\begin{aligned}\kappa_K &= \sigma_K = \nu_K = \lambda_K = 0, & \gamma_K &= -\frac{a(a + ir \cos \theta)}{\rho \bar{\rho}^2}, \\ \eta_K &= -\frac{\tilde{\Delta}}{2\bar{\rho}(a^2 + r^2)}, & \tau_K &= -\frac{ia \sin \theta}{\sqrt{2}|\rho|^2}, \\ \varepsilon_K &= -\frac{Ma^2 + rQ^2 - Mr^2}{2(a^2 + r^2)^2}, & \mu_K &= -\frac{a^2 + r^2}{\rho \bar{\rho}^2}, \\ \pi_K &= \frac{ia \sin \theta}{\sqrt{2}\bar{\rho}^2}, & a_K &= \frac{ia - r \cos \theta}{\sqrt{2}\bar{\rho}^2 \sin \theta},\end{aligned}\quad (5)$$

where we have denoted $a_K = \alpha_K - \bar{\beta}_K$; in addition, $\pi_K = \alpha_K + \bar{\beta}_K$. These relations can be geometrically interpreted as follows, cf. Appendixes A and B. The vector ℓ_K^a is tangent to a congruence of null curves, which are geodesics ($\kappa_K = 0$) and shear-free ($\sigma_K = 0$). The expansion and twist of ℓ_K^a vanish on the horizon ($\eta_K \dot{=} 0$). Similarly, n_K^a is tangent to null geodesics ($\nu_K = 0$), which are shear-free ($\lambda_K = 0$), but have nonvanishing expansion ($\text{Re}\mu_K$) and twist ($\text{Im}\mu_K$). Thus, the vector field n_K^a is not hypersurface orthogonal. In fact, vectors ℓ_K^a and n_K^a are, at each point, the principal null directions of the Weyl tensor, so that the only nonvanishing Weyl scalar is

$$\Psi_2^K = -\frac{M}{\bar{\rho}^3} + \frac{Q^2}{\bar{\rho}^3 \rho}. \quad (6)$$

In the presence of charge, the nonvanishing component of the trace-free part of the Ricci tensor is

$$\Phi_{11}^K = \frac{Q^2}{2|\rho|^4}, \quad (7)$$

while the scalar curvature Λ vanishes. Comparing the NP form of Einstein's equation $\Phi_{11}^K = |\phi_1^K|^2$ and (7), and using the Maxwell equations, we find

$$\phi_1^K = \frac{Q}{\sqrt{2}\bar{\rho}^2}, \quad (8)$$

while ϕ_0^K and ϕ_2^K vanish. Hence, the principal null directions of F_{ab} are aligned with ℓ_K^a and n_K^a . The four-potential of the electromagnetic field turns out to be

$$A_\mu dx^\mu = \frac{\sqrt{2}Qr}{|\rho|^2} (dv - a \sin^2 \theta d\varphi). \quad (9)$$

Finally, the Kerr-Newman metric admits a Killing-Yano form [28,29] which, in the Kinnersley tetrad (4), reads

$$Y^{ab} = -2a \cos \theta \ell_K^{[a} n_K^{b]} + 2i r m_K^{[a} \bar{m}_K^{b]}, \quad (10)$$

or, in coordinates,

$$\begin{aligned}Y &= d\varphi \wedge [a^2 \cos \theta \sin^2 \theta dr - r(a^2 + r^2) \sin \theta d\theta] \\ &\quad + dv \wedge [-a \cos \theta dr + ar \sin \theta d\theta].\end{aligned}\quad (11)$$

For any geodesic vector X^a , the one-form $k_a = Y_{ab} X^b$ is covariantly constant along X^a .

III. WEAKLY ISOLATED HORIZONS

A spacetime M is said to admit a *nonexpanding horizon* [4], if it contains a null hypersurface $\mathcal{H} \subset M$ with the topology $\mathbb{R} \times S^2$ on which Einstein's equations hold and the energy-momentum tensor T_{ab} satisfies the energy condition that $T_{ab} \ell^b$ is causal (i.e., timelike or null) and future pointing for any future null vector ℓ_a normal to \mathcal{H} . Moreover, any such normal is nonexpanding. It turns out that the spacetime connection ∇_a induces a preferred connection \mathcal{D}_a on \mathcal{H} and gives rise to a *rotational 1-form* ω_a defined by

$$\mathcal{D}_a \ell^b \doteq \omega_a \ell^b. \quad (12)$$

Since the choice of the null normal ℓ^a is not unique, it is natural to fix it by the requirement [3]

$$[\mathfrak{f}_\ell, \mathcal{D}_a] \ell^b \doteq 0 \quad (13)$$

which is equivalent to $\mathfrak{f}_\ell \omega_a \dot{=} 0$; here \mathfrak{f}_ℓ is the Lie derivative along ℓ^a . This leads to a definition of a *weakly isolated horizon* (WIH) as a nonexpanding horizon \mathcal{H} equipped with an equivalence class of null normals $[\ell^a]$, where elements of $[\ell^a]$ differ just by constant rescaling, satisfying the condition (13). This condition guarantees that the zeroth law of black hole thermodynamics is satisfied on \mathcal{H} and that the pull-back of ℓ^a is a Killing vector of the induced degenerate metric on \mathcal{H} .

Using the geometrical properties of the WIHs it is possible to construct Bondi-like coordinates and a null, Bondi-like, tetrad adapted to these coordinates. Details of this construction and perturbative solutions of the Einstein-Maxwell equations near a WIH were given in [17]. Here, we briefly review the main steps of the construction.

By definition, a given WIH $(\mathcal{H}, [\ell^a])$ has a preferred foliation by topological spheres. A coordinate v on \mathcal{H} is defined by the requirement that it is constant on each sphere S_v of the foliation and, in addition, $Dv \dot{=} 1$, where

$D = \ell^a \nabla_a$. Next, arbitrary coordinates x^I , $I = 2, 3$, are introduced on the sphere \mathcal{S}_0 along with two arbitrary complex null vectors m^a and \bar{m}^a tangent to \mathcal{S}_0 . To get a basis of the space tangent to \mathcal{H} , the vector m^a is propagated along ℓ^a by the condition [30]

$$\mathfrak{L}_\ell m^a \doteq 0. \quad (14)$$

Similarly, coordinates x^I are propagated off the sphere \mathcal{S}_0 along ℓ^a by the condition $Dx^I \doteq 0$ so that we have a coordinate system (v, x^2, x^3) for the entire horizon \mathcal{H} . The triad (ℓ^a, m^a, \bar{m}^a) can now be completed to a full NP tetrad $(\ell^a, n^a, m^a, \bar{m}^a)$ on the horizon. In order to obtain the tetrad in the neighborhood of the horizon, the vector field n^a on \mathcal{H} is extended geodesically off the horizon and the remaining vectors are parallelly transported along the resulting geodesic congruence, i.e., all vectors are propagated off the horizon by the conditions

$$\Delta n^a = \Delta \ell^a = \Delta m^a = 0, \quad (15)$$

where $\Delta = n^a \nabla_a$. The coordinate r is defined as an affine parameter along the geodesics n^a and the coordinates v and x^I are extended off the horizon by the conditions $\Delta v = \Delta x^I = 0$. In this way, the NP tetrad and the coordinates $x^\mu = (v, r, x^2, x^3)$ are introduced on the horizon and in its neighborhood. In these coordinates, the vectors of the null tetrad read

$$\begin{aligned} \ell &= \partial_v + U \partial_r + X^I \partial_I, & n &= -\partial_r, \\ m &= \Omega \partial_r + \xi^I \partial_I. \end{aligned} \quad (16)$$

By construction, the following functions vanish on \mathcal{H} :

$$U \doteq X^I \doteq \Omega \doteq 0. \quad (17)$$

Throughout the paper, we reserve the symbol x^μ for the coordinates introduced in this section. Adopting the terminology of [23], we will refer to the coordinates x^μ and the tetrad (16) as the ‘‘Bondi-like coordinates’’ and the ‘‘Bondi-like tetrad,’’ respectively.

By construction of the tetrad,

$$\gamma = \nu = \tau = 0, \quad \pi = \alpha + \bar{\beta}, \quad (18)$$

everywhere, and $\kappa \doteq 0$ on the horizon. Moreover,

$$\mu = \bar{\mu}, \quad (19)$$

which means that n^a is twist-free and, hence, orthogonal to hypersurfaces \mathcal{N}_v of constant v , which are transversal to the horizon and intersect it in the spherical cuts \mathcal{S}_v . Since the normal to the horizon ℓ^a is by assumption nonexpanding and orthogonal to \mathcal{H} , we have

$$\varrho \doteq 0, \quad (20)$$

which, together with the energy condition imposed on the energy-momentum tensor and the Ricci identities in the NP formalism, implies

$$\sigma \doteq 0, \quad \Psi_0 \doteq \Psi_1 \doteq 0, \quad \phi_0 \doteq 0, \quad (21)$$

i.e., the horizon is also shear-free and there is no gravitational or electromagnetic radiation crossing the horizon.

Having established the null tetrad and the coordinate system, it is possible to solve the Einstein equations perturbatively in the neighborhood of \mathcal{H} . More precisely, we regard the spacetime as a solution to a characteristic initial value problem with the initial data given on the horizon \mathcal{H} and any null hypersurface, say \mathcal{N}_0 , intersecting the horizon (see [31,32] for the precise formulation and existence results).

Adopting the notation of [4,17], the initial data on \mathcal{S}_0 consists of the following NP scalars:

$$\mathcal{S}_0: \pi^{(0)}, \quad \phi_1^{(0)}, \quad \mu^{(0)}, \quad \lambda^{(0)}, \quad \xi^I|_{\mathcal{S}_0}, \quad \kappa_{(\ell)}, \quad (22)$$

where the $\xi^I|_{\mathcal{S}_i}$ are the components of m^a on the sphere \mathcal{S}_0 which, in turn, define the two-dimensional metric on \mathcal{S}_0 . By the properties of WIHs, the surface gravity of the normal ℓ^a ,

$$\kappa_{(\ell)} \doteq \varepsilon^{(0)} + \bar{\varepsilon}^{(0)}, \quad (23)$$

is constant over the horizon—the zeroth law of thermodynamics. The quantity $\phi_1^{(0)}$ is the NP component of the electromagnetic field and its real and imaginary parts describe the electric as well as magnetic flux density through \mathcal{S}_0 , respectively. From the Ricci identities in the NP formalism one can then calculate the quantities

$$a^{(0)} = \alpha^{(0)} - \bar{\beta}^{(0)}, \quad \Psi_2^{(0)} \quad \text{and} \quad \Psi_3^{(0)} \quad (24)$$

on \mathcal{S}_0 . In the case of axisymmetric, stationary WIHs, the Weyl scalar $\Psi_2^{(0)}$ encodes the horizon multipole moments [8]. The quantity $a^{(0)}$ defines the connection on \mathcal{S}_0 . In addition, the Ricci identities determine the evolution of all these quantities along the horizon and, as it turns out, only the spin coefficients μ and λ and the Weyl scalar Ψ_3 depend on the coordinate v on \mathcal{H} .

The two remaining NP quantities, the Weyl scalar Ψ_4 and the component of the electromagnetic field ϕ_2 can be specified freely on the transversal null hypersurface \mathcal{N}_0 . The Ricci and Bianchi identities as well as the Maxwell equations in the NP formalism then determine the solution of the full Einstein-Maxwell equations in the neighborhood of \mathcal{H} . The geometrical setup is schematically sketched in Fig. 1.




FIG. 1. Characteristic initial value problem for WIH.

With this formulation of the initial value problem, one can expand all geometrical quantities into a series in the coordinate r and find a perturbative solution of the Einstein-Maxwell equations in a neighborhood of \mathcal{H} . This was done in [17]. The question now arises, how to choose the initial data on \mathcal{H} and \mathcal{N}_0 in order to reproduce any particular spacetime. For a Schwarzschild spacetime, this is a trivial task because the principal null directions of the Weyl tensor are already nontwisting and, therefore, the Kinnissley tetrad reduces to the desired Bondi-like one for $a = 0$. We will show that the physically more interesting solution, the Kerr-Newman spacetime, also allows the analytic construction of the Bondi-like tetrad.

IV. BONDI-LIKE TETRAD FOR KERR-NEWMAN SPACETIME

A. Nontwisting null geodesics

Let (v, r, θ, φ) be the standard ingoing null coordinates introduced in Sec. II. In order to construct a null tetrad which meets the criteria imposed in [17] for the Kerr-Newman spacetime, we first construct a nontwisting null geodesic congruence with the tangent vector n_B^a which will later be identified with the vector of the Bondi-like null tetrad. Using the well-known separability of the geodesic equation on the Kerr-Newman background [33], a generic null geodesic can be characterized by three constants E , L , and \mathcal{K} , in terms of which the components of n_B^a read

$$\begin{aligned} n_B^v &= -\frac{1}{|\rho|^2} \left(a^2 E \sin^2 \theta + aL + \frac{a^2 + r^2}{\tilde{\Delta}} (\sqrt{R} - P) \right), \\ n_B^r &= -\frac{1}{|\rho|^2} \sqrt{R}, \\ n_B^\theta &= -\frac{1}{|\rho|^2} \sqrt{\Theta}, \\ n_B^\varphi &= -\frac{1}{|\rho|^2} \left(aE + L \sin^{-2} \theta + \frac{a}{\tilde{\Delta}} (\sqrt{R} - P) \right), \end{aligned} \quad (25)$$

where functions Θ , P , and R are defined by

$$\begin{aligned} \Theta &= \mathcal{K} - (L + aE)^2 + (a^2 E^2 - L^2 \sin^{-2} \theta) \cos^2 \theta, \\ P &= aL + E(a^2 + r^2), \quad R = P^2 - \mathcal{K} \tilde{\Delta}. \end{aligned} \quad (26)$$

The constants of motion E and L represent the energy and the angular momentum of the geodesic and \mathcal{K} is the Carter constant which arises from the separation of the Hamilton-Jacobi equation or from the projection of the Killing tensor of the Kerr-Newman spacetime [34].

In the next step, we promote (25) to a geodesic *congruence*, where each geodesic of the congruence is parametrized by possibly different values of E , L , and \mathcal{K} . In this way, these parameters become functions of the position, assigning the corresponding values to a geodesic passing through a given point. We wish to choose the functions E , L , and \mathcal{K} in such a way that the resulting congruence is nontwisting. In order to accomplish that, we require that the covariant vector $(n_B)_a$ be a gradient, i.e.,

$$(n_B)_\mu dx^\mu \equiv Edv - \frac{a^2}{P + \sqrt{R}} dr + \sqrt{\Theta} d\theta + L d\varphi = dv \quad (27)$$

for some function v . Since ∂_v and ∂_φ are the Killing vectors of the Kerr-Newman metric, we assume $E = E(r, \theta)$ and similarly for L and \mathcal{K} . An inspection of the integrability condition for the existence of a function v in Eq. (27)

$$(dn_B)_\mu \wedge dx^\mu = 0, \quad (28)$$

then shows that E and L must be constant everywhere, while \mathcal{K} must satisfy the condition

$$\sqrt{\Theta} \frac{\partial \mathcal{K}}{\partial \theta} = -\sqrt{R} \frac{\partial \mathcal{K}}{\partial r}, \quad (29)$$

which is equivalent to $n_B^a \nabla_a \mathcal{K} = 0$ and does not impose a new condition, since \mathcal{K} is the constant of motion.

Because L must be a constant, the components of the geodesic (25) become singular on the axis $\theta = 0$. In fact, this is a singularity of a congruence rather than coordinate singularity, since, for example, the expansion $\nabla_a n_B^a$ diverges there. In order to avoid the singular behavior, we set $L = 0$. Notice that although it is natural to expect that nontwisting congruence has zero angular momentum, vanishing of the twist alone is compatible with any value of L .

Then, imposing that the congruence be symmetric under reflection across the equatorial plane [35], we have to choose $\mathcal{K} = a^2 E^2$. Finally, similarly to [23], we also choose $E = 1$ for convenience. Hence, we set

$$L = 0, \quad \mathcal{K} = a^2, \quad (30)$$

so that the congruence (25) simplifies to

$$n_B^v = -\frac{a^2}{|\rho|^2} \left(\sin^2 \theta - \frac{P}{P + \sqrt{R}} \right), \quad (31a)$$

$$n_B^r = -\frac{1}{|\rho|^2} \sqrt{R}, \quad (31b)$$

$$n_B^\theta = -\frac{a \cos \theta}{|\rho|^2}, \quad (31c)$$

$$n_B^\varphi = -\frac{a}{|\rho|^2} \left(1 - \frac{a^2}{P + \sqrt{R}} \right), \quad (31d)$$

where functions P and R now read

$$P = a^2 + r^2, \quad R = r^4 + a^2 r^2 + 2a^2 M r - a^2 Q^2. \quad (32)$$

B. Lorentz transformations

Having found an appropriate null congruence (31), we wish to complete n_B^a to a Bondi-like null tetrad, i.e. find vectors ℓ_B^a and m_B^a which are covariantly constant along n_B^a . The standard technique to parallelly propagate a frame along null geodesics using the Killing-Yano form (11) was developed in [36] and recently generalized to n -dimensional spacetimes admitting a conformal Killing-Yano form in [37]. The vector $k_a = Y_{ab} n_B^b$, where n_B^b is given by (31) is parallelly propagated along n_B^a and it is spacelike rather than null. Still, one can find another parallelly propagated spacelike vector \tilde{k}^a and take the complex linear combination of k^a and \tilde{k}^a in order to form the null vector m_B^a which is covariantly constant along n_B^a , and finally complete the tetrad with vector ℓ_B^a , which is then automatically covariantly constant too. Unfortunately, the frame obtained in this way does not satisfy the conditions imposed on the Bondi-like tetrad, i.e. the vector m_B^a is not tangent to the horizon and ℓ_B^a is not a normal to the horizon. Thus, one has to rotate the frame on the horizon to a desired Bondi-like tetrad with an appropriate matrix R and then require the matrix to be constant along n_B^a . However, the outlined procedure requires explicit solution of the geodesic equation at two stages: finding vector \tilde{k}^a and propagating the matrix R off the horizon. Such explicit solution can be found in terms of the elliptic integrals of the first kind but one has to solve the polynomial equation of the fourth order.

Because of these complications with the standard methods, we adopt a different approach in which we do not need the Killing-Yano form at all. In our approach we also get integrals which either cannot be calculated explicitly or the resulting formulas are too complex to be included in the paper. However, we present the tetrad in the form suitable both for symbolic manipulations and numerical calculations. We perform a sequence of Lorentz transformations which rotate the initial Kinnersley tetrad (4) to a Bondi-like

tetrad. Using the boost and null rotation about ℓ_K^a , we rotate the vector n_K^a to the direction (31), then by a spin and null rotation about n_K^a we eliminate the spin coefficients γ and τ , which yields the triad (ℓ_B, m_B, \bar{m}_B) tangent to the horizon and parallelly propagated along n_B^a . By this method we get not only the tetrad, but also the spin coefficients and the Weyl and Maxwell scalars, because it is easy to transform the spin coefficients step by step but very difficult to calculate them directly from the resulting tetrad, even with the help of computer algebra systems.

1. Boost

In the first step, we perform a boost (A8) in the plane spanned by ℓ_K^a and n_K^a of the Kinnersley tetrad (4) with the parameter

$$A^2 = \frac{2P}{P + \sqrt{R}}, \quad (33)$$

obtaining a new tetrad $(\ell_1, n_1, m_1, \bar{m}_1)$ and corresponding spin coefficients.

2. Null rotation about ℓ^a

Next, we rotate the tetrad about ℓ_1^a to a new tetrad $(\ell_2, n_2, m_2, \bar{m}_2)$ such that the vector field n_2^a coincides with the nontwisting vector field (31). We choose the parameter c of the null rotation (A16) to be

$$c = -\frac{ae^{-i\theta}}{\sqrt{2}\bar{\rho}}. \quad (34)$$

Then, n_2 is given by (31).

3. Spin

At this stage, the vector n_2^a is tangent to the desired nontwisting, affinely parametrized congruence of geodesics, i.e. $\Delta_2 n_2^a = 0$ with the respective radial derivative $\Delta_2 = n_2^a \nabla_a$. Moreover, the triad $(\ell_2^a, m_2^a, \bar{m}_2^a)$ is tangent to the horizon, where ℓ_2^a is a generator of the horizon satisfying (13). However, this triad is not covariantly constant along n_2^a , since we have [cf. the transport equations (A4c)]

$$\Delta_2 \ell_2^a = -\bar{\tau}_2 m_2^a - \tau_2 \bar{m}_2^a, \quad (35a)$$

$$\Delta_2 m_2^a = -\tau_2 n_2^a + (\gamma_2 - \bar{\gamma}_2) m_2^a. \quad (35b)$$

The coefficient γ_2 which is now purely imaginary (n_2^a is affinely parametrized) can be eliminated completely by the spin (A11) with the parameter $\chi = -\theta/2$, i.e., $m_3^a = e^{-i\theta} m_2^a$. The nonvanishing spin coefficients now are

$$\tau_3 = \frac{a}{\sqrt{2}|\rho|^2} \left(\frac{\tilde{\Delta}}{P + \sqrt{R}} - ie^{-i\theta} \sin \theta \right), \quad (36a)$$

$$\varrho_3 = -\frac{\tilde{\Delta}}{\bar{\rho}(P + \sqrt{R})}, \quad (36b)$$

$$\varepsilon_3 = \frac{r}{a^2} - \frac{1}{2\sqrt{R}} \left(\frac{2r^3}{a^2} + M + r \right), \quad (36c)$$

$$\begin{aligned} \pi_3 &= \frac{P - \sqrt{R}}{\sqrt{2}a\bar{\rho}^2} - \frac{\sqrt{2}a}{\bar{\rho}} \left(\frac{r}{a^2} - \frac{M + r + 2r^3/a^2}{2\sqrt{R}} \right) \\ &\quad + \left(\frac{2ia - \rho \cos \theta}{2\sqrt{2}\bar{\rho}^2 \sin \theta} + \frac{\cot \theta}{2\sqrt{2}\bar{\rho}} \right) e^{i\theta}, \end{aligned} \quad (36d)$$

$$\begin{aligned} \alpha_3 &= -\frac{a}{\sqrt{2}\bar{\rho}} \left(\frac{r}{a^2} - \frac{M + r + 2r^3/a^2}{2\sqrt{R}} - \frac{P - \sqrt{R}}{a^2\bar{\rho}} \right) \\ &\quad + \frac{e^{i\theta}}{2\sqrt{2}\bar{\rho}} \left(\frac{2ia - \rho \cos \theta}{\bar{\rho} \sin \theta} - i \right). \end{aligned} \quad (36e)$$

$$\begin{aligned} \beta_3 &= \frac{1}{2\sqrt{2}\bar{\rho}} \left[e^{-i\theta} (\cot \theta - i) \right. \\ &\quad \left. - 2a \left(\frac{r}{a^2} - \frac{M + r + 2r^3/a^2}{2\sqrt{R}} \right) \right], \end{aligned} \quad (36f)$$

$$\begin{aligned} \mu_3 &= \frac{1}{2\sqrt{R}|\rho|^2} (-2r^3 - a^2(M + r) \\ &\quad - a\sqrt{R} \cos 2\theta \csc \theta), \end{aligned} \quad (36g)$$

$$\begin{aligned} \lambda_3 &= \frac{a}{2\sqrt{R}\bar{\rho}^3} (-2aQ^2 + ar(3M + r) \\ &\quad + i(2r^3 + a^2(M + r)) \cos \theta \\ &\quad + \sqrt{R}(r \cos 2\theta - ia \cos \theta) \csc \theta). \end{aligned} \quad (36h)$$

By now, the only freedom in the choice of the tetrad is the null rotation about n^a . In order to preserve the property that the triad $(\ell_3^a, m_3^a, \bar{m}_3^a)$ be tangent to the horizon, the parameter d of null rotation (A19) must vanish on the horizon. The spin coefficient τ transforms, according to (A20), by $\tau_4 = \tau_3 - \Delta_3 d$, where $\Delta_3 = n_3^a \nabla_a$. Thus, in order to eliminate τ_3 we have to solve the equation

$$\Delta_3 d = \tau_3, \quad d \dot{=} 0. \quad (37)$$

It is exactly the initial condition $d \dot{=} 0$ which makes the problem difficult, otherwise the equation could be easily solved with the ansatz $d = f(r) + g(\theta)$. In order to implement the initial condition, we have to employ a coordinate

transformation which will eliminate the nonradial components of n_3^a .

4. Bondi-like coordinates

Having satisfied the integrability conditions (28) for Eq. (27), we can employ v as a new coordinate, eliminating the v -component of n_3^a . The angular coordinates which are constant along n_3^a can be conveniently introduced following the procedure of [23]. Hence, we define the new coordinates v , ϑ , and $\tilde{\phi}$ by

$$v = v - \int_{r_+}^r \frac{a^2 dr}{P + \sqrt{R}} + a \sin \theta, \quad (38a)$$

$$\sin \theta = \tanh X, \quad (38b)$$

$$\tilde{\phi} = \varphi + J(r), \quad (38c)$$

where

$$X = \alpha(r) + \text{arth} \sin \vartheta, \quad (39a)$$

$$\alpha(r) = \int_{r_+}^r \frac{adu}{\sqrt{u^4 + a^2u^2 + 2a^2Mu - a^2Q^2}}, \quad (39b)$$

$$J(r) = - \int_{r_+}^r \frac{a}{P(u) + \sqrt{R(u)}} \left(1 + \frac{u^2}{\sqrt{R(u)}} \right) du. \quad (39c)$$

With these choices of α and J , coordinates $(\vartheta, \tilde{\phi})$ coincide with the coordinates (θ, φ) on the horizon (in [23] they coincide at infinity), and equatorial plane is given everywhere by $\theta = \vartheta = \pi/2$. Moreover, ϑ and $\tilde{\phi}$ are constant along n_3^a . Using the relations

$$\begin{aligned} dv &= dv + \left(\frac{1}{P + \sqrt{R}} - \frac{1}{\sqrt{R} \cosh^2 X} \right) a^2 dr \\ &\quad - \frac{ad\vartheta}{\cosh^2 X \cos \vartheta}, \end{aligned} \quad (40a)$$

$$d\theta = \frac{1}{\cosh X} \left(\frac{adr}{\sqrt{R}} + \frac{d\vartheta}{\cos \vartheta} \right), \quad (40b)$$

$$d\varphi = \frac{a}{\sqrt{R}} \left(1 - \frac{a^2}{P + \sqrt{R}} \right) dr + d\tilde{\phi}, \quad (40c)$$

one can deduce the form of the metric tensor in these coordinates:

$$\begin{aligned} ds^2 &= \left(1 + \frac{Q^2 - 2Mr}{|\rho|^2} \right) \left(dv - \frac{2a}{\cosh^2 X \cos \vartheta} d\vartheta \right) dv - \frac{2|\rho|^2}{\sqrt{R}} dv dr - \frac{2a}{|\rho|^2} (Q^2 - 2Mr) \tanh^2 X dv d\varphi \\ &\quad + \frac{2a^2(Q^2 - 2Mr) \tanh^2 X}{|\rho|^2 \cosh^2 X \cos \vartheta} d\vartheta d\tilde{\phi} - \frac{r^4 + a^2(2Mr + r^2 - Q^2) \cosh^{-2} X}{|\rho|^2 \cosh^2 X \cos^2 \vartheta} d\vartheta^2 - \frac{\tanh^2 X}{|\rho|^2} (a^2 \tilde{\Delta} \operatorname{sech}^2 X + R) d\tilde{\phi}^2. \end{aligned} \quad (41)$$

This is the metric given in [23], except for that we used the horizon-penetrating Kerr coordinates instead of Boyer-Lindquist coordinates. The vector field n_3^a given by (31) has, in these coordinates, the simple form

$$n_3 = -\frac{\sqrt{R}}{|\rho|^2} \partial_r, \quad (42)$$

where ρ is now

$$\rho = r + i \operatorname{asech} X. \quad (43)$$

The remaining tetrad vectors read in these coordinates

$$\ell_3 = \partial_v + \frac{\tilde{\Delta}}{P + \sqrt{R}} \partial_r - \frac{a\tilde{\Delta} \cos \vartheta}{\sqrt{R}(P + \sqrt{R})} \partial_\vartheta + a^{-1} \left(1 - \frac{r^2}{\sqrt{R}} \right) \partial_{\tilde{\phi}}, \quad (44a)$$

$$m_3 = \frac{1}{\sqrt{2}\rho} \left[-\frac{a\tilde{\Delta}}{P + \sqrt{R}} \partial_r + \cos \vartheta \left(\frac{P}{\sqrt{R}} - i \sinh X \right) \partial_\vartheta + \left(\frac{r^2}{\sqrt{R}} + \frac{i}{\sinh X} \right) \partial_{\tilde{\phi}} \right]. \quad (44b)$$

The spin coefficient τ_3 has now the form

$$\tau_3 = \frac{a}{\sqrt{2}|\rho|^2} \left(\frac{\tilde{\Delta}}{P + \sqrt{R}} - 1 + \frac{1}{1 + i \sinh X} \right). \quad (45)$$

5. Null rotation about n^a

Now we can integrate Eq. (37) to get

$$d(r, \vartheta) = - \int_{r_+}^r \frac{|\rho(u, \vartheta)|^2}{\sqrt{R(u)}} \tau_3(u, \vartheta) du, \quad (46)$$

where τ_3 is given by (45), and perform null rotation about n_3^a , Eq. (A19), with the parameter d . This yields the desired Bondi-like tetrad $(\ell_B, n_B, m_B, \bar{m}_B)$. By construction d vanishes on the horizon and hence the triad (ℓ_B, m_B, \bar{m}_B) is tangent to the horizon and parallelly propagated along $n_B^a = n_3^a$.

Since d cannot be evaluated explicitly, we cannot give more explicit expressions for the spin coefficients than (36). In terms of those, the spin coefficients for the rotated tetrad are given by, cf. (A20),

$$\begin{aligned} \kappa_B &= d(2\varepsilon_3 + \varrho_3) + d^2(\pi_3 + 2\alpha_3) + d^3\lambda_3 \\ &\quad + (\tau_3 + 2\beta_3)|d|^2 + \mu_3 d^2 \bar{d} - |d|^2 \Delta_3 d \\ &\quad - D_3 d - d\bar{D}_3 d - \bar{d}\delta_3 d, \end{aligned} \quad (47a)$$

$$\sigma_B = d(\tau_3 + 2\beta_3) + \mu_3 d^2 - d\Delta_3 d - \delta_3 d, \quad (47b)$$

$$\varrho_B = \varrho_3 + 2d\alpha_3 + \bar{d}\tau_3 + d^2\lambda_3 - \bar{d}\Delta_3 d - \bar{\delta}_3 d, \quad (47c)$$

$$\varepsilon_B = \varepsilon_3 + d(\alpha_3 + \pi_3) + \beta_3 \bar{d} + \lambda_3 d^2 + \mu_3 |d|^2, \quad (47d)$$

$$\beta_B = \beta_3 + d\mu_3, \quad (47e)$$

$$\alpha_B = \alpha_3 + d\lambda_3, \quad (47f)$$

$$\pi_B = \pi_3 + d\lambda_3 + \bar{d}\mu_3, \quad (47g)$$

$$\mu_B = \mu_3, \quad (47h)$$

$$\lambda_B = \lambda_3, \quad (47i)$$

$$\tau_B = \gamma_B = \nu_B = 0, \quad (47j)$$

where the spin coefficients with subscript 3 are given by (36), the operators $D_3 = l_3^a \nabla_a$, $\Delta_3 = n_3^a \nabla_a$, and $\delta_3 = m_3^a \nabla_a$ are given by (42) and (44), and the derivatives of d are

$$\frac{\partial d}{\partial r} = -\frac{|\rho|^2}{\sqrt{R}} \tau_3, \quad (47k)$$

$$\frac{\partial d}{\partial \vartheta} = \int_{r_+}^r \frac{a}{\sqrt{2R(u)}} \frac{i \cosh X(u, \vartheta)}{(1 + i \sinh X(u, \vartheta))^2} du. \quad (47l)$$

In the last step we perform two remaining coordinate transformations. First we define a new radial coordinate by rescaling r ,

$$r = \int_{r_+}^r \frac{|\rho(u, \vartheta)|^2}{\sqrt{R(u)}} du, \quad (48)$$

so that r is an affine parameter along n^a

$$n_B = -\partial_r, \quad (49)$$

and vanishes on the horizon. In the rest of the paper, the variable $r = r(r)$ will always be understood as the function of this new coordinate r . Next, in order to eliminate the ϕ -component of ℓ^a on the horizon, we perform the last coordinate transformation

$$\phi = \tilde{\phi} - \frac{a\nu}{a^2 + r_+^2}, \quad (50)$$

which brings the tetrad into the form (16).

V. RESULTS

A. NP-quantities

Now we can summarize the obtained results. We have found the Bondi-like coordinates $x^\mu = (\nu, r, \vartheta, \phi)$ in which,

following the notation of [17], the Bondi-like null tetrad is of the form

$$\begin{aligned} \ell_B &= \partial_\nu + U\partial_r + X^I\partial_I, \\ n_B &= -\partial_r, \\ m_B &= \Omega\partial_r + \xi^I\partial_I, \end{aligned} \quad (51)$$

where $I = 2, 3$ and the components of the tetrad read

$$\begin{aligned} U &= \frac{\tilde{\Delta}}{\sqrt{2}\sqrt{R}(P + \sqrt{R})} \left(\sqrt{2}|\rho|^2 - a(\rho d + \bar{\rho}\bar{d}) \left(1 + \frac{a \cos \vartheta}{|\rho|^2} \frac{\partial r}{\partial \vartheta} \right) - \sqrt{2} \frac{\partial r}{\partial \vartheta} a \cos \vartheta \right) - |d|^2 \\ &\quad + \sqrt{2} \frac{\partial r}{\partial \vartheta} \operatorname{Re} \frac{d}{\bar{\rho}} (1 + i \sinh X), \end{aligned} \quad (52a)$$

$$X^2 = \frac{a\tilde{\Delta} \cos \vartheta}{\sqrt{2}\sqrt{R}(P + \sqrt{R})} \left(-\sqrt{2} + \frac{a\bar{d}}{\rho} + \frac{ad}{\bar{\rho}} \right) + \frac{\cos \vartheta}{\sqrt{2}} \left(\frac{\bar{d}}{\rho} (1 - i \sinh X) + \frac{d}{\bar{\rho}} (1 + i \sinh X) \right), \quad (52b)$$

$$X^3 = -\frac{a}{a^2 + r_+^2} + a^{-1} \left(1 - \frac{r^2}{\sqrt{R}} \right) + \frac{\bar{d}}{\sqrt{2}\rho} \left(\frac{r^2}{\sqrt{R}} + \frac{i}{\sinh X} \right) + \frac{d}{\sqrt{2}\bar{\rho}} \left(\frac{r^2}{\sqrt{R}} - \frac{i}{\sinh X} \right), \quad (52c)$$

$$\Omega = -d - \frac{a\tilde{\Delta}\bar{\rho}}{\sqrt{2}\sqrt{R}(P + \sqrt{R})} + \frac{\cos \vartheta}{\sqrt{2}\rho} \frac{\partial r}{\partial \vartheta} \left(\frac{P}{\sqrt{R}} - i \sinh X \right), \quad (52d)$$

$$\xi^2 = \frac{\cos \vartheta}{\sqrt{2}\rho} \left(\frac{P}{\sqrt{R}} - i \sinh X \right), \quad (52e)$$

$$\xi^3 = \frac{1}{\sqrt{2}\rho} \left(\frac{r^2}{\sqrt{R}} + \frac{i}{\sinh X} \right), \quad (52f)$$

and where X is given by (39a). Coordinates ν, r, ϑ , and ϕ are related to the standard ingoing null coordinates v, r, θ , and φ by (38a), (48), (38b), (38c), and (50), respectively; the functions α and d are given by (39b) and (46), respectively. Functions $\rho, \tilde{\Delta}, P$, and R are given by Eqs. (2) and (32), and the variable r is related to the coordinate r by (48). Finally,

$$\frac{\partial r}{\partial \vartheta} = - \int_{r_+}^r \frac{2a^2}{\sqrt{R(u)}} \frac{\sinh X(u, \vartheta)}{\cosh^3 X(u, \vartheta) \cos \vartheta} du. \quad (53)$$

In the tetrad (51), the following spin coefficients vanish:

$$\gamma_B = \nu_B = \tau_B = 0; \quad (54)$$

these equalities imply that n_B^a is an affinely parametrized geodesic, and both ℓ_B^a and m_B^a are covariantly constant along n_B^a . The remaining spin coefficients are given by (47).

In particular, the spin coefficient μ_B is real which means that the congruence n_B^a is nontwisting. The spin coefficients σ_B and λ_B which describe the shear of ℓ_B^a and n_B^a , respectively, do not vanish; another difference from the Kinnersley tetrad.

In type D spacetimes and in the tetrad adapted to the principal null directions, the sole nonvanishing Weyl scalar is Ψ_2 . The Bondi-like tetrad is not adapted to these null directions anymore so that the full set of Weyl scalars is given by

$$\Psi_0^B = 6d^2 \left(1 - \frac{ad}{\sqrt{2}\bar{\rho}} \right)^2 \Psi_2^K, \quad (55a)$$

$$\Psi_1^B = 3d \left(\frac{a^2 d^2}{\bar{\rho}^2} - \frac{3ad}{\sqrt{2}\bar{\rho}} + 1 \right) \Psi_2^K, \quad (55b)$$

$$\Psi_2^B = \left(\frac{3a^2 d^2}{\bar{\rho}^2} - \frac{3\sqrt{2}ad}{\bar{\rho}} + 1 \right) \Psi_2^K, \quad (55c)$$

$$\Psi_3^B = \frac{3a}{\bar{\rho}} \left(\frac{ad}{\bar{\rho}} - \frac{1}{\sqrt{2}} \right) \Psi_2^K, \quad (55d)$$

$$\Psi_4^B = \frac{3a^2}{\bar{\rho}^2} \Psi_2^K, \quad (55e)$$

where Ψ_2^K is given by (6). Similarly, for the components of the electromagnetic field we have

$$\phi_0^B = d \left(2 - \frac{\sqrt{2}ad}{\bar{\rho}} \right) \phi_1^K, \quad (56a)$$

$$\phi_1^B = \left(1 - \frac{\sqrt{2}ad}{\bar{\rho}} \right) \phi_1^K, \quad (56b)$$

$$\phi_2^B = -\frac{\sqrt{2}a}{\bar{\rho}} \phi_1^K \quad (56c)$$

where ϕ_1^K is given by (8).

In nonrotating limit, $a = 0$, the tetrad (51) reduces to the corresponding Kinnersley tetrad, since in this case all Lorentz transformations we applied are identities (except for the spin, whose purpose was to eliminate the coefficient γ , which vanishes for $a = 0$ and, hence, it is not necessary to perform the spin). In other words, for the Reissner-Nordström spacetime, the Kinnersley tetrad is already Bondi-like and $\Psi_3^B, \Psi_4^B, \phi_0^B$, and ϕ_2^B vanish.

B. Initial data

Although the tetrad and the corresponding spin coefficients we found are quite lengthy and containing the function d which cannot be integrated explicitly, we have provided all relations which are necessary to perform calculations in this tetrad. They can be done, for example, symbolically in *Mathematica* or similar software. The formulas are also suitable for numerical calculations, since the calculation of the tetrad components and the spin coefficients involves only numerical integration. In this section we use the Bondi-like tetrad to extract appropriate initial data given on the horizon and transversal null hypersurface which reproduce the Kerr-Newman solution, see Sec. III.

In order to formulate the initial value problem whose solution is the full Kerr-Newman spacetime, we start with the initial data on the initial sphere \mathcal{S}_0 . The only nontrivial components of the Bondi-like tetrad on the horizon are

$$\xi^2 \doteq \frac{e^{-i\vartheta}}{\sqrt{2}\rho^{(0)}}, \quad (57a)$$

$$\xi^3 \doteq \frac{1}{\sqrt{2}\rho^{(0)}} \left(\frac{r_+^2}{P^{(0)}} + i \cot \vartheta \right), \quad (57b)$$

which determine the metric on \mathcal{S}_0 ,

$$\begin{aligned} ds^2|_{\mathcal{S}_0} = & - \left(|\rho^{(0)}|^2 + \frac{a^4 \sin^2 \vartheta \cos^2 \vartheta}{|\rho^{(0)}|^2} \right) d\vartheta^2 \\ & - \frac{2a^2 (P^{(0)})^2}{|\rho^{(0)}|^2} \cos \vartheta \sin^2 \vartheta d\vartheta d\phi - \frac{(P^{(0)})^2 \sin^2 \vartheta}{|\rho^{(0)}|^2} d\phi^2, \end{aligned} \quad (58)$$

where the superscript (0) denotes the value of corresponding quantity on \mathcal{S}^0 . The area element acquires the standard form

$$dS = (r_+^2 + a^2) \sin \vartheta d\vartheta \wedge d\phi. \quad (59)$$

For the spin coefficients we have

$$\epsilon^{(0)} = \frac{r_+ - M}{2P^{(0)}}, \quad (60a)$$

$$a^{(0)} \equiv \alpha^{(0)} - \beta^{(0)} = -\frac{r_+ \cos \vartheta - ia}{\sqrt{2}(\bar{\rho}^{(0)})^2 \sin \vartheta}, \quad (60b)$$

$$\begin{aligned} \mu^{(0)} = & \frac{1}{2P^{(0)}|\rho^{(0)}|^2} (-2r_+^3 - a^2(M + r_+) \\ & - aP^{(0)} \cos 2\vartheta \csc \vartheta), \end{aligned} \quad (60c)$$

$$\begin{aligned} \lambda^{(0)} = & \frac{a}{2P^{(0)}(\bar{\rho}^{(0)})^3} (-2aQ^2 + ar_+(3M + r_+) \\ & + P^{(0)}(r_+ \cos 2\vartheta - ia \cos \vartheta) \csc \vartheta \\ & + i(2r_+^3 + a^2(M + r_+)) \cos \vartheta). \end{aligned} \quad (60d)$$

The surface gravity of the horizon is $\kappa_{(\ell)} = 2\epsilon^{(0)}$. To complete the formulation of the initial value problem, one has to specify the values of Ψ_4 and ϕ_2 on the null hypersurface \mathcal{N}_0 which intersects the horizon at the initial sphere \mathcal{S}_0 :

$$\Psi_4 = \frac{3a^2}{\bar{\rho}^5} \left(-M + \frac{Q^2}{\rho} \right), \quad \phi_2 = -\frac{aQ}{\bar{\rho}^3} \quad \text{on } \mathcal{N}_0. \quad (61)$$

For a general WIH, one has to provide also the values of $\pi^{(0)}$ and $\phi_1^{(0)}$ on \mathcal{S}_0 , which in our case are

$$\pi^{(0)} = \frac{a}{\sqrt{2}\bar{\rho}^{(0)}} \left(\frac{M - r_+}{P^{(0)}} e^{-i\vartheta} + \frac{i}{\bar{\rho}^{(0)}} \sin \vartheta \right), \quad (62a)$$

$$\phi_1^{(0)} = \frac{Q}{\sqrt{2}(\bar{\rho}^{(0)})^2}. \quad (62b)$$

However, for a stationary, axially symmetric horizon, these quantities are solutions to the constraints (cf. [38])

$$\bar{\delta}\pi^{(0)} + (\pi^{(0)})^2 = \kappa_{(\ell)}\lambda^{(0)}, \quad (63a)$$

$$\delta^{(0)}\phi_1^{(0)} + 2\pi^{(0)}\phi_1^{(0)} = \kappa_{(\ell)}\phi_2^{(0)}, \quad (63b)$$

where $\delta^{(0)} \doteq \xi^I \partial_I$, the operator $\bar{\delta}$ is defined by (A15) and $\phi_2^{(0)}$ is the value of ϕ_2 at \mathcal{S}_0 . These constraints were the main ingredients for the proof of the Meissner effect for WIHs in [21]. The Eqs. (60) determine $\Psi_2^{(0)}$ and $\Psi_3^{(0)}$ via the Ricci identities (see also [17])

$$\text{Re}\Psi_2^{(0)} = |a^{(0)}|^2 - \frac{1}{2}(\delta^{(0)}a^{(0)} + \bar{\delta}^{(0)}\bar{a}^{(0)}) + |\phi_1^{(0)}|^2, \quad (64a)$$

$$\text{Im}\Psi_2^{(0)} = -\text{Im}\bar{\delta}\pi^{(0)}, \quad (64b)$$

$$\Psi_3^{(0)} = (\bar{\delta} + \bar{\pi}^{(0)})\lambda^{(0)} - (\bar{\delta} + \pi^{(0)})\mu^{(0)}. \quad (64c)$$

The evolution of the components of the tetrad (51) is given by the NP commutators, the evolution of the spin coefficients is governed by the Ricci identities, the evolution of the Weyl scalars is determined by the Bianchi identities and the evolution of NP components of electromagnetic field is given by the NP form of Maxwell's equations. We do not list these equations here (see, e.g. [25]), because we know already that the solution is the Kerr-Newman spacetime in Bondi-like coordinates equipped with the Bondi-like null tetrad constructed in this paper.

It is worth noting that for a general WIH, Ψ_4 and ϕ_2 are functions given on \mathcal{N}_0 and they are independent from the data given on \mathcal{S}_0 . This is not the case for the Kerr-Newman spacetime, since these quantities are directly given by $\Psi_2^{(0)}$ and $\phi_1^{(0)}$ and their transformation properties under the Lorentz transformations. However, as pointed out already in [16], one can vary the data on \mathcal{N}_0 while keeping the initial data on \mathcal{S}_0 to produce a wide class of Kerr-like solutions in which the *intrinsic* geometries of the horizon coincide with the geometry of Kerr but differ *off* the horizon.

VI. CONCLUSIONS

The intrinsic properties of the Kerr-Newman black hole are well understood and have been exhaustively investigated in the formalism of (weakly) isolated horizons. Despite the relatively high degree of symmetry, the Kerr-Newman solution adequately describes isolated black holes in equilibrium in the absence of matter outside the black hole thanks to its uniqueness properties. As we explained in the introduction, the formalism of WIHs allows one to generate a large class of solutions representing black holes deformed by external matter or fields which can be prescribed in an arbitrary way; in particular, the external fields are not restricted to be weak. In order to accomplish this program and analyze properties of

deformed black holes analytically, one first needs the description of the full Kerr-Newman metric not only the part intrinsic to the horizon in the WIH formalism.

In this paper, we explicitly constructed a Bondi-like tetrad for the Kerr-Newman black hole satisfying the properties imposed in [17] for a general WIH. In this tetrad, we were able to find the initial data given on the horizon and, more importantly, on the transversal null hypersurface \mathcal{N}_0 . In this sense we completed the description of Kerr-Newman solution in the framework of WIHs. Having the standard example of a WIH at hand, the next step is to consider variations of the initial data and understand their physical implications. This work is in progress.

ACKNOWLEDGMENTS

Work of M. S. was financially supported by the Grant No. GAČR 17-16260Y of the Czech Science Foundation. A. F. acknowledges the support from Grant No. GAUK 588217 of Charles University in Prague. N. G. and M. S. gratefully acknowledge support from the DFG within the Research Training Group 1620 Models of Gravity. Partial support of N. G. comes also from NewCompStar, COST Action MP1304. A. F. and M. S. acknowledge useful discussions with professor Pavel Krtouš and Dr. David Kofroň and the hospitality of institute ZARM, University of Bremen, Germany.

APPENDIX A: NEWMAN-PENROSE FORMALISM

For the sake of completeness, in this section we list all relevant definitions and relations of the Newman-Penrose (NP) formalism. In this paper, we follow the conventions of [19,24,25] adapted to the metric signature $(+ - - -)$, while also other conventions are common [39,40]. In particular, our conventions differ from [17].

The “NP tetrad” is a four-tuple of null vectors $(\ell^a, n^a, m^a, \bar{m}^a)$ normalized by relations

$$\ell^a n_a = -m^a \bar{m}_a = 1, \quad (A1)$$

where all other possible contractions vanish. Covariant derivatives in the directions of vectors forming the null tetrad are denoted by

$$D = \ell^a \nabla_a, \quad \Delta = n^a \nabla_a, \quad \delta = m^a \nabla_a, \quad \bar{\delta} = \bar{m}^a \nabla_a, \quad (A2)$$

where bar denotes the complex conjugate. In the NP formalism, the connection is encoded in twelve complex spin coefficients defined by

$$\begin{aligned}\kappa &= m^a D\ell_a, \quad \tau = m^a \Delta\ell_a, \quad \varepsilon = \frac{1}{2} [n^a D\ell_a - \bar{m}^a Dm_a], \\ \sigma &= m^a \delta\ell_a, \quad \varrho = m^a \bar{\delta}\ell_a, \quad \beta = \frac{1}{2} [n^a \delta\ell_a - \bar{m}^a \delta m_a], \\ \pi &= n^a D\bar{m}_a, \quad \nu = n^a \Delta\bar{m}_a, \quad \gamma = \frac{1}{2} [n^a \Delta\ell_a - \bar{m}^a \Delta m_a], \\ \lambda &= n^a \bar{\delta}\bar{m}_a, \quad \mu = n^a \delta\bar{m}_a, \quad \alpha = \frac{1}{2} [n^a \bar{\delta}\ell_a - \bar{m}^a \bar{\delta}m_a].\end{aligned}\tag{A3}$$

Some of the spin coefficients have direct geometrical meaning [17]. Namely, real and imaginary parts of ϱ determine the expansion and twist [41] of the congruence ℓ^a , respectively; similarly, real and imaginary parts of μ describe the expansion and twist of n^a [26]; coefficients σ and λ describe the shear of ℓ^a and n^a . Definitions (A3) imply “transport equations”

$$D\ell^a = (\varepsilon + \bar{\varepsilon})\ell^a - \bar{\kappa}m^a - \kappa\bar{m}^a, \tag{A4a}$$

$$\Delta n^a = -(\gamma + \bar{\gamma})n^a + \nu m^a + \bar{\nu}\bar{m}^a, \tag{A4b}$$

$$\Delta\ell^a = (\gamma + \bar{\gamma})\ell^a - \bar{\tau}m^a - \tau\bar{m}^a, \tag{A4c}$$

which show that κ and ν describe the deviation of ℓ^a and n^a from being geodesics, while $\varepsilon + \bar{\varepsilon}$ and $\gamma + \bar{\gamma}$ measure the failure of ℓ^a and n^a being affinely parametrized. Covariant derivative of m^a along n^a is given by the transport equation

$$\Delta m^a = \bar{\nu}\ell^a - \tau n^a + (\gamma - \bar{\gamma})m^a. \tag{A5}$$

Components of the Weyl tensor with respect to the null tetrad are provided by the Weyl scalars

$$\begin{aligned}\Psi_0 &= C_{abcd}\ell^a m^b \ell^c m^d, & \Psi_1 &= C_{abcd}\ell^a n^b \ell^c m^d, \\ \Psi_2 &= C_{abcd}\ell^a m^b \bar{m}^c n^d, & \Psi_3 &= C_{abcd}\ell^a n^b \bar{m}^c n^d, \\ \Psi_4 &= C_{abcd}\bar{m}^a n^b \bar{m}^c n^d.\end{aligned}\tag{A6}$$

In electrovacuum spacetimes, the components of the trace-free part of the Ricci tensor are given by $\Phi_{mn} = \phi_m \bar{\phi}_n$, where

$$\begin{aligned}\phi_0 &= F_{ab}\ell^a m^b, & \phi_2 &= F_{ab}\bar{m}^a n^b, \\ \phi_1 &= \frac{1}{2}F_{ab}[\ell^a n^b - m^a \bar{m}^b].\end{aligned}\tag{A7}$$

are the components of the electromagnetic tensor F_{ab} . The scalar curvature is $\Lambda = 0$.

The actual field equations are provided by the set of Ricci identities, Bianchi identities and commutation relations. The full list of these equations as well as the Maxwell equations in the NP formalism can be found, e.g., in [24,25]. In the present paper, we need only few of those

equations and, hence, we show them in the appropriate context only.

On the other hand, we employ all freedom available in the choice of the tetrad and for that we need the complete set of transformation equations for the NP quantities. A “boost” in the plane spanned by ℓ^a and n^a with real parameter A is defined as the transformation

$$\ell^a \mapsto A^2 \ell^a, \quad n^a \mapsto A^{-2} n^a, \quad m^a \mapsto m^a. \tag{A8}$$

Under boost, the spin coefficients (A3) transform according to the formulas

$$\begin{aligned}\kappa &\mapsto A^4 \kappa, & \tau &\mapsto \tau, & \sigma &\mapsto A^2 \sigma, & \varrho &\mapsto A^2 \varrho, \\ \pi &\mapsto \pi, & \nu &\mapsto A^{-4} \nu, & \mu &\mapsto A^{-2} \mu, & \lambda &\mapsto A^{-2} \lambda, \\ \varepsilon &\mapsto A^2 \varepsilon + ADA, & \gamma &\mapsto A^{-2} \gamma + A^{-3} \Delta A, \\ \beta &\mapsto \beta + A^{-1} \delta A, & \alpha &\mapsto \alpha + A^{-1} \bar{\delta} A,\end{aligned}\tag{A9}$$

while the Weyl scalars (A6) and electromagnetic scalars (A7) transform as

$$\Psi_m \mapsto A^{2(2-m)} \Psi_m, \quad m = 0, 1, 2, 3, 4, \tag{A10a}$$

$$\phi_m \mapsto A^{2(1-m)} \phi_m, \quad m = 0, 1, 2. \tag{A10b}$$

Any quantity η which transforms as $\eta \mapsto A^{2w} \eta$ is said to have a “boost weight” w .

The next transformation is the spin in the spacelike plane spanned by m^a and \bar{m}^a with a real parameter χ defined by

$$\hat{\ell}^a \mapsto \ell^a, \quad \hat{n}^a \mapsto n^a, \quad \hat{m}^a \mapsto e^{2i\chi} m^a. \tag{A11}$$

Under spin, the spin coefficients (A3) transform as

$$\begin{aligned}\kappa &\mapsto e^{2i\chi} \kappa, & \tau &\mapsto e^{2i\chi} \tau, & \sigma &\mapsto e^{4i\chi} \sigma, & \varrho &\mapsto \varrho, \\ \pi &\mapsto e^{-2i\chi} \pi, & \nu &\mapsto e^{-2i\chi} \nu, & \mu &\mapsto \mu, & \lambda &\mapsto e^{-4i\chi} \lambda, \\ \varepsilon &\mapsto \varepsilon + iD\chi, & \gamma &\mapsto \gamma + i\Delta\chi, \\ \beta &\mapsto e^{2i\chi}(\beta + i\delta\chi), & \alpha &\mapsto e^{-2i\chi}(\alpha + i\bar{\delta}\chi).\end{aligned}\tag{A12}$$

The Weyl scalars (A6) as well as the electromagnetic scalars (A7) are then given by

$$\Psi_m \mapsto e^{2(2-m)i\chi} \Psi_m, \quad m = 0, 1, 2, 3, 4, \tag{A13}$$

$$\phi_m \mapsto e^{2(1-m)i\chi} \phi_m, \quad m = 0, 1, 2. \tag{A14}$$

Again, a quantity η is said to have a “spin weight” s if it transforms like $\eta \mapsto e^{2is\chi}$ under the spin. The associated spin raising/lowering operators δ and $\bar{\delta}$ are defined by [25,42]

$$\delta\eta = \delta\eta + s(\bar{\alpha} - \beta)\eta, \quad \bar{\delta}\eta = \bar{\delta}\eta - s(\alpha - \bar{\beta})\eta. \quad (\text{A15})$$

Another kind of transformation of the null tetrad is a “null rotation” about ℓ^a with complex parameter c :

$$\begin{aligned} \ell^a &\mapsto \ell^a, & m^a &\mapsto m^a + \bar{c}\ell^a, \\ n^a &\mapsto n^a + cm^a + \bar{c}\bar{m}^a + |c|^2\ell^a, \end{aligned} \quad (\text{A16})$$

under which the spin coefficients (A3) transform as follows:

$$\begin{aligned} \kappa &\mapsto \kappa, \\ \tau &\mapsto \tau + c\sigma + \bar{c}\varrho + \kappa|c|^2, \\ \sigma &\mapsto \sigma + \kappa\bar{c}, \\ \varrho &\mapsto \varrho + \kappa c, \\ \varepsilon &\mapsto \varepsilon + c\kappa, \\ \gamma &\mapsto \gamma + c(\beta + \tau) + \alpha\bar{c} + \sigma c^2 + (\varepsilon + \varrho)|c|^2 + \kappa c^2\bar{c}, \\ \beta &\mapsto \beta + c\sigma + \varepsilon\bar{c} + \kappa|c|^2, \\ \alpha &\mapsto \alpha + c(\varepsilon + \varrho) + \kappa c^2, \\ \pi &\mapsto \pi + 2ce + c^2\kappa + Dc, \\ \nu &\mapsto \nu + c(2\gamma + \mu) + \bar{c}\lambda + c^2(2\beta + \tau) + c^3\sigma \\ &\quad + |c|^2(\pi + 2\alpha) + c^2\bar{c}(2\varepsilon + \varrho) + c^3\bar{c}\kappa \\ &\quad + |c|^2Dc + \Delta c + c\delta c + \bar{c}\bar{\delta}c, \\ \mu &\mapsto \mu + 2c\beta + \bar{c}\pi + c^2\sigma + 2|c|^2\varepsilon + c^2\bar{c}\kappa + \bar{c}Dc + \delta c, \\ \lambda &\mapsto \lambda + c(\pi + 2\alpha) + c^2(\varrho + 2\varepsilon) + \kappa c^3 + cDc + \bar{\delta}c, \end{aligned} \quad (\text{A17})$$

The transformation rules for the Weyl and electromagnetic scalars (A6) and (A7) read

$$\begin{aligned} \Psi_0 &\mapsto \Psi_0, \\ \Psi_1 &\mapsto \Psi_1 + c\Psi_0, \\ \Psi_2 &\mapsto \Psi_2 + 2c\Psi_1 + c^2\Psi_0, \\ \Psi_3 &\mapsto \Psi_3 + 3c\Psi_2 + 3c^2\Psi_1 + c^3\Psi_0, \\ \Psi_4 &\mapsto \Psi_4 + 4c\Psi_3 + 6c^2\Psi_2 + 4c^3\Psi_1 + c^4\Psi_0, \\ \phi_0 &\mapsto \phi_0, \\ \phi_1 &\mapsto \phi_1 + c\phi_0, \\ \phi_2 &\mapsto \phi_2 + 2c\phi_1 + c^2\phi_0. \end{aligned} \quad (\text{A18})$$

Finally, a “null rotation” about n^a with complex parameter d is defined by the relations

$$\begin{aligned} n^a &\mapsto n^a, & m^a &\mapsto m^a + dn^a, \\ \ell^a &\mapsto \ell^a + \bar{d}m^a + d\bar{m}^a + |d|^2n^a. \end{aligned} \quad (\text{A19})$$

The spin coefficients (A3) now transform as

$$\begin{aligned} \kappa &\mapsto \kappa + d(2\varepsilon + \varrho) + \bar{d}\sigma + d^2(\pi + 2\alpha) + d^3\lambda \\ &\quad + |d|^2(\tau + 2\beta) + d^2\bar{d}(2\gamma + \mu) + d^3\bar{d}\nu \\ &\quad - |d|^2\Delta d - Dd - d\bar{\delta}d - \bar{d}\delta d, \\ \tau &\mapsto \tau + 2d\gamma + d^2\nu - \Delta d, \\ \sigma &\mapsto \sigma + d(\tau + 2\beta) + d^2(\mu + 2\gamma) + d^3\nu - d\Delta d - \delta d, \\ \varrho &\mapsto \varrho + 2d\alpha + \bar{d}\tau + d^2\lambda + 2|d|^2\gamma + d^2\bar{d}\nu - \bar{d}\Delta d - \bar{\delta}d, \\ \varepsilon &\mapsto \varepsilon + d(\alpha + \pi) + \beta\bar{d} + \lambda d^2 + (\mu + \gamma)|d|^2 + \nu d^2\bar{d}, \\ \gamma &\mapsto \gamma + d\nu, \\ \beta &\mapsto \beta + d(\gamma + \mu) + d^2\nu, \\ \alpha &\mapsto \alpha + d\lambda + \bar{d}\gamma + |d|^2\nu, \\ \pi &\mapsto \pi + d\lambda + \bar{d}\mu + |d|^2\nu, \\ \nu &\mapsto \nu, \\ \mu &\mapsto \mu + d\nu, \\ \lambda &\mapsto \lambda + \nu\bar{d}. \end{aligned} \quad (\text{A20})$$

For the Weyl scalars (A6) and the electromagnetic scalars (A7) we now have

$$\begin{aligned} \Psi_0 &\mapsto \Psi_0 + 4d\Psi_1 + 6d^2\Psi_2 + 4d^3\Psi_3 + d^4\Psi_4, \\ \Psi_1 &\mapsto \Psi_1 + 3d\Psi_2 + 3d^2\Psi_3 + d^3\Psi_4, \\ \Psi_2 &\mapsto \Psi_2 + 2d\Psi_3 + d^2\Psi_4, \\ \Psi_3 &\mapsto \Psi_3 + d\Psi_4, \\ \Psi_4 &\mapsto \Psi_4, \\ \phi_0 &\mapsto \phi_0 + 2d\phi_1 + d^2\phi_2, \\ \phi_1 &\mapsto \phi_1 + d\phi_2, \\ \phi_2 &\mapsto \phi_2. \end{aligned} \quad (\text{A21})$$

APPENDIX B: VISUALIZATION

In this section, we briefly present the visualization of the differences between the standard twisting Kinnersley tetrad and the nontwisting tetrad constructed in this paper. Standard treatment of optical scalars for null geodesics in the NP formalism can be found, e.g., in [25]. Here we need to consider slightly more general case.

Consider a general NP tetrad ℓ^a , n^a , and m^a , for which ℓ^a is not necessarily a geodesic (which is the case of ℓ_B^a given by V) and m^a is not necessarily parallelly propagated along either ℓ^a or n^a [which is the case for the Kinnersley tetrad (4)]. We can always introduce complex null vectors ξ_ℓ^a and ξ_n^a for which $D\xi_\ell^a = 0$ and $\Delta\xi_n^a = 0$, respectively. Let z_ℓ^a and z_n^a be deviation vectors orthogonal to ℓ^a and n^a , respectively, which are propagated by equations

$$Dz_\ell^a = z_\ell^b \nabla_b \ell^a, \quad \Delta z_n^a = z_n^b \nabla_b n^a. \quad (\text{B1})$$

We expand both connecting vectors as

$$\begin{aligned} z_\ell^a &= c_\ell \ell^a - \bar{z}_\ell \xi_\ell^a - z_\ell \bar{\xi}_\ell^a, \\ z_n^a &= c_n n^a - z_n \xi_n^a - \bar{z}_n \bar{\xi}_n^a, \end{aligned} \quad (\text{B2})$$

and interpret the component $z_\ell = x_\ell + iy_\ell$ ($z_n = x_n + iy_n$) as complex coordinate of the projection of z_ℓ^a (z_n^a) onto the spacelike plane orthogonal to ℓ^a (n^a). Corresponding evolution equations read

$$\begin{aligned} Dz_\ell &= -\sigma z_\ell - \bar{\sigma} \bar{z}_\ell + \kappa c_\ell, \\ \Delta z_n &= \mu z_n + \lambda \bar{z}_n - \nu c_n, \end{aligned} \quad (\text{B3})$$

and

$$\begin{aligned} Dc_\ell &= (\pi - \alpha - \bar{\beta})z + (\bar{\pi} - \bar{\alpha} - \beta)\bar{z}, \\ \Delta c_n &= (\bar{\alpha} + \beta - \tau)z + (\alpha - \bar{\beta} - \bar{\tau})\bar{z}. \end{aligned} \quad (\text{B4})$$

In what follows we consider vectors ℓ_K^a and n_K^a of the Kinnersley tetrad and ℓ_B^a and n_B^a of the Bondi-like tetrad. In each case we choose the initial spacetime point with coordinates (r_0, θ_0) and a sequence $\{z_{0,n} = e^{2\pi in/N}\}_{n=0}^N$ of initial coordinates of the deviation vector. For each $z_{0,n}$ we solve the deviation equations (B3) along the null geodesic and plot points z_n for given values of the parameter along the geodesic, which we interpret as the




FIG. 2. Nontwisting, nonexpanding and shear-free congruence ℓ_K^a with the initial position $r_0 = 0$, $\theta_0 = \pi/4$.




FIG. 3. Twisting, expanding and shear-free congruence ℓ_K^a with the initial position $r_0 = 0.5$, $\theta_0 = \pi/4$.

cross-sections of the family of nearby geodesics with initially circular cross-section.

We choose the parameters of the Kerr-Newman spacetime $M = 1.2$, $a = 1.1$, $Q = 0.2$. In Fig. 2 we plot the congruence ℓ_K^a on the horizon. By the properties of a WIH, this congruence is nontwisting, nonexpanding and shear-free. Congruence ℓ_K^a off the horizon, see Fig. 3, is expanding, twisting and shear-free, i.e. the cross-sections remain circular. On the other hand, congruence n_K^a , being transversal to the horizon and future pointing, is converging and twisting, see Fig. 4. Congruence n_B^a of the Bondi-like tetrad is also converging but has zero twist and nonvanishing shear, as Fig. 5 demonstrates. Finally, optical scalars of the congruence ℓ_B^a vanish on the horizon, Fig. 2, but off the horizon all optical scalars are nonvanishing, see Fig. 6.




FIG. 4. Twisting, converging and shear-free congruence n_K^a for the initial position $r_0 = 3$, $\theta_0 = \pi/4$.




FIG. 5. Nontwisting, converging and shearing congruence n_B^a for the initial position $r_0 = 5$, $\theta_0 = \pi/4$.




FIG. 6. Twisting, expanding and shearing congruence ℓ_B^a for the initial position $r_0 = 2$, $\theta_0 = \pi/4$.

- [1] A. Ashtekar, C. Beetle, and S. Fairhurst, *Classical Quantum Gravity* **16**, L1 (1999).
- [2] A. Ashtekar, C. Beetle, and S. Fairhurst, *Classical Quantum Gravity* **17**, 253 (2000).
- [3] A. Ashtekar, C. Beetle, and J. Lewandowski, *Phys. Rev. D* **64** (2001).
- [4] A. Ashtekar, C. Beetle, and J. Lewandowski, *Classical Quantum Gravity* **19**, 1195 (2002).
- [5] A. Ashtekar, J. Baez, A. Corichi, and K. Krasnov, *Phys. Rev. Lett.* **80**, 904 (1998).
- [6] A. Ashtekar and B. Krishnan, *Living Rev. Relativ.* **7** (2004).
- [7] A. Ashtekar, S. Fairhurst, and B. Krishnan, *Phys. Rev. D* **62**, 104025 (2000).
- [8] A. Ashtekar, J. Engle, T. Pawłowski, and C. V. D. Broeck, *Classical Quantum Gravity* **21**, 2549 (2004).
- [9] R. Geroch, *J. Math. Phys.* **11**, 2580 (1970).
- [10] R. O. Hansen, *J. Math. Phys.* **15**, 46 (1974).
- [11] N. Gürlebeck, *Phys. Rev. D* **90**, 024041 (2014).
- [12] N. Gürlebeck, *Phys. Rev. Lett.* **114**, 151102 (2015).
- [13] E. T. Newman, E. Couch, K. Chinnapared, A. Exton, A. Prakash, and R. Torrence, *J. Math. Phys.* **6**, 918 (1965).
- [14] T. Johannsen, A. E. Broderick, P. M. Plewa, S. Chatzopoulos, S. S. Doeleman, F. Eisenhauer, V. L. Fish, R. Genzel, O. Gerhard, and M. D. Johnson, *Phys. Rev. Lett.* **116** (2016).
- [15] D. Psaltis, N. Wex, and M. Kramer, *Astrophys. J.* **818**, 121 (2016).
- [16] J. Lewandowski, *Classical Quantum Gravity* **17**, L53 (2000).
- [17] B. Krishnan, *Classical Quantum Gravity* **29**, 205006 (2012).
- [18] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, *Proc. R. Soc. A Math. Phys. Eng. Sci.* **269**, 21 (1962).
- [19] E. Newman and R. Penrose, *J. Math. Phys.* **3**, 566 (1962).
- [20] E. T. Newman and T. W. J. Unti, *J. Math. Phys.* **3**, 891 (1962).
- [21] N. Gürlebeck and M. Scholtz, *Phys. Rev. D* **95**, 064010 (2017).
- [22] J. Lewandowski and T. Pawłowski, *Int. J. Mod. Phys. D* **11**, 739 (2002).
- [23] S. J. Fletcher and A. W. C. Lun, *Classical Quantum Gravity* **20**, 4153 (2003).
- [24] R. Penrose and W. Rindler, *Spinors and Space-time* (Cambridge University Press, Cambridge, England, 1984), Vol. 1.
- [25] J. Stewart, *Advanced General Relativity* (Cambridge University Press, Cambridge, England, 1993).
- [26] B. Krishnan, in *Springer Handbook of Spacetime*, edited by A. Ashtekar and V. Petkov (Springer, New York, 2014) Chap. 25, p. 527.
- [27] W. Kinnersley, *J. Math. Phys.* **10**, 1195 (1969).
- [28] J. Jezierski and M. Łukasik, *Classical Quantum Gravity* **23**, 2895 (2006).
- [29] E. G. Kalnins, W. Miller, Jr, and G. C. Williams, *J. Math. Phys.* **30**, 2360 (1989).
- [30] Notice that, thanks to the properties of nonexpanding horizon, Lie dragging preserves the normalization of the vectors.
- [31] I. Rácz, *Classical Quantum Gravity* **24**, 5541 (2007).
- [32] I. Rácz, *Classical Quantum Gravity* **31**, 035006 (2014).
- [33] B. Carter, *Phys. Rev.* **174**, 1559 (1968).
- [34] M. Walker and R. Penrose, *Commun. Math. Phys.* **18**, 265 (1970).
- [35] That is, radial geodesics emanating from the equatorial plane will remain in this plane.
- [36] J. A. Marck, *Phys. Lett.* **97A**, 140 (1983).
- [37] D. Kubizňák, V. P. Frolov, P. Krtouš, and P. Connell, *Phys. Rev. D* **79**, 024018 (2009).
- [38] J. Lewandowski and T. Pawłowski, *Classical Quantum Gravity* **20**, 587 (2003).
- [39] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2009).
- [40] J. B. Griffiths and J. Podolský, *Exact Space-times in Einstein's General Relativity* (Cambridge University Press, Cambridge, England, 2009).
- [41] E. Poisson, *A Relativist's Toolkit: The Mathematics of Black-Hole Mechanics* (Cambridge University Press, Cambridge, England, 2004).
- [42] J. N. Goldberg, *J. Math. Phys.* **8**, 2155 (1967).