

# Lidar soundings of the middle atmosphere: Temperature, gravity waves and noctilucent clouds

Natalie Kaifler

Institute of Atmospheric Physics, German Aerospace  
Center, Oberpfaffenhofen, Germany



Maynooth University  
15 February 2019



# Outline

## 1. The middle atmosphere


- Thermal state
- Dynamics



## 2. Observations with lidar instruments

## 3. Gravity waves

- Life cycle
- Observation and analysis




## 4. Noctilucent clouds

- History
- Balloon campaign




# The middle atmosphere

- Mesosphere
  - ~50-90 km
  - Decreasing T
  - Meteors and noctilucent clouds
- Stratosphere
  - ~10-50 km
  - Increasing T due to ozone absorption



SuW-Grafik

# Thermal structure of the atmosphere




- Dependence on season/latitude
- The atmosphere is far from radiative equilibrium

→ dominated by dynamics






# Waves in Fluids



Lighthill, 1967



# Gravity waves




Kim et al., 2003




# Thermal structure and circulation of the atmosphere



## 2. Observations with lidar instruments




# Sounding of the atmosphere with lidar



# Lidar data

## Measurement




## Data Processing

- Atmosphere is in hydrostatic equilibrium
- Ideal gas law
- Temperature  $T_0$  at top of profile is „known“

Integration of photon count profiles from top  $z_0$  to bottom  $z$ :


$$T(z) = \frac{S(z_0)}{S(z)} T_0 + \frac{M}{k_B} \int_{z_0}^z \frac{S(z')}{S(z)} g(z') dz$$

## Data Product




# From temperatures to waves

- Temperature




- Temperature perturbation  
= Gravity waves



Vertical filter



## Lidar technical description



TELMA, New Zealand, 2014



- 14 W laser power, 532 nm, 100 Hz
- 63 cm telescope, 200  $\mu$ rad FOV
- Multiple Rayleigh and Raman detectors
- Mobile lidar systems (11- and 8-foot container) of 1500 kg
- 2 kW power consumption




# DLR middle atmosphere lidars




# Thermal structure, Rio Grande, 54°S


Laser replacement:  
14 W



### 3. Gravity waves



# Gravity wave activity above Rio Grande, Argentina






- Extreme wave events in winter
- Strongly intermittent

Black symbols: observation period  $>3$  hours; grey symbols  $<3$  hours



# Mountain waves

- You need:
  - Strong winds
  - Perpendicular mountain range




# Extreme mountain wave event


- 80 K peak-to-peak amplitude (much more than previous observations)

- Long duration of several days

CORAL lidar observations at Rio Grande




Kaifler et al., in preparation for GRL




# Propagation within the polar vortex edge

- Generation and propagation inside the polar vortex edge
- profound disturbance of stratospheric circulation for several 1000 km






Temperature Perturbations (K) at 1 hPa  
Valid: 20180621 02 UTC



## Lidar and OH imager



- Strong mesospheric, propagating GW
- GW breaking generating small-scale waves




# Wavelet analysis

→ Gravity wave parameters


- e.g. 40 min period, 100 km horizontal wavelength, propagating north-west



## 4. Noctilucent clouds



## 4. Noctilucent clouds



- Nucleation around the mesopause
- Sedimentation in Earth's gravity field
- Growth to  $\sim 50$  nm
- Visible as NLC
- Sublimation at lower boundary



# History

- Discovery in 1885

July 16, 1885]

NATURE

be so well plotted on this last survey that the amount of gorge excavated since 1750 should be knowable to an acre. The west fall, then, only slightly the larger, has ever since been widening, lowering its edge, and getting more of the stream; so that the east one, comparatively stationary, retaining its height and decreasing in volume, must dry up, and its bed and all the isles become part of New York State.

E. L. GARRETT

July 11

## Sky Glows

EVER since the sunsets of 1883 and last year there has been at times an abnormal glare both before and after sundown. But I have seen nothing in the way of twilight effect so strange as that of Monday evening, the 6th, when about 10 p.m. a sea of luminous silvery white cloud lay above a belt of ordinary clear twilight sky, which was rather low in tone and colour. These clouds were wave-like in form, and evidently at a great elevation, and though they must have received their light from the sun, it was not easy to think so, as upon the dark sky they looked brighter and paler than clouds under a full moon. A friend who was with me aptly compared the light on these clouds to that which shines from white phosphor paint. This effect lasted for some time after 10 p.m., and extended from west to north, the lower edge of the clouds, which was sharply defined, was about  $12^{\circ}$  above the horizon.

ROBT. C. LESLIE

6, Moira Place, Southampton, July 8

well co  
now ref  
have al  
bility o  
change

Amor  
Astronc  
of Gree  
date an  
U.S. N  
order of  
U.S. N  
January  
Newcon  
*Almana*  
which w



"(1) T  
be practi  
everywhe

"(2) T  
longitude

" The  
the meth  
to us fro  
nomers.  
which the  
This pra




## Viewing geometry






## Satellite era

- Nadir-looking high-res CIPS camera on NASA's AIM satellite

→ small-scale structures:  
gravity waves



# Observation from a balloon platform!






# PMC-Turbo launch from Sweden




## NLC observation by lidars


- Laser (5 W, 532 nm, 100 Hz), optics and electronics inside pressure vessel
- 50 cm telescope for receiving backscattered photons




# NLC lidar soundings



# Wide-field images



video




## Narrow FOV images

- Turbulence and instability dynamics in high-resolution NLC images:

→ KHI  
→ vortices  
→ intrusions  
→ cusps

„the best place in the solar system to study turbulence“ – D.C. Fritts



# Summary

