


Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

## Autonomic Neuroscience: Basic and Clinical

journal homepage: [www.elsevier.com/locate/autneu](http://www.elsevier.com/locate/autneu)

## Recommendations for tilt table testing and other provocative cardiovascular autonomic tests in conditions that may cause transient loss of consciousness : Consensus statement of the European Federation of Autonomic Societies (EFAS) endorsed by the American Autonomic Society (AAS) and the European Academy of Neurology (EAN)<sup>☆,☆☆</sup>

Roland D. Thijs<sup>a,b,\*</sup>, Michele Brignole<sup>c,d</sup>, Cristian Falup-Pecurariu<sup>e</sup>, Alessandra Fanciulli<sup>f</sup>, Roy Freeman<sup>g</sup>, Pietro Guaraldi<sup>h</sup>, Jens Jordan<sup>i,j,k</sup>, Mario Habek<sup>l</sup>, Max Hilz<sup>m,n</sup>, Anne Pavy-LeTraon<sup>o</sup>, Iva Stankovic<sup>p</sup>, Walter Struhal<sup>q</sup>, Richard Sutton<sup>r</sup>, Gregor Wenning<sup>f</sup>, J. Gert van Dijk<sup>a</sup>

<sup>a</sup> Department of Neurology, Leiden University Medical Centre, Leiden, the Netherlands

<sup>b</sup> Stichting Epilepsie Instellingen Nederland (SEIN), Heemstede, the Netherlands

<sup>c</sup> Faint & Fall Programme, Department of Cardiology, Ospedale San Luca, IRCCS Istituto Auxologico Italiano, Milan, Italy

<sup>d</sup> Department of Cardiology and Arrhythmologic Centre, Ospedali del Tigullio, 16033 Lavagna, Italy

<sup>e</sup> Department of Neurology, County Emergency Clinic Hospital, Transilvania University, Brasov, Romania

<sup>f</sup> Department of Neurology, Medical University of Innsbruck, Innsbruck, Austria

<sup>g</sup> Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

<sup>h</sup> IRCCS Istituto delle Scienze Neurologiche di Bologna, Bologna, Italy

<sup>i</sup> German Aerospace Center (DLR), Institute of Aerospace Medicine, Cologne, Germany

<sup>j</sup> Chair of Aerospace Medicine, University of Cologne, Cologne, Germany

<sup>k</sup> University Hypertension Center, Cologne, Germany

<sup>l</sup> Referral Center for Autonomic Nervous System, Department of Neurology, University Hospital Center Zagreb, University of Zagreb, School of Medicine, Kispaticeva 12, HR-10000 Zagreb, Croatia

<sup>m</sup> Department of Neurology, University Erlangen-Nuremberg, Germany

<sup>n</sup> Dept. of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

<sup>o</sup> French reference center for MSA, Neurology department, University Hospital of Toulouse and INSERM U 1048, Toulouse, France

<sup>p</sup> Neurology Clinic, Clinical Center of Serbia, University of Belgrade, Belgrade, Serbia

<sup>q</sup> Department of Neurology, University Clinic Tulln, Karl Landsteiner University of Health Sciences, Tulln, Austria

<sup>r</sup> Department of Cardiology, National Heart & Lung Institute, Hammersmith Hospital, Duane Road, London W12 0NN, UK

## ARTICLE INFO

## Keywords:

Transient loss of consciousness  
Syncope  
Tilt table testing  
Vasovagal  
Reflex syncope  
Orthostatic hypotension  
Psychogenic pseudosyncope

## ABSTRACT

An expert committee was formed to reach consensus on the use of Tilt Table Testing (TTT) in the diagnosis of disorders that may cause transient loss of consciousness (TLOC) and to outline when other provocative cardiovascular autonomic tests are needed. While TTT adds to history taking, it cannot be a substitute for it. An abnormal TTT result is most meaningful if the provoked event is recognised by patients or eyewitnesses as similar to spontaneous ones. The minimum requirements to perform TTT are a tilt table, a continuous beat-to-beat blood pressure monitor, at least one ECG lead, protocols for the indications stated below and trained staff. This basic equipment lends itself to perform (1) additional provocation tests, such as the active standing test carotid sinus massage and autonomic function tests; (2) additional measurements, such as video, EEG, transcranial Doppler,

<sup>☆</sup> Consensus statement of the European Federation of Autonomic Societies (EFAS) endorsed by the American Autonomic Society (AAS) and the European Academy of Neurology (EAN). <sup>☆☆</sup> This article is co-published in the journals "Clinical Autonomic Research" and "Autonomic Neuroscience". <https://doi.org/10.1007/s10286-020-00738-6> and <https://doi.org/10.1016/j.autneu.2021.102792>. These articles are identical except for minor stylistic and spelling differences in keeping with each journal's style.

\* Corresponding author at: Department of Neurology, Leiden University Medical Centre, Leiden, the Netherlands.

E-mail address: [R.D.Thijs@lumc.nl](mailto:R.D.Thijs@lumc.nl) (R.D. Thijs).


<https://doi.org/10.1016/j.autneu.2021.102792>

Available online 19 March 2021

1566-0702/© 2021 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).


NIRS, end-tidal CO<sub>2</sub> or neuro-endocrine tests; (3) tailor-made provocation procedures in those with a specific and consistent trigger of TLOC.

TTT and other provocative cardiovascular autonomic tests are indicated if the initial evaluation does not yield a definite or highly likely diagnosis, but raises a suspicion of (1) reflex syncope, (2) the three forms of orthostatic hypotension (OH), i.e. initial, classic and delayed OH, as well as delayed orthostatic blood pressure recovery, (3) postural orthostatic tachycardia syndrome or (4) psychogenic pseudosyncope. A therapeutic indication for TTT is to teach patients with reflex syncope and OH to recognise hypotensive symptoms and to perform physical counter manoeuvres.



**Fig. 1.** Flow chart of the diagnostic-work-up following the initial evaluation of transient loss of consciousness (TLOC), i.e. history taking, ECG and bedside supine & standing blood pressure measurements.

Abbreviations: AST = active standing test with continuous blood pressure measurements, BP = blood pressure, cOH = classic orthostatic hypotension, CSM = carotid sinus massage, TLOC = transient loss of consciousness, TTT = tilt table testing, VVS = vasovagal syncope.



**Fig. 2.** Flow chart of the diagnostic-work-up of orthostatic intolerance, i.e. history taking and supine & standing blood pressure measurements. Abbreviations: AST = active standing test with continuous blood pressure measurements, classic OH = classical orthostatic hypotension, POTS = postural orthostatic tachycardia syndrome, TTT = tilt table testing.

## 1. Introduction

Tilt Table Testing (TTT), while initially developed to study physiological compensatory responses to orthostatic stress, proved useful as a diagnostic test for vasovagal syncope (VVS) in 1986 (Kenny et al., 1986). TTT is now widely used in clinical practice with a variety of protocols, variants and extensions. On 6 July 2018, the European Federation of Autonomic Societies (EFAS) organised a round table discussion to obtain consensus on the use of TTT in the diagnosis of disorders that may cause transient loss of consciousness (TLOC) and to outline when other provocative cardiovascular autonomic tests are needed. The 2018 guidelines of the European Society of Cardiology (Brignole et al., 2018a) and the 2011 EFAS/AAS Consensus paper (Freeman et al., 2011) were used as starting points. Definitions of various conditions and tests are used as presented in these sources; where necessary preference was given to the more recent ESC guidelines. Two authors (RDT and JGvD) searched PubMed for additional articles from 2014, with the keywords “tilt table” AND “syncope” OR “orthostatic intolerance” (449 hits). We also cite occasional earlier articles and reviews, when these were relevant. The resulting Consensus Statement was reviewed by all participants, discussed at a second session on 30 June 2019, a third session on 27 January 2020 and sent for endorsement to the autonomic nervous system (ANS) panel of the European Academy of Neurology (EAN). This Consensus Statement reflects the opinion of experts in the field, but is

not a formal evidence-based clinical guideline.

## 2. Indications

TTT and other provocative cardiovascular autonomic tests primarily aim to obtain a pathophysiological correlate for orthostatic intolerance and TLOC. The fundamental tool for the differential diagnosis is history taking of patients and eyewitnesses (Brignole et al., 2018a; Wieling et al., 2015a). TTT may provide an important addition to history taking, if the initial evaluation does not yield a definite or highly likely diagnosis. TTT should neither be used as a substitute for history taking nor isolated from history taking. Fig. 1 shows a pragmatic approach, beginning with the initial evaluation of TLOC including history taking, ECG and bedside orthostatic blood pressure measurements. Fig. 2 provides a flow chart for the work-up of patients with orthostatic intolerance.

### 2.1. To obtain a pathophysiological correlate for orthostatic intolerance and TLOC

The primary aim of TTT is to provoke an event with complaint recognition and to demonstrate the pathophysiological correlate (Brignole et al., 2018a; Saal et al., 2016). These aspects are both crucial: recognition may concern subjective sensations reported by patients as

well as visible aspects, such as changes in facial colour or movements, recognised by eyewitnesses as similar to spontaneous ones. Together with demonstration of pathophysiological measurements, a clinical-pathophysiological correlate is obtained, thereby proving the cause of TLOC. For most indications, TTT relies on prolonged orthostatic stress in the near-vertical position ('head-up tilt'). TTT is not useful for all TLOC forms, as epileptic seizures and cardiac syncope are not commonly provoked by the upright posture. In contrast, TTT is useful to diagnose syncope forms with an orthostatic component, i.e. reflex syncope and syncope due to orthostatic hypotension. TTT is also useful to provoke psychogenic TLOC, i.e. short-lasting apparent unconsciousness due to conversion. TTT can provide a definite diagnosis of psychogenic TLOC by excluding epilepsy or syncope. In the elderly it may be difficult to distinguish between falls with and without loss of consciousness. Hence, TTT may be used to investigate unexplained falls in the elderly, as these may be due to syncope (Heitterachi et al., 2002; Menant et al., 2016).

The role of TTT in diagnosing vasovagal syncope (VVS) has been debated (Sheldon, 2005; Kulkarni et al., 2020; Sutton et al., 2020; Shen et al., 2017). The yield of a certain or highly likely diagnosis with history taking following an initial evaluation by hospital physicians and long term follow up as a reference may amount to 60% with an accuracy of about 90% (van Dijk et al., 2008). The diagnostic yield can increase to as much as 85% with additional history taking by an expert in syncope (Sutton et al., 2014). In view of these findings TTT is not needed in the majority of cases presenting with syncope. Another concern is the specificity of TTT. Syncope during TTT reflects a tendency towards hypotension due to low preload in the upright position, rather than a specific diagnosis (Sutton and Brignole, 2014). TTT does not only modulate the occurrence of orthostatic VVS, but also of other forms of syncope such as cardiac syncope (Sutton and Brignole, 2014). TTT should therefore only be performed after a careful and detailed medical history and examination. The results of the test should be interpreted in the context of that history and examination. Estimates of sensitivity (overall 59%; 95 CI% 53–64%; range 21–72%) and specificity (overall 91%; 95 CI% 87–93%) of TTT in diagnosing VVS differ (Forleo et al., 2013). Important factors contributing to the variability relate to the specific test protocol (e.g. the use of pharmacological provocation, tilt angle, duration of TTT etc.), interpretation of the results, meaning whether complaint recognition or pathophysiological measurements are used as the standard to judge abnormality and the lack of an agreed reference standard other than long term follow-up (Brignole et al., 2018a; Saal et al., 2016; Forleo et al., 2013; Cheshire Jr. and Goldstein, 2019). The most marked contrasts are seen between 'passive' TTT, i.e. upright tilt without pharmacological provocation (sensitivity 37%; (95% CI 29–46%); specificity 99% (95CI% 97–99%)) vs. 'active' protocols (e.g. nitroglycerin provocation sensitivity 66% (95% CI 60–72%); specificity 89% (95% CI 84–92%), Forleo et al., 2013). As TTT relies predominantly on the orthostatic, literally meaning 'upright', position, TTT indications comprise the conditions bundled under 'orthostatic intolerance' (Brignole et al., 2018a; Saal et al., 2016; Cheshire Jr. and Goldstein, 2019). Orthostatic intolerance includes all three forms of OH (i.e. initial, classic and delayed OH), delayed orthostatic BP recovery, orthostatic VVS and the postural orthostatic tachycardia syndrome (POTS). Although some forms of reflex syncope, e.g. emotional VVS and cough syncope, primarily rely on other triggers, subjects with these forms are often also susceptible to orthostatic VVS (Humm and Z'Graggen, 2015; Mereu et al., 2016). TTT may therefore still help to provoke syncope in these cases. In other cases, specific additional provocations may be needed to provoke TLOC.

The role of TTT and other provocative cardiovascular autonomic tests is similar for the three forms of orthostatic hypotension (OH), i.e., the initial, classic and delayed ones, in that a definite diagnosis can be made when complaint recognition coincides with the type of BP decrease that is specific for each form (Table 1). The three forms may, however, require specific test protocols: (1) initial OH is commonly not provoked by passive tilt but requires an active standing test with

**Table 1**

Hemodynamic criteria for conditions causing orthostatic intolerance (Freeman et al., 2011; van Wijnen et al., 2017).

TTT/AST indication	Fall in SBP upon standing	Fall in DBP upon standing	Increase in HR upon standing	Timing
Initial OH	> 40 mmHg	> 20 mmHg	Not specified	Transient BP fall within 15 s upon standing.
Classic OH <sup>a,b</sup>	≥ 20 mmHg <sup>f</sup>	≥ 10 mmHg	Not specified <sup>f</sup>	Sustained BP fall within 3 min standing
Delayed OH <sup>a</sup>	≥ 20 mmHg	≥ 10 mmHg	Not specified	Sustained BP fall >3 min standing
POTS <sup>c</sup>	SBP fall not meeting OH criteria	DBP fall not meeting OH criteria	> 30 bpm <sup>d</sup> or > 120 bpm	Sustained HR increase within 10 min standing
Vasovagal presyncope	No formal criteria <sup>g</sup>	No formal criteria <sup>g</sup>	No formal criteria <sup>g</sup>	No formal criteria <sup>g</sup>
Delayed orthostatic BP recovery	Inability of SBP to recover to supine values within 30s of standing. Standing SBP should be ≥20 mmHg lower than supine values but not meet criteria of classic or initial OH.	Not meeting initial OH/classic OH criteria	Not specified	BP fall within 30 s upon standing

Abbreviations: AST = active standing test; classic OH = classical orthostatic hypotension, DBP = diastolic blood pressure, delayed OH = delayed orthostatic hypotension, HR = heart rate, initial OH = initial orthostatic hypotension, POTS = postural orthostatic tachycardia syndrome, SBP = systolic blood pressure, TTT = tilt table test.

<sup>a</sup> BP fall should be sustained to avoid confusion with transient BP falls seen in iOH or VVS.

<sup>b</sup> Please note that the ESC guidelines (Brignole et al., 2018a) added an additional (optional) criterion of SBP < 90 mmHg as these values are strongly associated with the occurrence of symptoms. An abnormal orthostatic fall in DBP without an abnormal fall in SBP is rare among patients with syncope and orthostatic intolerance (Fedorowski et al., 2017).

<sup>c</sup> A ΔHR/ΔSBP ratio < 0.5 bpm/mmHg argues for neurogenic OH (Norcliffe-Kaufmann et al., 2018a).

<sup>d</sup> For individuals aged 12–19 years the required increment is >40 bpm.

<sup>e</sup> POTS is a clinical diagnosis and requires symptoms of orthostatic intolerance and the documentation of an exaggerated postural tachycardia using TTT or an active standing test.

<sup>f</sup> For patients with supine hypertension, a reduction in SBP ≥ 30 mmHg is required, as the magnitude of the orthostatic BP fall is dependent on the baseline BP.

<sup>g</sup> No formal criteria to differentiate between vasovagal (pre)syncope and other causes of presyncope. The hemodynamic patterns associated with vasovagal (pre)syncope are heterogeneous (Brignole et al., 2018b) and may or may not be accompanied by HR slowing. Various criteria have been proposed to differentiate between subtypes.

continuous BP measurements (Finucane et al., 2019; Wieling et al., 2007); (2) classic OH is most commonly screened at bedside with intermittent BP measurements, but can also be assessed with TTT or an active standing test (Gibbons et al., 2017); delayed OH may require prolongation of the test protocol to allow a blood pressure decrease to occur (Gibbons et al., 2017; Gibbons and Freeman, 2006; Gibbons and Freeman, 2015). Note that classic OH and presumably delayed OH often fluctuate in severity, so the absence of complaints and a blood pressure decrease during the test do not exclude OH (Table 2) (Brignole et al., 2018a; Gibbons et al., 2017). Conversely, the finding of classic OH does not automatically provide an explanation for orthostatic intolerance or

**Table 2**

Association of orthostatic intolerance and orthostatic hypotension. Reproduced with permission from the 2018 ESC Guidelines for the diagnosis and management of syncope (Brignole et al., 2018a).

		History of syncope and orthostatic complaints	
		Highly suggestive of OH: syncope and pre-syncope are present during standing, absent while lying, and less severe or absent while sitting; a prediction for the morning; sitting or lying down must help; complaints may get worse immediately after exercise, after meals or in high temperature; no "automatic activation"	Possibly due to OH: not all of the features highly suggestive of OH are present
Supine and standing BP measurement	Symptomatic abnormal BP fall	Syncope is due to OH (Class I)	Syncope is likely due to OH (Class IIa)
	Asymptomatic abnormal BP fall	Syncope is likely due to OH (Class IIa)	Syncope may be due to OH (Class IIb)
	No abnormal BP drop	Unproven	Unproven

Abbreviations: BP= blood pressure, OH= orthostatic hypotension.

TLOC, as it is a common finding affecting up to one in five community-dwelling elderly (Brignole et al., 2018a; Saedon et al., 2020). When there is such a discrepancy between measurement results and complaints, the question whether the measured OH represents a clinically relevant finding critically depends on the clinical presentation. When doubt remains, home measurements during complaints may settle the issue.

A diagnosis of psychogenic TLOC preferably requires documentation of an event, for which video or video-EEG recordings provide the most convincing evidence. History taking often raises a strong suspicion of psychogenic TLOC, but may not always be reliable enough to rule out other causes of TLOC (LaFrance Jr. et al., 2013). The use of video-EEG is particularly important to exclude epilepsy when the clinical event is accompanied by positive motor phenomena (LaFrance Jr. et al., 2013). Note that a normal EEG during a provoked event excludes syncope with certainty, but need not exclude all possible forms of epilepsy. It does, however, exclude those forms of epilepsy that present with TLOC, i.e. tonic, clonic and tonic-clonic seizures (van Dijk et al., 2009). A diagnosis of psychogenic TLOC can also be established on clinical grounds if a clinician witnesses the event and observes features favouring psychogenic TLOC (e.g. eye closure, resisted eye-opening, partial responsiveness during the event etc.) (LaFrance Jr. et al., 2013).

## 2.2. To classify classic OH

TTT and other autonomic tests (active standing test, Valsalva manoeuvre and deep breathing) may help to distinguish between neurogenic and non-neurogenic causes for classic OH and to identify sympathetic or parasympathetic dysfunction in those with neurogenic OH (Brignole et al., 2018a; Gibbons et al., 2017; Norcliffe-Kaufmann et al., 2018a; Hilz and Dutsch, 2006). A blunted heart rate increase during classic OH makes a neurogenic cause more likely. A  $\Delta\text{HR}/\Delta\text{SBP}$  ratio  $< 0.5$  bpm/mmHg argues for neurogenic OH (sensitivity 91%; specificity 88% AUC = 0.96) (Norcliffe-Kaufmann et al., 2018a). It should be noted, however, that cardiac disorders (e.g., pacemaker, arrhythmias etc.) or certain drugs may preclude heart rate augmentation, thus an ECG, cardiac history and a medication review should be part of the evaluation of classic OH (Gibbons et al., 2017). Specialised autonomic function tests including responses to deep breathing and Valsalva manoeuvre may also help to diagnose neurogenic OH as well as to identify sympathetic or parasympathetic dysfunction in those with neurogenic OH (Brignole et al., 2018a; Gibbons et al., 2017; Hilz and Dutsch, 2006; Goldstein and Cheshire Jr., 2017; Brignole et al., 2018b; Low, 2003). The absence of a blood pressure overshoot and the absence of a heart rate increase during the Valsalva manoeuvre is

pathognomonic for neurogenic classic OH. There is also consensus that a blunted or abolished heart rate variation during deep breathing is suggestive of neurogenic classic OH. Interpretation of these findings requires age-adjusted, normative values (Low, 2003).

## 2.3. Treatment

TTT is not completely restricted to diagnosis. Patients may exhibit a decrease in syncope frequency after TTT, which may be due to patients having learned to recognise early signs of syncope, allowing them to take measures to prevent it (Sheldon et al., 2007). TTT may easily be expanded with a biofeedback session to teach the 'counter manoeuvres' (van Dijk et al., 2006; Wieling et al., 2015b). This session can be performed when subjects are still symptomatic after syncope, thus allowing strong feedback regarding the effectiveness of these interventions. The session is also extremely important for patients' education, as patients can see their own blood pressure fluctuate and discover which manoeuvres decrease (e.g., squat-to-stand) and which manoeuvres increase (e.g., leg crossing) their blood pressure. Applying these measures using biofeedback reduces the recurrence risk in patients with reflex syncope by 39% compared to conventional treatment only, i.e. explanation and life-style advice (van Dijk et al., 2006). Patients with definite VVS and no need for additional testing could be referred to an animated video on physical counter manoeuvres (). TTT also allows to study the temporal relation between onset of asystole and TLOC in those with asystolic VVS and help to guide pacemaker recommendations. In one-third of cases of tilt-induced asystolic reflex syncope, asystole occurred too late to have been the primary cause of TLOC, thus making pacemaker implantation likely ineffective (Saal et al., 2017).

## 2.4. Recommendations

### 2.4.1. TTT should be considered

- To increase the probability of a diagnosis of reflex syncope

This holds for those in whom the initial evaluation, including history taking, ECG and supine and standing blood pressure measurements raised a suspicion but not a definite or highly likely (probability 80–100%) diagnosis, or in those in whom confirmation of a diagnosis is required for other purposes, such as convincing patients, parents or medicolegal requirements.

- To assess classic OH and delayed OH

The recommended screening test for classic OH is a bedside active standing test with conventional intermittent BP measurements. TTT or an active standing test with continuous BP measurements should be considered if the bedside test does not show classic OH while the symptoms suggest classic OH (Table 2) (Brignole et al., 2018a). TTT is particularly useful to detect delayed OH, i.e. those with a sustained blood pressure fall of the magnitude of classic OH, but occurring later than 3 min upon standing (Freeman et al., 2011; Gibbons and Freeman, 2006). TTT may also be used to diagnose classic OH in subjects for whom active standing is difficult or unlikely to succeed, such as frail elderly and patients with significant motor impairments.

- To support a clinical diagnosis of POTS (Freeman et al., 2011; Sheldon et al., 2015)

Documentation of an exaggerated postural tachycardia using TTT or an active standing test may support a clinical diagnosis of POTS.

- To differentiate between syncope with myoclonus ('convulsive' syncope) and tonic-clonic seizures (Shmueli et al., 2018).
- To discriminate between neurogenic and non-neurogenic classic OH

A blunted heart rate increase during classic OH makes a neurogenic cause more likely. A  $\Delta HR/\Delta SBP$  ratio  $< 0.5$  bpm/mmHg argues for neurogenic OH (Norcliffe-Kaufmann et al., 2018a).

- To differentiate between vasovagal syncope and psychogenic TLOC (Tannemaat et al., 2013; Blad et al., 2015).
- To study the timing between asystole and the onset of TLOC

Asystole in VVS may occur too late to have been the prime cause of TLOC, making pacemaker implantation likely ineffective (Saal et al., 2017).

- To teach patients with reflex syncope and orthostatic hypotension to recognise hypotensive symptoms and how to perform physical counter manoeuvres (van Dijk et al., 2006; Wieling et al., 2015b).

Counter manoeuvres are effective measures to counteract impending reflex syncope or syncope due to orthostatic hypotension. Applying these measures using biofeedback reduces recurrence risk in patients with reflex syncope by 39% compared to conventional treatment (i.e. explanation and life-style advice) (van Dijk et al., 2006).

2.4.2. TTT should not be used

- To evaluate the treatment of reflex syncope

TTT should not be used for the routine clinical evaluation of treatment efficacy, although it might have a role in the evaluation of treatment approaches for reflex syncope in the research context (Brignole et al., 2018a; Shen et al., 2017).




Fig. 3. Choice of type of tilt table testing (TTT) protocol.

# pharmacological provocation with sublingual trinitroglycerin (TNG) may be considered to increase the sensitivity for VVS but should be avoided in those for whom delayed OH is suspected.

Abbreviations: AST = active standing test with continuous blood pressure measurements, CSM = carotid sinus massage, OH = orthostatic hypotension, POTS = postural orthostatic tachycardia syndrome, PPS = psychogenic pseudo-syncope, TTT = tilt table testing.

**Table 3**  
TTT protocols and supplementary measurements for each TTT indication.

TTT indication	Duration of tilt <sup>a</sup>	Pharmacological provocation	Additional measurements
Orthostatic intolerance <sup>b</sup>	At least 10 min	Not recommended	<i>Optional</i> Video, EEG, Respiratory, TCD, Catecholamines
- Classic OH	- 3 min		
- POTS	- 10 min		
- Delayed OH	- Up to 40 min		
- Vasovagal presyncope	- Up to 45 min		
Transient loss of consciousness	Up to 45 min <sup>c</sup>	Optional	<i>Optional</i> Video, EEG, Respiratory, NIRS, TCD
- Reflex syncope			
- PPS			

Abbreviations: classic OH = classical orthostatic hypotension, delayed OH = delayed orthostatic hypotension, LBNP = lower body negative pressure, NIRS = near infrared spectroscopy, POTS = postural orthostatic tachycardia syndrome, PPS = psychogenic pseudosyncope, TCD = transcranial doppler.

<sup>a</sup> TTT should be terminated earlier in case a clinical endpoint (i.e. (pre)syncope) is reached.

<sup>b</sup> Please note that an active standing test may also be used to establish a diagnosis in patients with orthostatic intolerance (see Section 2 of the main manuscript).

<sup>c</sup> In case of pharmacological provocation (TNG) TTT may be shortened to 40 min (20 min before and 20 min after TNG administration).

- To exclude cardiac syncope

Syncope during TTT suggests the presence of a hypotensive susceptibility. This tendency is common and does not only modulate the occurrence of orthostatic VVS, but also of other forms of syncope such as cardiac syncope (Sutton and Brignole, 2014).

- To exclude VVS

The sensitivity of TTT, especially TTT without pharmacological provocation, is too low to exclude VVS (Sutton and Brignole, 2014; Forleo et al., 2013).

- To diagnose initial OH

Documentation of initial OH requires an active standing test with a beat-to-beat blood pressure monitor (see paragraph 5.2) (Finucane et al., 2019; Wieling et al., 2007; van Wijnen et al., 2017).

### 3. Indications for additional provocations and measurements

The committee recommends that additional provocations and measurements are tailored to the patient's needs. Fig. 3 provides guidance when these auxiliary measures could be considered. Recommendations for TTT protocols and supplementary measurements are summarised in Table 3 and detailed in the following section.

## 4. Equipment

### 4.1. Basic equipment

#### 4.1.1. Beat-to-beat blood pressure monitor

In the context of syncope blood pressure (BP) can change substantially in a few seconds, so it must be measured with an appropriate temporal resolution. Conventional intermittent BP measurements allow one measurement per minute at best and are therefore unsuitable for syncope assessment. Classic OH and delayed OH are, however, accompanied by sustained decrease in BP lasting minutes, and hence can be detected with a conventional intermittent BP measurements (Finucane

et al., 2019; Gibbons et al., 2017; Pavy-Le Traon et al., 2016). Delayed OH, however, often occurs after 10 min of passive standing (Gibbons and Freeman, 2006). TTT is therefore more appropriate to identify delayed OH. In frail elderly and those with major motor impairments, TTT may also be preferred over active standing to prevent falling. Several commercial devices are available for non-invasive beat-to-beat measurements of blood pressure and heart rate using finger photoplethysmography and the volume clamp method (Imholz et al., 1998). Additional software yields estimates of other circulatory parameters including stroke volume, cardiac output and total peripheral resistance; these may be used to quantify the relative contributions of vasodilation and cardioinhibition (van Dijk et al., 2020), but this is not required for routine clinical testing.

#### 4.1.2. ECG

At least one ECG channel is required to detect bradycardia or asystole and to identify its underlying mechanism (e.g. intermittent AV block) (Taal et al., 2003; Brignole et al., 2011).

#### 4.1.3. Tilt table

The tilt-down time should preferably be short for syncope, as longer times may lengthen circulatory arrest and hereby the duration of cerebral hypoperfusion (Zysko et al., 2016). The optimal time is not known. The panel recommends the tilt back period from 70 degrees to horizontal should be less than 15 s. The subject should be protected against falling with safety straps with at least one strap at the chest and one above the knee.

## 4.2. Equipment extensions

### 4.2.1. Video and EEG recordings

For reflex syncope, adding video to TTT has the advantage that key clinical events that usually last less than 20 s can be studied in detail after the fact. Adding EEG provides additional guidance in control over the tilt procedure in that slowing of the EEG should always result in tilting the patient back; in reverse, tilting back is not yet needed if the EEG is still normal. The EEG pattern consists of either slowing or a slow-flat-slow pattern; this provides information about the degree of cerebral perfusion (van Dijk et al., 2014; Shmuelly et al., 2018). Adding video and EEG has limited clinical value in OH because TTT in OH forms rarely induces syncope, and if complaints occur these usually concern presyncope only, not usually accompanied by EEG changes.

For psychogenic TLOC, video and EEG recordings during TTT are very important as they allow a definite diagnosis of psychogenic TLOC (Brignole et al., 2018a; Tannemaat et al., 2013; Blad et al., 2015) and to recognise mixed presentations of psychogenic pseudosyncope and VVS (Blad et al., 2015). A complete clinical identification of the presence of TLOC requires assessment of loss of responsiveness, abnormal motor control, amnesia for the period of apparent unconsciousness and a short duration (Brignole et al., 2018a; van Dijk et al., 2009). Adding video allows these phenomena to be assessed with certainty. The addition of EEG helps to confirm the presence of normal brain activity during psychogenic TLOC and hereby to exclude TLOC due to syncope (slowing and/or flattening) or epilepsy (epileptiform abnormalities). Video without EEG does allow a highly probable diagnosis of psychogenic pseudosyncope, i.e. attacks with loss of muscle tone but without jerking movements, by proving that the recorded event fulfils the clinical criteria of TLOC in the absence of hypotension (LaFrance Jr. et al., 2013; Tannemaat et al., 2013; Blad et al., 2015). If TLOC is accompanied by stiffening or jerks, however, additional EEG recordings are recommended to discriminate between PNES and epileptic seizures (LaFrance Jr. et al., 2013). A diagnosis of psychogenic TLOC can also be established on clinical grounds if a clinician witness observed TLOC and documented examination findings typically found in psychogenic TLOC (e.g. eye closure, resisted eye-opening, partial responsiveness during the event etc.) (LaFrance Jr. et al., 2013).

Adding video-EEG to TTT can be accomplished by using an EEG machine to record and store all signals, including the output from the beat-to-beat BP monitor. As this is easily accomplished in a neurological setting, the panel encourages adding video and EEG to most TTT procedures. The video should record facial expression to assess eye closure and skin colour and, preferably, the upper extremities to evaluate the presence of jerking movements.

#### 4.2.2. Respiratory recordings

Respiration can be recorded using strain gauges around the thorax or abdomen, blood oxygen saturation and end-tidal CO<sub>2</sub>. Of these, CO<sub>2</sub> recordings may offer most insight. CO<sub>2</sub> recordings are useful to detail the pathophysiological cascade as hypocapnia may precede reflex syncope (Norcliffe-Kaufmann et al., 2008). Adding CO<sub>2</sub> recordings to TTT may also be of clinical interest as it may help to identify POTS subtypes that are consequent upon postural hyperventilation (Stewart et al., 2018) or episodes of hyperventilation in those who report dizziness without concomitant hypotension (Naschitz et al., 2000).

#### 4.2.3. Blood sampling

Blood sampling of neuro-endocrine substances holds promise as a means to improve pathophysiological understanding (Gibbons et al., 2017; Goldstein and Cheshire, 2018; Goldstein et al., 2003; Benditt et al., 2012). Measuring catecholamines before and after head-up tilt offers insights into baroreflex-mediated sympathetic activation, mostly to help distinguish between causes of neurogenic OH such as pure autonomic failure, Parkinson's disease and multiple system atrophy and to classify POTS variants (Gibbons et al., 2017; Goldstein and Cheshire, 2018; Goldstein et al., 2003; Benditt et al., 2012; Raj, 2013). Catecholamine assessments may also help to predict conversion from pure autonomic failure to multiple system atrophy (Singer et al., 2017). The measurements require placement of an indwelling venous line and established protocols regarding the material (cooling, centrifugation and type of assay) and TTT as the catheter implies a needle to obtain venous or arterial access and may hereby provoke VVS. To avoid this, a longer period of supine rest is recommended before tilt up. The panel recommends restricting the use of catecholamine assays to those with a diagnosis of neurogenic OH where the clinical work-up fails to discriminate between pre and postganglionic causes.

#### 4.2.4. Transcranial Doppler (TCD)

TCD monitoring during TTT aids to assess alterations in cerebral blood flow velocity at a high temporal resolution. This is of interest to investigate the pathophysiology of syncope (Norcliffe-Kaufmann et al., 2018b), but was so far of no help in the differential diagnosis of orthostatic intolerance or TLOC. TCD monitoring has been advocated as an additional tool to help establish psychogenic pseudosyncope (Raj et al., 2014).

#### 4.2.5. Near-infrared spectroscopy (NIRS)

Near-infrared spectroscopy of the brain reflects the relative amount of deoxygenated blood in the scalp and brain. At present, the technique shows promise as it suggests that decreased brain perfusion may be identified prior to clinical manifestations (Bachus et al., 2018; Kharraziha et al., 2019).

### 5. TTT protocols

#### 5.1. Classic TTT

We first present classic TTT and then discuss all variants and extensions that can be performed using the same equipment. The choice of the TTT protocol will depend on the clinical presentation (Figs. 2 and 3).

##### 5.1.1. Indication reflex syncope

Several protocols for TTT have been reported with or without

pharmacological challenges (Forleo et al., 2013; Sutton, 2015). The available evidence suggests that 30–60 min is optimal for the diagnosis of VVS (Brignole et al., 2018a; Saal et al., 2016; Forleo et al., 2013). The panel did not reach consensus as to whether TTT should be performed with or without pharmacological provocation. While the majority of the panel preferred to perform TTT for 45 min duration without pharmacological provocation, there is evidence that pharmacological provocation increases sensitivity with a minimal effect on the number of false positive results (Forleo et al., 2013; Bartoletti et al., 1999). For pharmacological provocation, we advocate against the use of intravenous isoproterenol and instead recommend sublingual trinitroglycerin (TNG) (300 to 400 µg): TNG does not require cannulation and is easier and quicker to administer, while sensitivity and specificity of TNG and isoproterenol are similar (Forleo et al., 2013). TNG is usually administered in a fixed dose (0.3 or 0.4 mg) rather than a dose based on body mass. It is not known at which mass side effects become a limiting factor, but the panel discourages the use of TNG in those with a mass below 50 kg. Arguments favouring TNG provocation include the increased sensitivity for VVS (sensitivity 66% (95% CI 60–72%) vs. passive TTT (37%; (95% CI 29–46%)) (Forleo et al., 2013). TNG may also, albeit to a lesser extent, impact specificity. The lower specificity of the TNG protocols 89% (95% CI 84–92%) vs. passive 99% (95% CI 97–99%) requires a critical evaluation whether the provoked event resembles the spontaneous one. Another drawback of TNG provocation is that it may provoke delayed OH. This is of particular concern in those with orthostatic (pre) syncope without specific clues for VVS or delayed OH (e.g. orthostatic (pre)syncope without autonomic activation in an elderly subject). Another factor to consider is the health care setting, as in some countries application of TNG for diagnostic purposes may be restricted or require a specific clinical environment.

Lower body negative pressure (LBPN) is an alternative non-pharmacological measure to provoke syncope during TTT. While LBPN has been studied extensively to assess haemodynamic responses to orthostatic stress, little is known of the validity of LBPN to detect VVS in patients with unexplained TLOC (El-Bedawi and Hainsworth, 1994; Goswami et al., 2019). This does not mean that LBPN cannot be useful in this context, but its clinical value has not yet been proven.

- Supine rest phase of at least 5 min with reliable and constant BP and HR values as a baseline.
- Head-up tilt angle between 60° and 80°, measured from the horizontal (Forleo et al., 2013). At 60°, 70° and 80° the force component pulling blood towards nether parts of the body presents 0.87, 0.94 and 0.98 of that of the fully vertical position. The positive yield of TTT seems optimal at 70° as the sensitivity is higher compared with TTT at 60° and 80° (Forleo et al., 2013).
- TTT without pharmacological provocation: head-up tilt for 45 min
- TTT with pharmacological provocation: start with 20 min head-up tilt, followed by administration of TNG in the tilted position and continuation for another 20 min
- TTT can be terminated when clinical events occur or, in case no symptoms are provoked, when the end of the recording period is reached. Which symptoms can serve as an endpoint will critically depend on the clinical context. Presyncope may be used as an endpoint in cases with a high pre-test likelihood of reflex syncope; patients may then be tilted back when recognised complaints are accompanied by hypotension. When the a priori probability is low, or when syncope with myoclonus needs to be distinguished from a convulsive seizure, syncope should be the endpoint. The decision to tilt back should be taken by someone with direct access to the patient and all recorded signals. The presence of asystole or EEG slowing, if recorded, is always a reason to tilt back immediately.
- A key element of TTT is to ask patients for recognition of symptoms. Eyewitnesses may confirm whether the provoked event resembled the spontaneous episodes, which may help to reduce false positive results.

### 5.1.2. Indication OH

Pharmacological provocation should not be used for any form of OH. Initial OH is not a reason for a TTT, but requires an active standing test with continuous blood pressure measurements. TTT may be used for those with orthostatic intolerance (1) when the active standing test does not reveal classic OH, (2) when delayed OH is suspected at history taking or (3) when a complaint correlation is needed. The TTT duration may be tailored to individual needs. Syncope occurs rarely in classic or delayed OH during TTT and should not be the primary endpoint.

- Supine rest phase before head-up tilt of at least 5 min with reliable and constant BP and HR values as a baseline. Supine measurements can be used to assess neurogenic supine hypertension (Fanciulli et al., 2018).
- Head-up tilt angle between 60° and 80°, measured from the horizontal.
- Head-up tilt of 3 min would suffice to establish a diagnosis of classic OH but longer TTT may be needed to rule out other causes of OH (e.g. delayed OH, POTS). The diagnosis of delayed OH may require a head-up tilt of up to 45 min (Gibbons and Freeman, 2006; Gibbons and Freeman, 2015). Patients may be tilted back when a complaint correlation is obtained, when syncope occurs, or when patients can no longer withstand the head-up position.
- Complaint recognition: see above. It is important to note that a mean standing BP <75 mmHg had an optimal sensitivity (97%) and specificity (98%) for detecting symptomatic OH in patients with Parkinson's disease (Palma et al., 2015). The symptom threshold may, however, vary between causes of OH. In a study of patients with diverse causes of OH, many patients were asymptomatic despite substantial SBP falls and low orthostatic blood pressures, suggesting that symptoms may not always be a reliable indicator of impaired cerebral perfusion or risk of syncope (Freeman et al., 2020).

### 5.1.3. Indication psychogenic TLOC

For psychogenic TLOC, pharmacological provocation may be used as for reflex syncope for two reasons: firstly, VVS occurs more often in psychogenic pseudosyncope than chance predicts (Blad et al., 2015) and the occurrence of psychogenic TLOC may depend on suggestion (Whitehead et al., 2017), which may be aided by a provocation procedure. The same protocol may be used for psychogenic TLOC as for reflex syncope, except for the endpoint.

- Ensure a supine rest phase before head-up tilt of at least 5 min.
- Head-up tilt angle between 60° and 80°, measured from the horizontal.
- Head-up tilt for up to 40 min (pharmacological provocation) or 45 min (no pharmacological provocation)
- Pharmacological provocation may be chosen and performed as for reflex syncope
- The test ends when the allotted time has passed or when psychogenic TLOC occurs. As blood pressure is not low and cerebral hypoperfusion is absent in psychogenic TLOC, tilting back may safely be postponed until all symptoms and signs are recognised.
- Complaint recognition: see above.

### 5.1.4. Indication POTS

Documentation of an exaggerated postural tachycardia using TTT or an active standing test may support a clinical diagnosis of POTS. The panel recommends the use of TTT in those with complaints of both orthostatic intolerance and TLOC as TTT has the advantage to provoke VVS as well. The diagnostic criteria of POTS require a head-up tilt period of 10 min. If other causes of orthostatic intolerance are considered (notably vasovagal presyncope or delayed OH) a longer head-up tilt is needed to establish a diagnosis. As the pathophysiological relation between orthostatic tachycardia and symptoms is as yet unclear, the panel stresses that orthostatic tachycardia without symptoms is a nonspecific

finding. Dehydration should be considered as an explanation. The test does not aim to provoke syncope, but if it occurs, it should be considered that the initial high heart rate upon standing may be a reflection of compensatory mechanisms and thus an expression of VVS. In such case it would be worthwhile to repeat TTT to see whether POTS is a consistent finding.

- Ensure a supine rest phase before head-up tilt of at least 5 min with reliable and constant BP and HR values as a baseline.
- Head-up tilt angle between 60° and 80°, measured from the horizontal.
- Head-up tilt for 10 min
- The test ends when the allotted time has passed, when 10 min have passed or when syncope inadvertently occurs, in which case the rules for tilting back of reflex syncope apply.
- Complaint recognition: see above.

## 5.2. TTT: possible extensions & variants

### 5.2.1. Active standing test with continuous blood pressure measurements

The active standing test can be used to provoke initial orthostatic hypotension (initial OH), classical OH (classic OH), POTS or to demonstrate delayed orthostatic BP recovery. It includes supine rest for at least 5 min followed by 3 min of standing. For a more detailed description of the active standing test, see the recent review by Finucane et al. (Finucane et al., 2019). A diagnosis of initial OH can be established if a transient BP drop (> 40 mmHg systolic blood pressure or >20 mmHg diastolic blood pressure) occurs within 15 s of standing (Freeman et al., 2011). Delayed orthostatic BP recovery is defined as the inability of systolic BP to recover to supine baseline values within 30 s of standing; the orthostatic BP should be  $\geq 20$  mmHg lower than the supine baseline values, but should not meet the criteria of classic or initial OH (van Wijnen et al., 2017). In view of the short duration of the blood pressure decrease in initial OH, the active standing test requires a beat-to-beat blood pressure measurement device. The active standing test can also be used to detect classic OH, as the criteria only require a measurement period of 3 min after standing up. In contrast to initial OH, classic OH or delayed OH can be detected with conventional intermittent BP measurements (Pavy-Le Traon et al., 2016).

### 5.2.2. Valsalva manoeuvre and deep breathing

Autonomic function tests may help to (1) identify autonomic failure as the underlying cause of classic OH (2) to characterize cardiovascular sympathetic and parasympathetic autonomic function and (3) quantify the severity of autonomic dysfunction (Brignole et al., 2018a; Hilz and Dutsch, 2006; Brignole et al., 2018b; Low, 2003). To ensure valid results it is extremely important that the tests are performed by trained personnel and under controlled circumstances, meaning a quiet and temperature controlled room; no meals for 3 h before the test, and no vasoactive substances (nicotine or caffeine-, theine-, or taurine-containing drinks) or medications on the day of examination. During the Valsalva manoeuvre, the patient is asked to conduct a maximally forced expiration for 15 s with an open glottis, i.e. with open nose and mouth, or into a closed loop system with a resistance of 40 mmHg. During the deep breathing test, the patient is asked to breathe deeply at 6 breaths per minute for 1 min.

### 5.2.3. Carotid sinus massage (CSM)

The diagnostic yield of the TTT can be enhanced by adding carotid sinus massage (CSM). CSM has been advocated in adults over 40 years of age with a history of unexplained reflex syncope or falls, and those in whom TLOC is triggered by head rotation or pressure on carotid sinus (Brignole et al., 2018a; Pasquier et al., 2017). The need for a definite diagnosis should always be weighed against the risk for complications (see Section 7).

#### 5.2.4. Venepuncture

Adding venepuncture provocation may increase diagnostic yield especially in those with a history of syncope during instrumentation (Humm and Z'Graggen, 2015). If intended for blood sampling during the test it is recommended to be performed at least 20 min before TTT, so as to avoid provocation.

#### 5.2.5. Meal provocation

A meal provocation can be used to diagnose postprandial hypotension (Pavelic et al., 2017). Postprandial hypotension can be detected with conventional intermittent BP measurements. The panel recognises that there is no consensus regarding the definition and the assessment of postprandial hypotension. Most authors define postprandial hypotension as a fall of  $\geq 20$  mmHg in systolic blood pressure or a SBP drop from  $>100$  mmHg to SBP  $<90$  mmHg within 2 h after completion of the meal (Pavelic et al., 2017). Various provocations have been used including orally administered glucose, a standard (normal) meal or a mixed liquid meals (Pavelic et al., 2017). Consensus is also lacking regarding the frequency of measurements and the body position during the test (supine, sitting) (Pavelic et al., 2017). The panel recognises the need for uniform definitions and protocols to foster our understanding of postprandial hypotension.

#### 5.2.6. Tailored provocations

A personally tailored provocation may be carried out in those with a specific and consistent trigger of TLOC. Examples include standing still following exercise (Thijs et al., 2003; Kim et al., 2017), coughing (Mereu et al., 2016), or bending forward (Taal et al., 2003). It should, however, be noted that many specific triggers suggest situational reflex syncope with high confidence and usually do not require diagnostic confirmation. Tailored provocation should therefore be reserved for those with unusual triggers or specific triggers but conflicting ictal signs or symptoms. Some provocation may need precautions to avoid injuries, such as floor mats.

#### 5.2.7. Counter pressure manoeuvres

The beat-to-beat blood pressure monitor can be used to teach patients suffering from reflex syncope or orthostatic hypotension to recognise hypotensive symptoms and to perform physical counter manoeuvres (van Dijk et al., 2006; Wieling et al., 2015b). Various manoeuvres including leg crossing, handgrip or arm tensing can be taught while patients see the blood pressure response on the monitor. Such training sessions allow tailoring of the manoeuvres. For example, leg crossing may not be feasible for those with movement disorders, but handgrip, arm tensing or tensing of the buttocks may help to improve orthostatic tolerance. Ideally, these manoeuvres are taught when patients experience mild symptoms of OH without the risk of immediate syncope or falls, e.g. immediately following TTT or active standing test.

### 6. Patient information

The panel recognises that there is ample literature to support specific patient recommendations to prepare for TTT. It could be considered to instruct patients to refrain from easily preventable vasoactive substances (e.g. caffeinated beverages, tobacco smoking) as this may limit the ability of TTT to provoke symptoms. The recommendation regarding medication fully depends upon the clinical context: if symptoms are likely provoked by medication the patient should continue the drug regimen. In other cases, it may be needed to test the patients off medication as certain drugs may confound the assessment of autonomic functions. In either case, reporting of the drug regimen is important to properly interpret TTT, as drug-induced orthostatic intolerance is a major cause of all three patterns of OH, particularly among elderly (Brignole et al., 2018a; Saedon et al., 2020). The panel recommends to advise patients to empty the bladder prior to test to avoid sympathetic stress (Fagius and Karhuvaara, 1989) and incontinence in case TTT provokes syncope.

Another consideration is the timing of the tests, as TTT more likely provokes syncope in the morning hours (Hu et al., 2011).

### 7. Safety

TTT is safe and major complications are very rare (Brignole et al., 2018a; Shen et al., 2017; Baron-Esquivias et al., 2002; Gieroba et al., 2004). Although we found no evidence that TTT is harmful, the low blood pressure during syncope might theoretically harm those with ischaemic disorders of the heart or brain, so risks and benefits should be weighed in such patients. There is an association between frequent syncope, defined as five or more syncopal spells during life, with more white matter lesions (Kruit et al., 2013). The panel felt that the advantages of a definite diagnosis, with possibly fewer future syncopal spells as a result, outweigh the theoretical risk of adding white matter lesions by one syncopal spell during TTT. TTT may provoke atrial fibrillation but this is usually self-limiting (Brignole et al., 2018a). Other side effects include headache or migraine following TNG challenge.

Complications following CSM are uncommon ( $<0.5\%$ ) and predominantly involve transient neurological symptoms while some cases with stroke have been reported (Bastulli and Orłowski, 1985; Munro et al., 1994; Richardson et al., 2000; Puggioni et al., 2002; Curro Dossi et al., 2010; Ungar et al., 2016). One death after CSM resulting from massive brain infarction has, however, been reported (van Munster et al., 2017). The risks and benefits of CSM therefore have to be weighed carefully. According to the ESC guidelines CSM should be undertaken with caution in patients with previous stroke, TIA or known carotid stenosis  $>70\%$  (Brignole et al., 2018a). The American Heart Association (AHA) and the American College of Cardiology (ACC) guidelines discourage the use of CSM in patients with a carotid bruit, TIA, stroke or myocardial infarction in the preceding months, except if carotid doppler excludes a significant stenosis (Shen et al., 2017). The expert group noticed that the more stringent criteria by the AHA/ACC more closely resemble the selection criteria employed in the studies assessing complication risks (bruits: 3 out of 4 studies; myocardial infarction: 2 out of 4 studies). The expert group feels that the need for a definite diagnosis of carotid sinus hypersensitivity should always be weighed against the risk for complications. Ultrasound investigation of the carotid arteries may be carried out to screen for atherosclerosis. There is, however, no evidence that this approach prevents complications.

### 8. Clinical environment

It is strongly recommended that TTT is an integral part of a clinical facility for the diagnosis and management of syncope and related symptoms, with dedicated staff and access to appropriate diagnostics and therapies. A consensus statement of the European Heart Rhythm Association (EHRA) offers guidance how to set up such facility (Kenny et al., 2015). The panel endorses the ESC recommendations regarding TTT performance: the test should be conducted by a physician, technician or a nurse trained in syncope and resuscitation (Brignole et al., 2018a). Clinical TTT should typically be conducted within a clinical environment with established procedures for life-threatening events (Kenny et al., 2015).

#### Declaration of competing interest

Dr. Thijs reports personal fees from Medtronic, Union Chimique Belge (UCB), Glaxo Smith Kline (GSK), Theravance, Novartis and grants from Dutch National Epilepsy Fund, Christelijke Vereniging voor de Verpleging van Lijders aan Epilepsie, AC Thomson Foundation, Medtronic, The Netherlands Organisation for Health Research and Development (843002707), outside the submitted work. Dr. Fanciulli reports royalties from Springer Nature Publishing Group and Thieme Verlag, speaker fees from the Austrian Autonomic Society, Austrian Neurology Society, Austrian Parkinson Society, Ordensklinikum Linz,

International Parkinson's disease and Movement Disorders Society and Theravance Biopharma, research grants from the Stichting Parkinson-Fond, MSA Coalition and the Österreichischer Austausch Dienst, outside of the submitted work. Prof Sutton serves as consultant to Medtronic Inc., is a member of Speakers Bureau SJM Abbott Laboratories, and shareholder in Edwards Lifesciences and Boston Scientific Corp, outside of the submitted work. Prof Pavy-Le Traon reports research grant from the French Ministry of Health, outside of submitted work, outside of the submitted work. Prof Jordan served as advisor for Novo-Nordisk, Bayer, received research support from Boston-Scientific and Boehringer Ingelheim and is cofounder of Eternygen GmbH, outside the submitted work. Prof Habek participated as clinical investigator, speaker and/or advisor for Biogen, Sanofi Genzyme, Merck, Bayer, Novartis, Pliva/Teva, Roche, Alvogen, Actelion, Alexion Pharmaceuticals, outside the submitted work. Prof Struhal reports consultancy for Boehringer, Eli Lilly, royalties from Manz-Rechtsverlag, Springer, Oxford University Press, speaker fees from Boehringer, Merz, Medconvent, Donau-Universität Krems, Austrian Society of Neurology, Manz-Rechtsverlag. Prof. Hilz received consulting honoraria from Alnylam and Sanofi-Genzyme, lecturing honoraria and travel support from Bayer Health Care, Amicus, Sanofi-Genzyme, and research funding from Novartis Pharma and Bayer Health Care.

## References

- Bachus, E., Holm, H., Hamrefors, V., Melander, O., Sutton, R., Magnusson, M., Fedorowski, A., 2018. Monitoring of cerebral oximetry during head-up tilt test in adults with history of syncope and orthostatic intolerance. *Europace: European pacing, arrhythmias, and cardiac electrophysiology: journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology* 20, 1535–1542.
- Baron-Eskivias, G., Pedrote, A., Cayuela, A., Valle, J.I., Fernandez, J.M., Arana, E., Fernandez, M., Morales, F., Burgos, J., Martinez-Rubio, A., 2002. Long-term outcome of patients with asystole induced by head-up tilt test. *Eur. Heart J.* 23, 483–489.
- Bartoletti A, Gaggioli G, Menozzi C, Bottoni N, Del Rosso A, Mureddu R, Musso G, Foglia-Manzillo G, Bonfigli B, Brignole M (1999) Head-up tilt testing potentiated with oral nitroglycerin: a randomized trial of the contribution of a drug-free phase and a nitroglycerin phase in the diagnosis of neurally mediated syncope. *Europace: European pacing, arrhythmias, and cardiac electrophysiology: journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology* 1:183–186.
- Bastulli, J.A., Orlowski, J.P., 1985. Stroke as a complication of carotid sinus massage. *Crit. Care Med.* 13, 869.
- Benditt, D.G., Detloff, B.L., Adkisson, W.O., Lu, F., Sakaguchi, S., Schussler, S., Austin, E., Chen, L.Y., 2012. Age-dependence of relative change in circulating epinephrine and norepinephrine concentrations during tilt-induced vasovagal syncope. *Heart Rhythm.* 9, 1847–1852.
- Blad, H., Lamberts, R.J., van Dijk, G.J., Thijs, R.D., 2015. Tilt-induced vasovagal syncope and psychogenic pseudosyncope: overlapping clinical entities. *Neurology* 85, 2006–2010.
- Brignole, M., Deharo, J.C., De Roy, L., Menozzi, C., Blommaert, D., Dabiri, L., Ruf, J., Guieu, R., 2011. Syncope due to idiopathic paroxysmal atrioventricular block: long-term follow-up of a distinct form of atrioventricular block. *J. Am. Coll. Cardiol.* 58, 167–173.
- Brignole, M., Moya, A., de Lange, F.J., Deharo, J.C., Elliott, P.M., Fanciulli, A., Fedorowski, A., Furlan, R., Kenny, R.A., Martin, A., Probst, V., Reed, M.J., Rice, C.P., Sutton, R., Ungar, A., van Dijk, J.G., 2018a. 2018 ESC Guidelines for the diagnosis and management of syncope. *Eur. Heart J.* 39, 1883–1948.
- Brignole, M., Moya, A., de Lange, F.J., Deharo, J.C., Elliott, P.M., Fanciulli, A., Fedorowski, A., Furlan, R., Kenny, R.A., Martin, A., Probst, V., Reed, M.J., Rice, C.P., Sutton, R., Ungar, A., van Dijk, J.G., 2018b. Practical instructions for the 2018 ESC Guidelines for the diagnosis and management of syncope. *Eur. Heart J.* 39, e43–e80.
- Cheshire Jr., W.P., Goldstein, D.S., 2019. Autonomic uprising: the tilt table test in autonomic medicine. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 29, 215–230.
- Curro Dossi, R., Roscia, G., Turri, E., Dall'ora, E., Sansone, S., Stockner, I., Wiedermann, C.J., 2010. Acute ischemic stroke complicating carotid sinus massage in the absence of carotid artery disease and failure of thrombolytic therapy. *Minerva Med.* 101, 193.
- El-Bedawi, K.M., Hainsworth, R., 1994. Combined head-up tilt and lower body suction: a test of orthostatic tolerance. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 4, 41–47.
- Fagius J, Karhuvaara S (1989) Sympathetic activity and blood pressure increases with bladder distension in humans. *Hypertension (Dallas, Tex. : 1979)* 14:511-517.
- Fanciulli A, Jordan J, Biaggioni I, Calandra-Buonaura G, Cheshire WP, Cortelli P, Eschlboeck S, Grassi G, Hilz MJ, Kaufmann H, Lahrmann H, Mancina G, Mayer G, Norcliffe-Kaufmann L, Pavy-Le Traon A, Raj SR, Robertson D, Rocha I, Struhal W, Thijs R, Tsioufis KP, van Dijk JG, Wenning GK (2018) Consensus statement on the definition of neurogenic supine hypertension in cardiovascular autonomic failure by the American Autonomic Society (AAS) and the European Federation of Autonomic Societies (EFAS): endorsed by the European Academy of Neurology (EAN) and the European Society of Hypertension (ESH). *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 28:355–362.
- Fedorowski, A., Hamrefors, V., Sutton, R., van Dijk, J.G., Freeman, R., Lenders, J.W., Wieling, W., 2017. Do we need to evaluate diastolic blood pressure in patients with suspected orthostatic hypotension? *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 27, 167–173.
- Finucane C, van Wijnen VK, Fan CW, Soraghan C, Byrne L, Westerhof BE, Freeman R, Fedorowski A, Harms MPM, Wieling W, Kenny R (2019) A practical guide to active stand testing and analysis using continuous beat-to-beat non-invasive blood pressure monitoring. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society.*
- Forleo, C., Guida, P., Iacoviello, M., Resta, M., Monitillo, F., Sorrentino, S., Favale, S., 2013. Head-up tilt testing for diagnosing vasovagal syncope: a meta-analysis. *Int. J. Cardiol.* 168, 27–35.
- Freeman, R., Wieling, W., Axelrod, F.B., Benditt, D.G., Benarroch, E., Biaggioni, I., Cheshire, W.P., Chelmsky, T., Cortelli, P., Gibbons, C.H., Goldstein, D.S., Hainsworth, R., Hilz, M.J., Jacob, G., Kaufmann, H., Jordan, J., Lipsitz, L.A., Levine, B.D., Low, P.A., Mathias, C., Raj, S.R., Robertson, D., Sandroni, P., Schatz, I., Schondorff, R., Stewart, J.M., van Dijk, J.G., 2011. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 21, 69–72.
- Freeman R IB, Lapusca R, Campagnolo M, Abuzinadah AR, Bonyhay I, Sinn DI, Miglis M, White J, Gibbons CH. (2020) Symptom recognition is impaired in patients with orthostatic hypotension. *Hypertension (Dallas, Tex. : 1979)* in press.
- Gibbons, C.H., Freeman, R., 2006. Delayed orthostatic hypotension: a frequent cause of orthostatic intolerance. *Neurology* 67, 28–32.
- Gibbons, C.H., Freeman, R., 2015. Clinical implications of delayed orthostatic hypotension: a 10-year follow-up study. *Neurology* 85, 1362–1367.
- Gibbons, C.H., Schmidt, P., Biaggioni, I., Frazier-Mills, C., Freeman, R., Isaacson, S., Karabin, B., Kuritzky, L., Lew, M., Low, P., Mehdird, A., Raj, S.R., Vernino, S., Kaufmann, H., 2017. The recommendations of a consensus panel for the screening, diagnosis, and treatment of neurogenic orthostatic hypotension and associated supine hypertension. *J. Neurol.* 264, 1567–1582.
- Gieroba, Z.J., Newton, J.L., Parry, S.W., Norton, M., Lawson, J., Kenny, R.A., 2004. Unprovoked and glyceryl trinitrate-provoked head-up tilt table test is safe in older people: a review of 10 years' experience. *J. Am. Geriatr. Soc.* 52, 1913–1915.
- Goldstein, D.S., Cheshire Jr., W.P., 2017. Beat-to-beat blood pressure and heart rate responses to the Valsalva maneuver. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 27, 361–367.
- Goldstein, D.S., Cheshire, W.P., 2018. Roles of catechol neurochemistry in autonomic function testing. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 28, 273–288.
- Goldstein, D.S., Holmes, C., Sharabi, Y., Brentzel, S., Eisenhofer, G., 2003. Plasma levels of catechols and metanephrines in neurogenic orthostatic hypotension. *Neurology* 60, 1327–1332.
- Goswami, N., Blaber, A.P., Hinghofer-Szalkay, H., Convertino, V.A., 2019. Lower body negative pressure: physiological effects, applications, and implementation. *Physiol. Rev.* 99, 807–851.
- Heitlerachi, E., Lord, S.R., Meyerkort, P., McCloskey, I., Fitzpatrick, R., 2002. Blood pressure changes on upright tilting predict falls in older people. *Age Ageing* 31, 181–186.
- Hilz, M.J., Dutsch, M., 2006. Quantitative studies of autonomic function. *Muscle Nerve* 33, 6–20.
- Hu, K., Scheer, F.A., Laker, M., Smales, C., Shea, S.A., 2011. Endogenous circadian rhythm in vasovagal response to head-up tilt. *Circulation* 123, 961–970.
- Humm, A.M., Z'Graggen, W.J., 2015. Venepuncture during head-up tilt testing in patients with suspected vasovagal syncope - implications for the test protocol. *Eur. J. Neurol.* 22, 389–394.
- Imholz, B.P., Wieling, W., van Montfrans, G.A., Wesseling, K.H., 1998. Fifteen years experience with finger arterial pressure monitoring: assessment of the technology. *Cardiovasc. Res.* 38, 605–616.
- Kenny RA, Ingram A, Bayliss J, Sutton R (1986) Head-up tilt: a useful test for investigating unexplained syncope. *Lancet (London, England)* 1:1352-1355.
- Kenny, R.A., Brignole, M., Dan, G.A., Deharo, J.C., van Dijk, J.G., Doherty, C., Hamdan, M., Moya, A., Parry, S.W., Sutton, R., Ungar, A., Wieling, W., 2015. Syncope unit: rationale and requirement—the European Heart Rhythm Association position statement endorsed by the Heart Rhythm Society. *Europace: European pacing, arrhythmias, and cardiac electrophysiology: journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology* 17, 1325–1340.
- Kharrazzaha, I., Holm, H., Bachus, E., Ricci, F., Sutton, R., Fedorowski, A., Hamrefors, V., 2019. Cerebral oximetry in syncope and syndromes of orthostatic intolerance. *Frontiers in Cardiovascular Medicine* 6, 171.
- Kim, T.H., Jang, H.J., Kim, S., Cho, S.Y., Song, K.S., Pickett, C., Schmitt, H.J., Lee, J., 2017. A new test for diagnosing vasovagal syncope: standing after treadmill test with sublingual nitrate administration. *PLoS One* 12, e0179631.
- Kruit, M.C., Thijs, R.D., Ferrari, M.D., Launer, L.J., van Buchem, M.A., van Dijk, J.G., 2013. Syncope and orthostatic intolerance increase risk of brain lesions in migraineurs and controls. *Neurology* 80, 1958–1965.
- Kulkarni, N., Mody, P., Levine, B.D., 2020. Abolish the tilt table test for the workup of syncope! *Circulation* 141, 335–337.

- LaFrance Jr., W.C., Baker, G.A., Duncan, R., Goldstein, L.H., Reuber, M., 2013. Minimum requirements for the diagnosis of psychogenic nonepileptic seizures: a staged approach: a report from the International League Against Epilepsy Nonepileptic Seizures Task Force. *Epilepsia* 54, 2005–2018.
- Low, P.A., 2003. Testing the autonomic nervous system. *Semin. Neurol.* 23, 407–422.
- Menant, J.C., Wong, A.K., Trollor, J.N., Close, J.C., Lord, S.R., 2016. Depressive symptoms and orthostatic hypotension are risk factors for unexplained falls in community-living older people. *J. Am. Geriatr. Soc.* 64, 1073–1078.
- Mereu, R., Taraborrelli, P., Sau, A., Di Toro, A., Halim, S., Hayat, S., Bernardi, L., Francis, D.P., Sutton, R., Lim, P.B., 2016. Diagnostic role of head-up tilt test in patients with cough syncope. *Europace: European pacing, arrhythmias, and cardiac electrophysiology: journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology* 18, 1273–1279.
- Munro, N.C., McIntosh, S., Lawson, J., Morley, C.A., Sutton, R., Kenny, R.A., 1994. Incidence of complications after carotid sinus massage in older patients with syncope. *J. Am. Geriatr. Soc.* 42, 1248–1251.
- Naschitz, J.E., Rosner, I., Rozenbaum, M., Gaitini, L., Bistrizki, I., Zuckerman, E., Sabo, E., Yeshurun, D., 2000. The capnography head-up tilt test for evaluation of chronic fatigue syndrome. *Semin. Arthritis Rheum.* 30, 79–86.
- Norcliffe-Kaufmann, L., Kaufmann, H., Palma, J.A., Shibao, C.A., Biaggioni, I., Peltier, A. C., Singer, W., Low, P.A., Goldstein, D.S., Gibbons, C.H., Freeman, R., Robertson, D., 2018a. Orthostatic heart rate changes in patients with autonomic failure caused by neurodegenerative synucleinopathies. *Ann. Neurol.* 83, 522–531.
- Norcliffe-Kaufmann, L., Galindo-Mendez, B., Garcia-Guariz, A.L., Villarreal-Vitorica, E., Novak, V., 2018b. Transcranial Doppler in autonomic testing: standards and clinical applications. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 28, 187–202.
- Norcliffe-Kaufmann, L.J., Kaufmann, H., Hainsworth, R., 2008. Enhanced vascular responses to hypocapnia in neurally mediated syncope. *Ann. Neurol.* 63, 288–294.
- Palma, J.A., Gomez-Esteban, J.C., Norcliffe-Kaufmann, L., Martinez, J., Tijero, B., Berganzo, K., Kaufmann, H., 2015. Orthostatic hypotension in Parkinson disease: how much you fall or how low you go? *Movement disorders: official journal of the Movement Disorder Society* 30, 639–645.
- Pasquier, M., Clair, M., Pruvot, E., Hugli, O., Carron, P.N., 2017. Carotid sinus massage. *N. Engl. J. Med.* 377, e21.
- Pavelic, A., Krbot Skoric, M., Cmosija, L., Habek, M., 2017. Postprandial hypotension in neurological disorders: systematic review and meta-analysis. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 27, 263–271.
- Pavy-Le Traon, A., Piedvache, A., Perez-Lloret, S., Calandra-Buonaura, G., Cochen-De Cock, V., Colosimo, C., Cortelli, P., Debs, R., Duerr, S., Fanciulli, A., Foubert-Samier, A., Gerdelat, A., Gurevich, T., Krismer, F., Poewe, W., Tison, F., Tranchant, C., Wenning, G., Rascol, O., Meissner, W.G., 2016. New insights into orthostatic hypotension in multiple system atrophy: a European multicentre cohort study. *J. Neurol. Neurosurg. Psychiatry* 87, 554–561.
- Puggioni, E., Guiducci, V., Brignole, M., Menozzi, C., Oddone, D., Donato, P., Croci, F., Solano, A., Lollì, G., Tomasi, C., Bottoni, N., 2002. Results and complications of the carotid sinus massage performed according to the “method of symptoms”. *Am. J. Cardiol.* 89, 599–601.
- Raj, S.R., 2013. Postural tachycardia syndrome (POTS). *Circulation* 127, 2336–2342.
- Raj, V., Rowe, A.A., Fleisch, S.B., Paranjape, S.Y., Arain, A.M., Nicolson, S.E., 2014. Psychogenic pseudosyncope: diagnosis and management. *Autonomic neuroscience: basic & clinical* 184, 66–72.
- Richardson, D.A., Bexton, R., Shaw, F.E., Steen, N., Bond, J., Kenny, R.A., 2000. Complications of carotid sinus massage—a prospective series of older patients. *Age Ageing* 29, 413–417.
- Saal, D.P., Thijs, R.D., van Dijk, J.G., 2016. Tilt table testing in neurology and clinical neurophysiology. *Clinical neurophysiology: official journal of the International Federation of Clinical Neurophysiology* 127, 1022–1030.
- Saal, D.P., Thijs, R.D., van Zwet, E.W., Bootsma, M., Brignole, M., Benditt, D.G., van Dijk, J.G., 2017. Temporal relationship of asystole to onset of transient loss of consciousness in tilt-induced reflex syncope. *JACC. Clinical electrophysiology* 3, 1592–1598.
- Saedon, N.I., Tan, M.P., Frith, J., 2020. The prevalence of orthostatic hypotension: a systematic review and meta-analysis. *J. Gerontol. A Biol. Sci. Med. Sci.* 75, 117–122.
- Sheldon, R., 2005. Tilt testing for syncope: a reappraisal. *Curr. Opin. Cardiol.* 20, 38–41.
- Sheldon, R.S., Sheldon, A.G., Serletis, A., Connolly, S.J., Morillo, C.A., Klinghenben, T., Krahn, A.D., Koshman, M.L., Ritchie, D., 2007. Worsening of symptoms before presentation with vasovagal syncope. *J. Cardiovasc. Electrophysiol.* 18, 954–959.
- Sheldon, R.S., Grubb 2nd, B.P., Olshansky, B., Shen, W.K., Calkins, H., Brignole, M., Raj, S.R., Krahn, A.D., Morillo, C.A., Stewart, J.M., Sutton, R., Sandroni, P., Friday, K.J., Hachul, D.T., Cohen, M.I., Lau, D.H., Mayuga, K.A., Moak, J.P., Sandhu, R.K., Kanjwal, K., 2015. 2015 heart rhythm society expert consensus statement on the diagnosis and treatment of postural tachycardia syndrome, inappropriate sinus tachycardia, and vasovagal syncope. *Heart Rhythm.* 12, e41–e63.
- Shen, W.K., Sheldon, R.S., Benditt, D.G., Cohen, M.I., Forman, D.E., Goldberger, Z.D., Grubb, B.P., Hamdan, M.H., Krahn, A.D., Link, M.S., Olshansky, B., Raj, S.R., Sandhu, R.K., Sorajja, D., Sun, B.C., Yancy, C.W., 2017. 2017 ACC/AHA/HRS guideline for the evaluation and management of patients with syncope: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Heart Rhythm.* 14, e155–e217.
- Shmueli, S., Bauer, P.R., van Zwet, E.W., van Dijk, J.G., Thijs, R.D., 2018. Differentiating motor phenomena in tilt-induced syncope and convulsive seizures. *Neurology* 90, e1339–e1346.
- Singer, W., Berini, S.E., Sandroni, P., Fealey, R.D., Coon, E.A., Suarez, M.D., Benarroch, E.E., Low, P.A., 2017. Pure autonomic failure: predictors of conversion to clinical CNS involvement. *Neurology* 88, 1129–1136.
- Stewart, J.M., Pianosi, P., Shaban, M.A., Terilli, C., Svistunova, M., Visintainer, P., Medow, M.S., 2018. Postural hyperventilation as a cause of postural tachycardia syndrome: increased systemic vascular resistance and decreased cardiac output when upright in all postural tachycardia syndrome variants. *Journal of the American Heart Association* 7.
- Sutton, R., 2015. The value of tilt testing and autonomic nervous system assessment. *Cardiol. Clin.* 33, 357–360.
- Sutton, R., Brignole, M., 2014. Twenty-eight years of research permit reinterpretation of tilt-testing: hypotensive susceptibility rather than diagnosis. *Eur. Heart J.* 35, 2211–2212.
- Sutton, R., van Dijk, N., Wieling, W., 2014. Clinical history in management of suspected syncope: a powerful diagnostic tool. *Cardiol. J.* 21, 651–657.
- Sutton, R., Fedorowski, A., Benditt, D.G., 2020. Letter by Sutton et al regarding article, “abolish the tilt table test for the workup of syncope!”. *Circulation* 141, e944–e945.
- Taal, W., van der Dussen, D.H., van Erven, L., van Dijk, J.G., 2003. Neurally-mediated complete heart block. *The Lancet. Neurology* 2, 255–256.
- Tannemaat, M.R., van Niekerk, J., Reijntjes, R.H., Thijs, R.D., Sutton, R., van Dijk, J.G., 2013. The semiology of tilt-induced psychogenic pseudosyncope. *Neurology* 81, 752–758.
- Thijs, R.D., Reijntjes, R.H., van Dijk, J.G., 2003. Water drinking as a potential treatment for idiopathic exercise-related syncope: a case report. *Clinical autonomic research: official journal of the Clinical Autonomic Research Society* 13, 103–105.
- Ungar, A., Rivasi, G., Rafanelli, M., Toffanello, G., Mussi, C., Ceccofiglio, A., McDonagh, R., Drumm, B., Marchionni, N., Alboni, P., Kenny, R.A., 2016. Safety and tolerability of tilt testing and carotid sinus massage in the octogenarians. *Age Ageing* 45, 242–248.
- van Dijk, J.G., Thijs, R.D., Benditt, D.G., Wieling, W., 2009. A guide to disorders causing transient loss of consciousness: focus on syncope. *Nat. Rev. Neurol.* 5, 438–448.
- van Dijk, J.G., Thijs, R.D., van Zwet, E., Tannemaat, M.R., van Niekerk, J., Benditt, D.G., Wieling, W., 2014. The semiology of tilt-induced reflex syncope in relation to electroencephalographic changes. *Brain* 137, 576–585.
- van Dijk, J.G., Ghariq, M., Kerkhof, F.I., Reijntjes, R., van Houwelingen, M.J., van Rossum, I.A., Saal, D.P., van Zwet, E.W., van Lieshout, J.J., Thijs, R.D., Benditt, D.G., 2020. Novel methods for quantification of vasodepression and cardioinhibition during tilt-induced vasovagal syncope. *Circ. Res.* 127, e126–e138.
- van Dijk, N., Quartieri, F., Blanc, J.J., Garcia-Civera, R., Brignole, M., Moya, A., Wieling, W., 2006. Effectiveness of physical counterpressure maneuvers in preventing vasovagal syncope: the Physical Counterpressure Manoeuvres Trial (PC-Trial). *J. Am. Coll. Cardiol.* 48, 1652–1657.
- van Dijk, N., Boer, K.R., Colman, N., Bakker, A., Stam, J., van Grieken, J.J., Wilde, A.A., Linzer, M., Reitsma, J.B., Wieling, W., 2008. High diagnostic yield and accuracy of history, physical examination, and ECG in patients with transient loss of consciousness in FAST: the Fainting Assessment study. *J. Cardiovasc. Electrophysiol.* 19, 48–55.
- van Munster CE, van Ballegoij WJ, Schroeder-Tanka JM, van den Berg-Vos RM (2017) [A Severe Stroke Following Carotid Sinus Massage]. *Nederlands tijdschrift voor geneeskunde* 161:D826.
- van Wijnen, V.K., Finucane, C., Harms, M.P.M., Nolan, H., Freeman, R.L., Westerhof, B. E., Kenny, R.A., Ter Maaten, J.C., Wieling, W., 2017. Noninvasive beat-to-beat finger arterial pressure monitoring during orthostasis: a comprehensive review of normal and abnormal responses at different ages. *J. Intern. Med.* 282, 468–483.
- Whitehead, K., Kane, N., Wardrope, A., Kandler, R., Reuber, M., 2017. Proposal for best practice in the use of video-EEG when psychogenic non-epileptic seizures are a possible diagnosis. *Clinical neurophysiology practice* 2, 130–139.
- Wieling W, Krediet CT, van Dijk N, Linzer M, Tschakovsky ME (2007) Initial orthostatic hypotension: review of a forgotten condition. *Clinical science (London, England : 1979)* 112:157-165.
- Wieling, W., van Dijk, N., de Lange, F.J., Olde Nordkamp, L.R., Thijs, R.D., van Dijk, J.G., Linzer, M., Sutton, R., 2015a. History taking as a diagnostic test in patients with syncope: developing expertise in syncope. *Eur. Heart J.* 36, 277–280.
- Wieling, W., van Dijk, N., Thijs, R.D., de Lange, F.J., Krediet, C.T., Halliwill, J.R., 2015b. Physical countermeasures to increase orthostatic tolerance. *J. Intern. Med.* 277, 69–82.
- Zysko, D., Fedorowski, A., Nilsson, D., Rudnicki, J., Gajek, J., Melander, O., Sutton, R., 2016. Tilt testing results are influenced by tilt protocol. *Europace: European pacing, arrhythmias, and cardiac electrophysiology: journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology* 18, 1108–1112.