

Potential for Phototrophy in Venus' Clouds

Rakesh Mogul,¹ Sanjay S. Limaye,² Yeon Joo Lee,³ and Michael Pasillas¹

Abstract

We show that solar irradiances calculated across Venus' clouds support the potential for Earth-like phototrophy and that treatment of Venus' aerosols containing neutralized sulfuric acid favor a habitable zone. The phototrophic potential of Venus' atmosphere was assessed by calculating irradiances (200–2000 nm, 15° solar zenith angle, local noon) using a radiative transfer model that accounted for absorption and scattering by the major and minor atmospheric constituents. Comparisons to Earth's surface (46 W m⁻², 280–400 nm) suggest that Venus' middle and lower clouds receive ~87% less normalized UV flux (6–7 W m⁻²) across 200–400 nm, yet similar normalized photon flux densities (~4400–6200 μmol m⁻² s⁻¹) across 350–1200 nm. Further, Venus' signature phototrophic windows and subwindows overlap with the absorption profiles of several photosynthetic pigments, especially bacteriochlorophyll *b* from intact cells and phycocyanin. Therefore, Venus' light, with limited UV flux in the middle and lower clouds, is likely quite favorable for phototrophy. We additionally present interpretations to refractive index and radio occultation measures for Venus' aerosols that suggest the presence of lower sulfuric abundances and/or neutralized forms of sulfuric acid, such as ammonium bisulfate. Under these considerations, the aerosols in Venus' middle clouds could harbor water activities (≥0.6) and buffered acidities (Hammett acidity factor, H_0 -0.1 to -1.5) that lie within the limits of acidic cultivation (≥ H_0 -0.4) and are tantalizingly close to the limits of oxygenic photosynthesis (≥ H_0 0.1). Together, these photophysical and chemical considerations support a potential for phototrophy in Venus' clouds. Key Words: Venus—Photosynthesis—Phototrophs—Aerosols—Irradiance—pH. *Astrobiology* 21, 1237–1249.

1. Introduction

WE PRESENT PHOTOPHYSICAL and chemical arguments for the potential of phototrophy in Venus' clouds or the harnessing of light for metabolic purposes. We show that the wavelength-dependent photon fluxes calculated across Venus' cloud layers are sufficient for terrestrial-like phototrophy. With regard to cloud layer habitability, we also present novel interpretations of the refractive index values obtained *in situ* and sulfuric acid vapor abundance profiles obtained through radio occultation that suggest, due to the presence of partly neutralized sulfuric acid, that acidity and water activity values in Venus' aerosols may potentially be suitable for microbial growth.

2. Photophysical Considerations

Solar radiances in the venusian atmosphere were measured by the Venera atmospheric entry probes (Economov

et al., 1984) and Pioneer Venus Large Probe (Tomasko *et al.*, 1980) during descents to the surface. Across the altitudes of 60–48 km (400–1000 nm; reported solar zenith angle 45°), normalized irradiances ranged from 840–1390 W m⁻² or 1080 ± 250 W m⁻² (Moroz *et al.*, 1985). For this study, venusian fluxes were normalized from the reported solar zenith angle (SZA) to 0° and adjusted for anisotropic scattering by the clouds using Equation 5a from the work of Moroz *et al.* (1985). In contrast, irradiances of ~930 W m⁻² (400–1000 nm, reported SZA 48.19°) are found on Earth's surface (ASTM G-173-03, standard reference) when considering cloudless conditions and normalization through division by the cosine of the reported SZA. These comparisons reveal that irradiances in Venus' middle and lower clouds are holistically similar to those at Earth's surface, where photosynthesis is abundant, thus providing tangible support for the potential for phototrophy at Venus.

¹Chemistry & Biochemistry Department, California State Polytechnic University, Pomona, California, USA.

²Space Science and Engineering Center, University of Wisconsin, Madison, Wisconsin, USA.

³Zentrum für Astronomie und Astrophysik, Technische Universität Berlin, Berlin, Germany.

When considering phototrophy, wavelength-dependent photon fluxes are critical due to a dependence on photon stoichiometry. Photon fluxes, however, could not be accurately estimated from Venera 11 and 14 spectra (Moroz *et al.*, 1982; Ignatiev *et al.*, 1997), which were low resolution and available only in graphical format, or extracted from the integrated irradiances from Pioneer Venus (Moroz *et al.*, 1985). Additionally, transmittance windows within the clouds could not be accurately established from Venera 11 and 14 spectra due to log-scale representations of the published spectra and reduced relative peak heights (Moroz *et al.*, 1982; Ignatiev *et al.*, 1997). Accordingly, we obtained radiances with high spectral and vertical resolution using radiative transfer calculations (Lee *et al.*, 2016, 2019) based on a commonly adopted Venus cloud model (Ragent *et al.*, 1985).

3. Calculated Irradiances

The radiative transfer calculations (local noon, SZA 15°) were described in the works of Lee *et al.* (2016) and Lee *et al.* (2019). For this study, we calculated solar irradiances and photon fluxes from 0.2–5 μm at 0.0001 nm resolution at altitudes ranging from the atmospheric tops (99.5 km) to the surface. Altitude profiles of the calculated spectra enabled delineation of the transmittance windows for solar irradiance within Venus' middle (56.5–50.5 km) and lower (50.5–47.5 km) clouds, along with identification of the wavelengths available for phototrophy.

Briefly, the calculations yielded spectra for downward direct and diffused irradiances that accounted for absorption and scattering by major and minor atmospheric constituents (CO_2 , H_2O , N_2 , SO_2 , COS , HCl , CO , HF , H_2S), along with scattering by aerosolized sulfuric acid, which has a minimal absorption at $\lambda < 3 \mu\text{m}$ (Hummel *et al.*, 1988). Optical depths ranged between 20 and 50 in the visible range, as observed *in situ* (Ragent *et al.*, 1985), and the vertical structure for the equatorial clouds (optical depth of 31 in the visible range) was from the work of Crisp (1986).

To account for the ultraviolet and blue radiation absorbers within Venus' clouds that, to date, remain unidentified (Pérez-Hoyos *et al.*, 2018; Titov *et al.*, 2018), the absorbers were treated as (abiotic) components of the aerosols, where the single scattering albedo of the aerosols was forced to < 1.0 between 310 and 780 nm across the altitudes of 57–71 km (Crisp, 1986). This treatment involved optimization to yield a scale factor of 1.18, which successfully fit the mean UV albedo at low latitudes (0–30°S) observed from 2006 through 2017 (Lee *et al.*, 2019). The resulting spectra are shown in Fig. 1A for the top of the atmosphere (TOA, 99.5 km); the upper (65.5 km), middle (53.5 km), and lower (49.5 km) clouds; and below the clouds (44.5 km).

Comparison with integrated irradiances measured *in situ* revealed similar values after correction for SZA and anisotropic scattering, as described. Calculated and normalized irradiances across 450–1200 nm of 998 W m^{-2} (53.5 km) and 741 W m^{-2} (49.5 km) were very close or within error to averaged irradiances from Venera 11, 13, and 14 (450–1200 nm) of $1085 \pm 86 \text{ W m}^{-2}$ (55 km) and $719 \pm 162 \text{ W m}^{-2}$ (50 km). However, normalized values from Pioneer Venus across 400–1000 nm of 1150 W m^{-2} (55 km) and 956 W m^{-2} (50 km) were ~ 20 – 30% higher than the calculated values (400–1000 nm) of 932 W m^{-2} (53.5 km) and 728 W m^{-2}

(49.5 km). These differences are suggestive of variations in cloud opacity across *in situ* measures and/or an indication of lower *in situ* abundances (along the incoming beam) of the unidentified blue-radiation absorbers in comparison to our model. Similarly, the wavelength of peak irradiance at 607.5 nm (53.5 km) from our calculations was $\sim 10\%$ higher than the value of 558 nm (52 km) recorded by Venera 14, which was suggestive of lower absorption coefficients and/or lower *in situ* abundances of the unidentified blue-radiation absorbers.

4. Comparisons of Venus and Earth Spectra

Expectedly, our calculations show that irradiances below Venus' cloud tops are substantially attenuated with decreasing altitude through absorption by SO_2 ($\leq 320 \text{ nm}$) and the unknown absorbers. Calculations conducted across the UV (200–400 nm) show that irradiances of 205 W m^{-2} at the TOA decrease by $\sim 97\%$ to ~ 6 – 7 W m^{-2} in the middle and lower clouds (Table 1). In comparison, Moroz (1983) reported a $\sim 90\%$ reduction of *in situ* UV flux at 370 nm in the middle clouds to yield 0.6–2.8 W m^{-2} (58–48 km). Our calculations show that (A) $>99.99\%$ of the total UV flux in the middle clouds arises in the UV-A region (320–400 nm), (B) $\sim 66\%$ of the total UV flux arises at wavelengths (370–400 nm) longer than those measured by the Venera probes, and (C) fluxes in the UV-B (280–320 nm) and UV-C (200–280 nm) decrease to negligible levels in the middle clouds from the respective fluxes of $\sim 20\%$ and $\sim 7\%$ in the TOA.

On Earth's surface (46 W m^{-2} , 280–400 nm), integrated irradiances across the UV-A and UV-B are ~ 7 -fold higher than those in Venus' middle clouds (6.7 W m^{-2} , 280–400 nm), after normalization as described. We note that measured UV irradiances on Earth range from ~ 12 – 29 W m^{-2} (normalized from SZA 30°) in the UV-A across North America (Grant and Slusser, 2005), where microbial activity is abundant, and can rise to $\sim 73 \text{ W m}^{-2}$ (normalized from SZA 16°) across the UV-A and UV-B (295–385 nm) in the Atacama Desert (Piacentini *et al.*, 2003), where microbial abundance and activity are limited but measurable (Warren-Rhodes *et al.*, 2006; Orellana *et al.*, 2020). Further, Lee *et al.* (2019) reported $\sim 30\%$ variations in global UV albedo of Venus on a decadal timescale from analysis of images from Venus Express (2006–2014) and Akatsuki (2016–present) orbiters. Therefore, given the observed variability and low flux, UV radiation is likely not a limiting constraint for habitability in Venus' middle and lower clouds.

We additionally compared photon fluxes ($\text{m}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$) at the wavelength of peak photon flux (λ_{peak}) across Venus' clouds and Earth's surface. As displayed in Fig. 1B and listed in Table 1, normalized photon fluxes in Venus' middle and lower clouds peak at 607.5 nm with values of 8.2×10^{18} and $7.0 \times 10^{18} \text{ m}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$, respectively. In contrast, normalized fluxes on Earth's surface peak at 669 nm with values of $6.6 \times 10^{18} \text{ m}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$, representing ~ 0.8 -fold of the flux in Venus' middle clouds, and a $\sim 60 \text{ nm}$ shift in the λ_{peak} toward the red. These differences are due to the modeled absorption by the unknown absorbers, and Rayleigh scattering in the atmosphere.

To account for cloud opacity variability, we estimated variances in the downward flux using *in situ* measures obtained by Venera 11, 13, and 14 (Table 6–7, Moroz *et al.*,




FIG. 1. (A) Irradiance spectra for Venus across the top of the atmosphere (99.5 km) and upper (66.5 km), middle (53.5 km), and lower (49.5 km) clouds, and below the clouds (44.5 km). (B) Photon flux spectra of Venus' middle and lower clouds and Earth's surface. (C) Photon flux spectra of Venus' middle and lower clouds with demarcations of bands associated with CO₂ and H₂O stretching.

TABLE 1. LIST OF SELECTED PHOTOPHYSICAL PARAMETERS¹ FOR VENUS (NORMALIZED TO SZA 0° FROM 15°C AND ADJUSTED FOR ANISOTROPIC SCATTERING) AT THE TOP OF THE ATMOSPHERE (TOA; 99.5 km) AND IN THE UPPER (65.5 km), MIDDLE (53.5 km), AND LOWER (49.5 km) CLOUDS, INCLUDING BELOW THE CLOUDS (BTC; 44.5 km), AS WELL AS EARTH'S SURFACE (NORMALIZED TO SZA 0° FROM 48.19°)

| Normalized flux and irradiances | λ_{peak} (nm) | Peak photon flux ^a ($\text{m}^{-2} \text{s}^{-1} \text{nm}^{-1}$) | Integrated UV photon flux ^b ($\text{m}^{-2} \text{s}^{-1}$) | Peak irradiance ^a ($\text{W m}^{-2} \mu\text{m}^{-1}$) | Integrated UV irradiance ^b (W m^{-2}) |
|---------------------------------|--|--|--|---|---|
| Venus' TOA (99.5 km) | 555.2 ^c 450.7 ^d | 1.07×10^{19} | 3.77×10^{20} | 4410 | 218 |
| Venus' upper clouds (65.5 km) | 607.2 | 1.17×10^{19} | 1.52×10^{20} | 3820 | 86.9 |
| Venus' middle clouds (53.5 km) | 607.5 | 8.18×10^{18} | 1.20×10^{19} | 2670 | 6.73 |
| Venus' lower clouds (48.5 km) | 607.5 | 7.01×10^{18} | 1.12×10^{19} | 2290 | 5.90 |
| Venus BTC (44.5 km) | 607.5 | 6.79×10^{18} | 1.08×10^{19} | 2220 | 5.69 |
| Earth's surface | 669.0 ^c 531.0 ^d | 6.58×10^{18} | 8.44×10^{19} | 2150 | 45.8 |

¹Selected parameters include λ_{peak} (for photon flux and irradiance), normalized peak photon flux ($\text{m}^{-2} \text{s}^{-1} \text{nm}^{-1}$), integrated and normalized UV photon flux between 200 and 400 nm ($\text{W m}^{-2} \mu\text{m}^{-1}$), normalized peak irradiance ($\text{m}^{-2} \text{s}^{-1}$), and integrated and normalized UV irradiance (W m^{-2}) between 200 and 400 nm.

^aNormalized peak photon flux or normalized peak irradiance at λ_{peak} .

^bIntegrated over 200–400 nm for Venus and 280–400 nm for Earth.

^c λ_{peak} for photon flux.

^d λ_{peak} for irradiance.

1985). Integrated downward fluxes (450–1200 nm) at the respective altitudes of 60, 55, 50, and 48 km were averaged across the Venera data and resulting standard deviations expressed as percentages to reveal variances ranging from 8–23% (or an average of $15 \pm 6\%$) across the middle and lower clouds. As inferred from these uncertainties (or variances in cloud opacity), fluxes over time at the respective λ_{peak} are likely comparable across Venus' middle and lower clouds and Earth's surface. This suggests that photon abundances and energies in Venus' middle and lower clouds are sufficient for Earth-like phototrophy and are potentially more favorable due to lower UV flux.

5. Earth Life in Venus' Light

On Earth, solar irradiance is regarded as a selective biological pressure for phototrophy, where the λ_{peak} (669 nm) correlates to the wavelength of maximum visible absorption by chlorophyll *a* (λ_{max} 669 nm), the dominant pigment used in photosynthesis (Nishio, 2000; Kiang *et al.*, 2007b). Similarly, solar irradiance on Venus could theoretically serve as a selective pressure and favor phototrophs that harbor pigments with maximal absorbances at the λ_{peak} of ~ 608 nm (or between 558 and 608 nm, when considering the range from Venera 14 and calculated spectra).

On Earth, the wavelengths available for phototrophy are also constrained by the absorption of atmospheric species such as water, O₂, O₃, and to a lesser degree CO₂. In turn, the light that is transmitted to Earth's surface overlaps with the absorption profiles of pigments used in photosynthesis (*e.g.*, chlorophylls, bacteriochlorophylls, and carotenoids) (Govindjee, 1974; Kiang *et al.*, 2007a, 2007b; Stomp *et al.*, 2007; Holtrop *et al.*, 2021). Analogously, the dominant absorption associated with CO₂ stretching in Venus' clouds (see Fig. 1C)—and to a lesser degree H₂O stretching—also defines regions of transmitted solar irradiance or potential windows for phototrophy.

Highlighted in Fig. 2A are the hypothetical phototrophic windows in Venus' clouds. As adapted from the work of Kiang *et al.* (2007a), the photosynthetic windows on Earth comprise three signature regions that relate to peak absorbances by the photosynthetic pigments. These regions comprise the λ_{peak} (W_P), as the principal window for energy input and selective biological pressure, and the longest (W_L) and shortest (W_S) wavelengths available for absorption by photosynthetic pigments. On Earth, the signature photosynthetic windows, as listed in Table 2, comprise (1) red light in the W_P (~ 590 – 685 nm), which overlaps with the absorption of chlorophylls *a* and *b*, (2) near-infrared wavelengths in the W_L (~ 950 – 1100 nm), which are absorbed by bacteriochlorophyll *b* (and related protein complexes), and (3) blue light in the W_S (~ 350 – 500 nm), which is absorbed by all pigments. Relatedly, absorption across W_P (red) and W_S (blue) by chlorophylls *a* and *b* results in the observed green reflectance of plants. Earth's spectra, of course, harbor phototrophic subwindows that overlap with pigments from niche environments (*e.g.*, chlorophyll *f*, bacteriochlorophyll *a*, and bacteriochlorophyll *b*) (Madigan, 2003; Stomp *et al.*, 2007).

For Venus—when using Earth pigments as a reference—the primary phototrophic windows, as listed in Table 2, are nominally ~ 590 – 656 nm (W_P), ~ 900 – 1200 nm (W_L), and ~ 400 – 500 nm (W_S), as constrained by the absorption of

CO₂ and H₂O in the atmosphere. Under such selective biological pressures, we speculate that dominant phototrophs within Venus' solar radiation might appear brown/maroon after absorption in the respective W_P (orange) and W_S (blue) regions. Demarcations for potential phototrophic subwindows in Venus' spectra are provided in Fig. 2A (asterisks). Potential low-energy niches include subwindows in the infrared of ~ 1210 – 1300 , ~ 1400 – 1425 , and ~ 1675 – 1875 nm (asterisks in Fig. 2A); however, plausible photosynthesis would require absorption of increased numbers of photons per fixed CO₂ (or conversion of CO₂ to (CH₂O)_{*n*})—or Venus equivalent) to attain energies equivalent to those from the W_P region (Wolstencroft and Raven, 2002; Kiang *et al.*, 2007a; Lehmer *et al.*, 2018).

Solar photon fluxes within Venus' middle clouds are compared in Fig. 2B to the absorption spectra of selected terrestrial anoxygenic phototrophs (intact cells) and photosynthetic pigments. The comparisons show that solar irradiances that transmit through W_P , W_L , and W_S directly overlap with the absorption bands of intact cells of purple nonsulfur and sulfur bacteria (such as *Blastochloris viridis* and *Ectothiorhodospira halochloris* BN 1850, respectively), which harbor macromolecular complexes that contain bacteriochlorophyll *b* and differing pigments (see Fig. 2B legend). Irradiances transmitting through W_P also overlap with the absorption of phycocyanin, a carotenoid from cyanobacteria, while irradiances that transmit through W_S overlap with the absorbances of all assessed biochemical/biological spectra (similar to Earth). In fact, as indicated in Fig. 2C, the absorption features from all listed terrestrial anoxygenic phototrophs (intact cells) overlap with Venus' phototrophic windows or subwindows.

Photosynthetic photon flux densities (PPFDs; $\mu\text{mol m}^{-2} \text{s}^{-1}$) were calculated across 400–700 nm (photosynthetically active radiation on Earth), 350–1200 nm (the wavelengths spanning W_S to W_L for Venus and Earth), and 700–1200 nm (near-infrared region). Normalized PPFD values (Table 2) in Venus' middle clouds were $2700 \mu\text{mol m}^{-2} \text{s}^{-1}$ (400–700 nm), $6230 \mu\text{mol m}^{-2} \text{s}^{-1}$ (350–1200 nm), and $3520 \mu\text{mol m}^{-2} \text{s}^{-1}$ (700–1200 nm); while those at Earth were $2610 \mu\text{mol m}^{-2} \text{s}^{-1}$ (400–700 nm), $6480 \mu\text{mol m}^{-2} \text{s}^{-1}$ (350–1200 nm), and $3780 \mu\text{mol m}^{-2} \text{s}^{-1}$ (700–1200 nm). Normalized values for Venus' lower clouds are provided in Table 2. Despite variances in the flux, Venus' cloud PPFD values are very similar to those at Earth's surface, while values in Venus' lower clouds are likely lower in the near infrared as suggested by decreases of $\sim 40\%$ and $\sim 45\%$ when compared to Venus' middle clouds and Earth's surface, respectively.

Intriguingly, appreciable fluxes of photons also arise from thermal emissions (~ 1000 – 1750 nm) from the hot surface and atmosphere below Venus' clouds (Crisp *et al.*, 1989; Meadows and Crisp, 1996; Mueller *et al.*, 2008; Longobardo *et al.*, 2012). As displayed in Fig. 2B (spectra #6, black dashed lines), Venus' thermal radiances, as measured on the nightside between 900 and 1320 nm (Meadows and Crisp, 1996), overlap with the absorption of intact cells of purple nonsulfur and sulfur bacteria that harbor bacteriochlorophyll *b*. Using spectra from the works of Mueller *et al.* (2008) and Meadows and Crisp (1996), nightside photon fluxes within W_L were calculated to be $8.3 \times 10^{16} \text{ m}^{-2} \text{ s}^{-1}$ (1000–1200 nm) and $6.7 \times 10^{16} \text{ m}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ (900–1200 nm), respectively; with an additional $1.3 \times 10^{16} \text{ m}^{-2} \text{ s}^{-1}$ and $1.1 \times 10^{16} \text{ m}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$




FIG. 2. (A) Potential phototropic windows of W_S , W_P , and W_L (green-tinted boxes) and subwindows (asterisks) in Venus' middle (red) and lower (green) clouds. (B) Comparison of the photon flux (left y-axis) and phototropic windows in Venus' middle clouds to the absorption profiles (right y-axis, relative absorbance) of (1) purified chlorophyll a; (2) purified chlorophyll b; (3) phycocyanin (lysozyme extract of *Alphanizomenon ovalisporum*); (4) *Halorhodospira halochloris* BN 1850 (intact cells; bacteriochlorophyll (Bchl) b + bacteriopheophytin b + methoxy-hydroxylycopene glucoside + dihydroxylycopene diglucoside + others); and (5) *Blastochloris viridis* (intact cells; Bchl b + dihydroneurosporene + dihydrolycopene); along with (6) the photon flux profile for Venus' nightside thermal emissions across 900–1320 nm from Meadows and Crisp (1996) (dashed black lines; right y-axis, relative photon flux). Block arrows represent the direction of photon flux into the cloud layer through the signature windows. (C) Comparison of the photon flux (left y-axis) in Venus' middle clouds (red) and on Earth's surface (blue) to the absorption profiles (right y-axis, relative absorbance) of (1a and 1b) chlorophylls a and b (purified); (2) *Pelodictyon phaeoclastratiforme* sp. nov. (intact cells; Bchl e and isorenieratenes); (3) *Chlorobium phaeobacteroides* MN1 (intact cells; Bchl e + Bchl a + isorenieratenes); (4) *Halorhodospira halophila* BN 9910 (intact cells; Bchl a); (5) *Roseospirillum parvum* (intact cells; Bchl a + spirilloxanthin + lycopenal); (6) *Halorhodospira halochloris* BN 1850 (intact cells; Bchl b + bacteriopheophytin b + methoxy-hydroxylycopene glucoside + dihydroxylycopene diglucoside + others); and (7) *Blastochloris viridis* (intact cells; Bchl b + dihydroneurosporene + dihydrolycopene) (Imhoff and Trüper, 1977; Thornber *et al.*, 1980; Overmann and Pfennig, 1989; Overmann *et al.*, 1992; Glaeser and Overmann, 1999; Takaichi *et al.*, 2001; Stomp *et al.*, 2007; Yacobi *et al.*, 2015; Imhoff, 2017).

TABLE 2. SIGNATURE PHOTOTROPHIC WINDOWS AND PHOTOSYNTHETIC PHOTON FLUX DENSITY (PPFD) FOR VENUS' MIDDLE (53.5 KM) AND LOWER (48.5 KM) CLOUDS AND EARTH'S SURFACE

| <i>Spectral windows</i> | W_P (nm) | W_L (nm) | W_S (nm) |
|--|-------------------|--------------------|--------------------|
| Venus' middle and lower clouds | ~590–656 | ~900–1200 | ~400–500 |
| Earth's surface | ~590–685 | ~950–1100 | ~350–500 |
| <i>PPFD</i> ($\mu\text{mol m}^{-2} \text{s}^{-1}$) | <i>400–700 nm</i> | <i>350–1200 nm</i> | <i>700–1200 nm</i> |
| Venus' middle clouds (53.5 km) | 2700 | 6230 | 3520 |
| Venus' lower clouds (49.5 km) | 2330 | 4430 | 2090 |
| Earth's surface | 2610 | 6480 | 3780 |

Phototrophic windows represent the regions of highest photon flux (W_P) and the shortest (W_S) and longest (W_L) suitable wavelengths of available solar radiation.

transmitting through the subwindow of ~1210–1300 nm, respectively. Consistent with our radiative transfer model (Figs. 1C and 2B), thermal emissions would be detectable by orbiter or Earth-based measures after transmission through W_L (for emissions at ~1030, 1100, and 1180 nm) and the subwindow of ~1210–1300 nm (for emissions at ~1280 and 1300 nm); additionally, the abrupt decrease in emission between ~1035 and 1050 nm, as observed in the spectra of Mueller *et al.* (2008) and Meadows and Crisp (1996), is due to attenuation by atmospheric CO₂ (or, absorption related to CO₂ stretching at 1037 and 1049 nm).

On Earth, anoxygenic low-light survival is associated with green sulfur bacteria in (1) the deep sea at hydrothermal vents in the East Pacific Rise (depth 2391 m), where photon fluxes of ~10¹⁵ m⁻² s⁻¹ sr⁻¹ (600–1000 nm) arising from geothermal emissions are reported to support phototrophy (Beatty *et al.*, 2005); (2) the Black Sea at depths of 120 m where the presence of phototrophs is associated with downwelling photon fluxes that range from 4.5 × 10¹⁴ to 1.3 × 10¹⁵ m⁻² s⁻¹, as adapted from PPFD values of 0.00075 to 0.0022 μmol m⁻² s⁻¹ (400–700 nm) reported by Manske *et al.* (2005); and (3) laboratory studies where fixation of CO₂ occurs at fluxes as low as 9.0 × 10¹⁵ m⁻² s⁻¹ (0.015 μmol m⁻² s⁻¹) (Manske *et al.*, 2005; Marschall *et al.*, 2010). In comparison, Venus' photon fluxes and PPFD values that arise from thermal radiances (0.14 μmol m⁻² s⁻¹, 1000–1200 nm; 0.11 μmol m⁻² s⁻¹ sr⁻¹, 900–1200 nm) are well above the minimum fluxes that support anoxygenic photosynthesis on Earth, albeit at longer wavelengths, as phototrophs near hydrothermal vents and deep in the Black Sea are associated with measurable abundances of bacteriochlorophylls *c* and *e*, respectively, which absorb between ~650 and 850 nm (Beatty *et al.*, 2005; Manske *et al.*, 2005).

Thus, for a hypothetical cloud-based microbiome at Venus, the absorption of solar irradiance with minimal UV flux, along with the absorption of continual thermal radiance from the surface/atmosphere, could together provide several unique advantages—that is, when compared to the selective pressures on Earth—including an expanded chemical and structural parameter space for photobiochemistry and novel diurnal cycles for photosynthesis. In turn, such attributes could hypothetically yield increased rates of microbial growth and division, and enhanced microbial diversity due to niche differentiation in an expanded photophysical habitat that includes solar flux during the day and illumination from below the clouds throughout the day and night. For photosynthesis, the anaerobic reduction of CO₂ could also hypothetically


proceed through electron donors arising from coupled sulfur and iron-based cycles (Limaye *et al.*, 2018), which incorporate nitrate reduction and hydrogen oxidation (Limaye *et al.*, 2018; Cockell *et al.*, 2021). Additionally, nitrite, a potential chemical in Venus' middle clouds (Mogul *et al.*, 2021), could contribute to nitrogen cycles on Venus (to yield nitrate) by serving as an electron donor for anoxygenic photosynthesis.

6. Habitability Limitations

In Venus' clouds, however, the potentials for phototrophy and/or chemotrophy are significantly constrained by the abundances of acid and water. Discussions of photosynthesis and photoassimilation are further restricted as simple carbohydrates ((CH₂O)_{*n*}) have yet to be reported in Venus' atmosphere. In contrast, abundances of species containing nitrogen, sulfur, phosphorus, chlorine, and iron have been reported in several studies (Knollenberg *et al.*, 1980; Krasnopolsky, 1989, 2013, 2017; Johnson and de Oliveira, 2019).

Acid abundances are regarded to be ≥75% w/w sulfuric acid in the aerosols in Venus' middle clouds with values peaking at ~98% w/w in the lower clouds (Kliore *et al.*, 1979; Grinspoon and Bullock, 2007; Cottini *et al.*, 2012; Krasnopolsky, 2015). From studies on Earth, solutions of ≥75% w/w sulfuric acid (~13–18 M H₂SO₄) are considered to exhibit water activities of ≤0.01 (Staples, 1981; Halls-worth *et al.*, 2021), where measured concentrations of free (or unreacted) water are ≤9 M (Walrafen *et al.*, 2000). Due to this water limitation, estimates of pH (-log [H₃O⁺]) greatly underestimate the functional acidity of the solution, or ability of a Brønsted-Lowry acid to transfer a proton. Instead, better measures of acidity are obtained when using the Hammett acidity factor (H_0), where relative degrees of acid strength can be interpreted using the typical pH scale (Hammett and Deyrup, 1932; Johnson *et al.*, 1969). As is indicated in Fig. 3, abundances of 75–99% w/w sulfuric acid relate to H_0 values of -6 to -11 across ~25–60°C (the temperatures found across 51.5 to 56 km) and correspond to acidity levels that are several orders of magnitude greater than those suggested by Grinspoon and Bullock (2007) (pH ~0 to -1), stoichiometric calculations of pH (pH -1.1 to -1.3), and Raman spectral measures of hydronium ion concentrations (pH -0.6 to -1.1) in 30–98% sulfuric acid solutions (Young *et al.*, 1959; Walrafen *et al.*, 2000).

In perspective, such extreme acidities and low water activities preclude any known biochemistry. On Earth, microbial growth is typically associated with water activities



of ≥ 0.6 (Rummel *et al.*, 2014) with the lowest known limit being 0.585 (Hallsworth *et al.*, 2021), which, in the context of sulfuric acid, correspond to $\leq 39\%$ w/w or (≤ 4.5 M), as indicated in the works of Staples (1981, Table 42) and Hallsworth *et al.* (2021). Microbial growth is further constrained by acidity as inferred from the current limits for cultivation and oxygenic and anoxygenic photosynthesis. For acidic cultivation, lower limits are under chemoheterotrophic anaerobic conditions at an apparent pH of -0.06 (60°C), or an added amount of 1.2 M H_2SO_4 ($H_0 -0.43$), by *Picrophilus*, an aerobic archaea (Schleper *et al.*, 1995). For oxygenic photosynthesis (22–25°C), the lower limits are pH ~ 0.3 ($H_0 \sim 0.1$; adapted from pH) by *Dunaliella acidophila* and pH 1.5 ($H_0 \sim 0.7$; adapted from pH) by *Chlamydomonas acidophila*, which are green algae (Gimmler and Weis, 1992; Gerloff-Elias *et al.*, 2005). For anoxygenic photosynthesis, lower limits for activity ($\sim 50^\circ\text{C}$) are at pH 2.2–2.4, or 5–7 mM H_2SO_4 ($H_0 = \text{pH} \sim 2$), where molecular measurements support anoxygenic photoassimilation of carbon and the presence of transcripts associated with bacteriochlorophyll biosynthesis (Hamilton *et al.*, 2019).

7. Chemical Considerations

Given these constraints, and as a chemical argument towards habitability, we suggest that treatments of Venus' aerosols, to date, may underestimate the presence of bisulfate and/or sulfate, which would directly impact the functional acidity and water activities of the aerosols. Bisulfate (HSO_4^-) and sulfate (SO_4^{2-}) are structurally and spectrally related to sulfuric acid (H_2SO_4), as they represent the respective conjugate bases of neutralized sulfuric acid. Aqueous solutions of bisulfate and sulfate salts, such as ammonium bisulfate and copper sulfate, are also optically refractive, which is relevant since Venus' clouds have been inferred to harbor $\sim 75\%$ w/w sulfuric acid due to similarities in refractive index (Hansen and Hovenier, 1974; Barstow *et al.*, 2012; Arney *et al.*, 2014). At Venus' cloud tops, refractive index (RI) values were found to be 1.44 ± 0.02 (550 nm) and 1.44 ± 0.01 (900 nm) by ground and orbiter-based polarimetry, respectively (Hansen and Hovenier, 1974; Kawabata *et al.*, 1980; Knollenberg *et al.*, 1980; Sato *et al.*, 1996). Similarly, ~ 75 – 76% w/w sulfuric acid solutions exhibit RI values

of ~ 1.44 per calculations (550 nm) at -23 and 15°C (Hansen and Hovenier, 1974) and measurements (556 nm) at 27°C (Palmer and Williams, 1975).

In comparison, solutions of ~ 35 – 62% w/w ammonium bisulfate (~ 3.7 – 7.5 M; NH_4HSO_4) at 25°C yield measured RI values (633 nm) of 1.38 – 1.41 (Tang and Munkelwitz, 1994) with $\sim 89\%$ w/w (~ 11 M) yielding an extrapolated RI of 1.44 ; as calculated when using the linear relationship ($R^2 > 0.99$) between measured RI and concentration of NH_4HSO_4 . Additionally, solutions of 1.5 M copper sulfate (CuSO_4) yield RI values (589 nm) of ~ 1.37 with minimal variation (1.365 – 1.370) between 25°C and 50°C (Nieto and Olcina, 1999). These similar RI values are noteworthy as models and *in situ* measures in Venus' atmosphere suggest the presence of ammonia/ammonium (Surkov *et al.*, 1974; Mogul *et al.*, 2021), copper (Schaefer and Fegley, 2004), and iron (Krasnopolsky, 2017), though relative abundances remain unconstrained. Theoretically, such chemicals could contribute to the neutralization of sulfuric acid to yield bisulfate and/or sulfate salts.

Measures of cloud layer RI were also obtained by the Pioneer Venus and Venera spacecraft (Knollenberg *et al.*, 1980; Ragent *et al.*, 1985). For this study, measures possessing defined uncertainties across the spacecraft measures (Pioneer Venus, 900 nm; Venera 9, 920 nm) were aggregated to yield averaged RI values of 1.41 ± 0.03 (propagated variance) across 51 – 55 km (~ 33 – 63°C) for the middle clouds, and 1.39 ± 0.04 (propagated variance) across 48 – 49 km (~ 79 – 87°C) for the lower clouds. When considering the propagated variances, the range in RI values for the middle clouds of 1.38 – 1.44 translated to abundances ranging ~ 40 – 85% w/w sulfuric acid, while RI values for the lower clouds (1.34 – 1.42) surprisingly translated to decreased values of < 25 – 70% w/w sulfuric acid—these estimates were obtained by (1) interpolation of measurements (909 nm) from the work of Palmer and Williams (1975) using the linear relationship ($R^2 > 0.99$) between RI across 25 – 75% w/w sulfuric acid and (2) coarsely assuming that the impacts of temperature were minimal, as suggested by the $< 10\%$ decrease in sulfuric acid densities between 25°C and 100°C , the $\leq 10\%$ change in calculated RI for 75% w/w sulfuric acid at -53°C and 27°C , and $\leq 10\%$ change in RI for 0.1 – 1.5 M copper sulfate across 25 – 50°C (interpreted as a proxy for sulfuric acid) (Massie, 1994; Nieto and Olcina, 1999; Haynes, 2016).

In summary (Fig. 3), these observations and calculations, when solely considering RI, suggest that aerosols in the middle clouds could harbor wide-ranging abundances of $\sim 40\%$ to 85% w/w sulfuric acid with corresponding acidities (H_0) of -2.5 to -8.5 . Our comparisons also show that solutions of bisulfate and/or sulfate salts—and quite likely mixtures with sulfuric acid—exhibit optical properties like those observed at Venus. For example, similar RI values are shared between Venus' middle clouds, 40 – 85% w/w sulfuric acid (~ 5.4 – 16 M), and 40 – 89% w/w ammonium bisulfate (~ 4.5 – 11 M).

8. Sulfuric Acid Abundances

Abundances of sulfuric acid in Venus' aerosols have additionally been inferred from radio occultation measurements, which involve microwave absorption and a dependence on the index of refraction of the clouds. Several studies indicate that sulfuric acid vapor is ~ 0 – 3 ppm in Venus' middle

clouds and ~ 1 – 10 ppm in the lower clouds (Kolodner and Steffes, 1998; Imamura *et al.*, 2017; Oschlisniok *et al.*, 2021). In fact, across 51 – 56 km (middle clouds), multiple measures of ~ 0 ppm vapor have been obtained from varying spacecraft including Akatsuki (Imamura *et al.*, 2017), Magellan across three orbits (Jenkins *et al.*, 1994), Venus Express (Oschlisniok *et al.*, 2012), and Mariner 10 (Kliore *et al.*, 1979).

To relate vapor and solution-phase abundances for sulfuric acid at Venus, Krasnopolsky (2015) performed thermodynamics calculations, which related ~ 2 ppm H_2SO_4 vapor to $\sim 98\%$ w/w sulfuric acid solutions at 50 km ($\sim 66^\circ\text{C}$ and 1 atm) and 30 ppm water vapor. Under these constraints, radio occultation measures of ~ 10 ppm H_2SO_4 vapor are then suggestive of the lower clouds harboring supersaturated sulfuric acid. Supersaturation, however, is contrary to the models of Esposito *et al.* (1983) and James *et al.* (1997), which suggest nominal saturating conditions between 50 and 45 km. Supersaturation is also inconsistent with gas chromatography measures of water vapor, which suggest abundances of 86% w/w sulfuric acid at 51.6 km (Oyama *et al.*, 1980). In addition, apparent RI values from the lower clouds (1.38 ± 0.04) are suggestive of $\leq 70\%$ w/w sulfuric acid (48 – 49 km). These comparisons, therefore, reveal potential uncertainties in calculated estimates for solution-phase abundances, where ~ 10 ppm H_2SO_4 vapor may instead relate to ≤ 98 – 100% w/w sulfuric acid, or nominally saturating conditions in the lower clouds.

We also estimated sulfuric acid abundances using vapor phase diagrams measured by Marti *et al.* (1997). These measures showed that solutions of $\sim 75\%$ w/w sulfuric acid (25°C , 1 atm) liberate ~ 3 ppm H_2SO_4 vapor at ~ 910 ppm water (2.9% relative humidity). In contrast, calculations by Krasnopolsky (2015) suggest that $\sim 75\%$ w/w sulfuric acid solutions liberate 0 ppm H_2SO_4 vapor at 2000 and $10,000$ ppm water under conditions relating to 54 km ($\sim 25^\circ\text{C}$, ~ 0.5 atm) and 50 km (66°C , 1 atm), respectively. These comparisons again suggest that thermodynamics calculations and measurements possess inherent uncertainties. Potential examples of these uncertainties include: (1) the ~ 1 -log difference between measured H_2SO_4 vapor pressures of Gmitro and Vermeulen (1964) and Ayers *et al.* (1980) (as compared in Marti *et al.* [1997]); (2) the ~ 3.5 -fold higher H_2SO_4 vapor pressures when comparing calculations of Krasnopolsky (2015) and measurements of Ayers *et al.* (1980); (3) the 5 – 6% lower H_2SO_4 activity values of Rard *et al.* (1976) when compared to those of Giauque *et al.* (1960); and (4) the $\sim 30\%$ uncertainty in Venus' cloud water vapor, as reported by Hallsworth *et al.* (2021), which for ~ 10 – 1000 ppm water correspond to ± 5 – 10% variances in solution-phase sulfuric acid abundances, as estimated using H_2O vapor phase diagrams of Gmitro and Vermeulen (1964).

Additional sources of potential uncertainty are the nonvolatile and volatile chemicals in Venus' atmosphere (Hoffman *et al.*, 1980; Krasnopolsky, 2013, 2017; Mogul *et al.*, 2021) that may dissolve within the aerosols (similar to our treatment of the unknown absorber) and decrease the partial vapor pressures of sulfuric acid and water (Lovelace *et al.*, 1921). For the lower clouds, uncertainties regarding the vertical variability in RI also include assumptions of particle shape. For example, Knollenberg *et al.* (1980) indicated that nephelometry data were best fit at 50 km (presumed altitude)

with very low RI values of ≤ 1.3 , which was indicative of limited backscatter due to conditions including absorbance of the nephelometer radiation by the particles and/or irregular (or non-spherical) particle shapes. Interestingly, intact cells that contain bacteriochlorophyll *a* (*Roseospirillum parvum*, Fig. 2C) maximally absorb near the nephelometer wavelengths of 900 and 920 nm; theoretically, therefore, biotic pigments could explain such observations, as could abiotic absorbers in the near infrared.

Alternatively, radio occultation measurements could support the presence of partly neutralized sulfuric acid. According to Marti *et al.* (1997), solutions of 55–77% w/w sulfuric acid (30°C) exhibit similar H₂SO₄ vapor pressures both before and after partial neutralization by ammonia gas (≤ 0.5 mole ratio of NH₃ to H₂SO₄). For example, for ~72% w/w sulfuric acid, vapor pressures of ~2–3 ppm H₂SO₄ marginally decrease to 0.9 ± 0.4 ppm after neutralization by 50% mole ratio of ammonia. As displayed in Fig. 3, the impacts of such neutralizations for Venus were estimated assuming 40–85% w/w sulfuric acid for the middle clouds (in accordance with RI measures) and a 0.5 mole ratio of ammonia. Such hypothetical neutralizations yield mixtures of 23–46% w/w ammonium bisulfate (~2.8–6.4 M) with 19–40% w/w sulfuric acid (~2.2–5.3 M)—concentrations assumed densities equal to sulfuric acid at 25°C. By means of measures on Earth, water activities of ≥ 0.6 (25°C) are obtained from solutions of 23–46% w/w ammonium bisulfate (Tang and Munkelwitz, 1994) or 19–40% w/w sulfuric acid (Staples, 1981).

Under these buffering conditions, in accordance with Fig. 3, adjusted acidity (H_0) values for the middle clouds range from -0.1 to -1.5 with slightly more basic values projected at higher temperatures (Johnson *et al.*, 1969). We found limited information in the literature regarding H_0 values for 1–10 M mixtures of bisulfate and sulfuric acid. To account for equimolar conjugate base, therefore, H_0 values (-1.0 and -2.4) at 19% and 40% w/w sulfuric acid (~2.2 and 5.3 M), as adapted from the work of Johnson *et al.* (1969), were corrected by using a ΔH_0 of 0.9, which yielded values of -0.1 to -1.5. As reported by Paul and Long (1957), solutions of 2–6 M H₂SO₄ and HClO₄ display similar H_0 profiles, while comparison of 6 M HClO₄ to mixtures of 3 M HClO₄ with 3 M NaClO₄ revealed a ΔH_0 of ~0.9. Hence, given that H₂SO₄ and HClO₄ display similar acidic behaviors, a ΔH_0 of ~0.9 was used to correct for 2–6 M solutions that contained equimolar sulfuric acid and bisulfate. For Venus' lower clouds, such treatments for 25–98% w/w sulfuric acid (lower and higher limits from RI and vapor phase calculations, respectively) yielded an adjusted H_0 range of +0.4 to -2 and water activities of ≥ 0.41 (for 46% w/w sulfuric acid). Thus, these hypothetical acidities and water activities, especially for the middle clouds, fall within the limits of terrestrial microbial growth.

9. Potential for Phototrophy

Our photophysical results indicate that transmitted solar irradiances in Venus' clouds (Table 1) are suitable for Earth-like phototrophy with similar photon fluxes (Table 2) across the visible and near-infrared (350–1200 nm) when compared to Earth's surface (ASTM G-173-03). As potential selective biological pressures, the solar irradiances in Venus' clouds could also be more favorable due to ~87% less flux in

UV-A, with negligible flux in UV-B and UV-C. Hypothetical phototrophs, moreover, could appear brown/maroon in the middle and lower cloud after absorption of light transmitting through Venus' signature phototrophic windows (Table 2), inclusive of maximal absorption at the wavelength of peak solar irradiance (608 nm). Lastly, our calculations suggest that thermal emissions from below Venus' clouds could sustain phototrophy. Hence, our analyses are supportive of a photophysical habitat that includes diurnal input of solar energy through the cloud tops, along with the persistent input of lower-power and lower-flux thermal energy from below the clouds.

10. Potential for Habitability

When considering chemical limits to habitability, our interpretations of measured refractive index and radio occultation data suggest that aerosols in the middle clouds could harbor neutralized forms of sulfuric acid, such as ammonium bisulfate. Such interpretations of partly neutralized sulfuric acid are parallel to those of Rimmer *et al.* (2021), who suggest that aerosols with pH values of ~1 (due to the presence of substantial base) yield water and SO₂ vapors that match Venus observations. When assuming the presence of mixtures of sulfuric acid and ammonium bisulfate (0.5 mole ratio), our results suggest that the aerosols could harbor water activity (≥ 0.6) and acidity values (H_0 -0.1 to -1.5) that lie within the limits of terrestrial acidic cultivation ($H_0 \geq -0.4$). The acidity limits are also tantalizingly close to the limits for oxygenic photosynthesis ($\geq H_0$ 0.1). However, for anoxygenic photosynthesis ($\geq H_0$ 2), acidity remains to be a constraining variable, which suggests the need for additional basicity, similar to the model of Rimmer *et al.* (2021), to yield terrestrial-type conditions.

When considering water, a consensus of low water vapor abundances (~20–100 ppm) are obtained through optical measures (Johnson and de Oliveira, 2019). In contrast, *in situ* measures suggest ~200–2000 ppm (Vega 1, and Venera 9, 10, 13, and 14) in the middle/lower clouds (58–48 km) and ~5000 ppm (Pioneer Venus) just below the clouds (41.7 km) (Hoffman *et al.*, 1980; Oyama *et al.*, 1980; Surkov *et al.*, 1983, 1987; Von Zahn *et al.*, 1983). Apart from instrumental issues, we posit that these high values represent partial sampling of the aerosols, which hint at larger solution-phase reservoirs of water, per se, that may harbor solutes, as noted, that could lower the partial pressures of water and/or sulfuric acid. We speculate, therefore, that microbial survival in an aerosolized and water-restricted environment could include bio/chemical strategies to reduce vapor pressure and loss of solution-phase water, similar to microbial strategies associated with freezing point and temperature depression (Scotter *et al.*, 2006; Möhlmann, 2012), inhibition of ice formation (Krembs and Deming, 2008; Raymond *et al.*, 2008), and cloud condensation (Lazaridis, 2019).

11. Conclusions

Solar irradiances calculated across Venus' clouds support the potential for Earth-like phototrophy, while treatment of Venus' aerosols as containing neutralized sulfuric acid favors a habitable zone in the clouds. To date, most studies suggest that the aerosols in Venus' middle and lower clouds

harbor high acidities (Seager *et al.*, 2020; Limaye *et al.*, 2021) with commensurate low water activities or availabilities (Cockell, 1999; Izenberg *et al.*, 2020; Seager *et al.*, 2020; Cockell *et al.*, 2021; Hallsworth *et al.*, 2021; Limaye *et al.*, 2021). In contrast, as presented in this study, alternative interpretations to *in situ* measures yield potentially habitable conditions with water activities (≥ 0.585) and buffered acidities (Hammett acidity factor, H_0 -0.1 to -1.5) that lie within the limits of terrestrial microbial growth.

Venus' clouds, therefore, may not be as inhospitable as suggested by the models of Krasnopolsky (2015) and Hallsworth *et al.* (2021) given the lack of observational constraints for Venus' middle and lower clouds and since differing chemicals may share RI and microwave absorption properties. Looking ahead, mass spectrometers amenable to sampling vapors of sulfuric and phosphoric acids (Wurz *et al.*, 2012; Ren *et al.*, 2020), ionized chemical species (Baines *et al.*, 2021), and/or sublimated salts (Hänni *et al.*, 2019) could help in detailing the acid, conjugate base, and water abundances in the aerosols, and thereby directly address the potential for habitability in Venus' clouds. Furthermore, when considering potential life-detection strategies, our results suggest that the wavelength range of ~ 558 – 608 nm, which accounts for uncertainties with the unknown absorbers, may serve as a reasonable excitation window for fluorescence spectroscopy.

Author Contributions

All authors (RM, SSL, YJL, and MP) contributed to the acquisition, analysis, or interpretation of the data, and drafting of critical revisions/reports of the work. The primary investigator and corresponding author is RM. Spectra were calculated by YJL. Calculation of photon flux and PPF, and assignment of spectral regions for phototrophy, were performed by RM. Refractive index, acidity, and water activities were calculated or estimated by RM and MP.

Acknowledgments

We thank Jaime A. Cordova for editing of the manuscript. RM acknowledges support from the National Aeronautics and Space Administration (NASA) Research Opportunities in Space and Earth Sciences (NNH18ZDA001N). SSL acknowledges support from NASA (NNX16AC79G). YJL acknowledges funding from EU Horizon 2020 MSCA-IF No. 841432.

References

Arney G, Meadows V, Crisp D, *et al.* (2014) Spatially resolved measurements of H₂O, HCl, CO, OCS, SO₂, cloud opacity, and acid concentration in the Venus near-infrared spectral windows. *J Geophys Res Planets* 119: 1860–1891.

Ayers G, Gillett R, and Gras J (1980) On the vapor pressure of sulfuric acid. *Geophys Res Lett* 7:433–436.

Baines KH, Nikolić D, Cutts JA, *et al.* (2021) Investigation of Venus cloud aerosol and gas composition including potential biogenic materials via an aerosol-sampling instrument package. *Astrobiology* 21:1316–1323.

Barstow J, Tsang C, Wilson C, *et al.* (2012) Models of the global cloud structure on Venus derived from Venus Express observations. *Icarus* 217:542–560.

Beatty JT, Overmann J, Lince MT, *et al.* (2005) An obligately photosynthetic bacterial anaerobe from a deep-sea hydrothermal vent. *Proc Natl Acad Sci USA* 102:9306–9310.

Cockell CS (1999) Life on Venus. *Planet Space Sci* 47:1487–1501.

Cockell CS, Higgins PM, and Johnstone AA (2021) Biologically available chemical energy in the temperate but uninhabitable venusian cloud layer: what do we want to know? *Astrobiology* 21:1224–1236.

Cottini V, Ignatiev N, Piccioni G, *et al.* (2012) Water vapor near the cloud tops of Venus from Venus Express/VIRTIS dayside data. *Icarus* 217:561–569.

Crisp D (1986) Radiative forcing of the Venus mesosphere: I. Solar fluxes and heating rates. *Icarus* 67:484–514.

Crisp D, Sinton W, Hodapp K-W, *et al.* (1989) The nature of the near-infrared features on the Venus night side. *Science* 246:506–509.

Ekonomov A, Moroz V, Moshkin B, *et al.* (1984) Scattered UV solar radiation within the clouds of Venus. *Nature* 307:345–347.

Esposito L, Knollenberg R, Marov M, *et al.* (1983) The clouds and hazes of Venus. In *Venus*, edited by DM Hunten, University of Arizona Press, Tucson, pp 484–564.

Gerloff-Elias A, Spijkerman E, and Pröschold T (2005) Effect of external pH on the growth, photosynthesis and photosynthetic electron transport of *Chlamydomonas acidophila* Negoro, isolated from an extremely acidic lake (pH 2.6). *Plant Cell Environ* 28:1218–1229.

Giauque W, Hornung E, Kunzler J, *et al.* (1960) The thermodynamic properties of aqueous sulfuric acid solutions and hydrates from 15 to 300°K. *J Am Chem Soc* 82: 62–70.

Gimmler H and Weis U (1992) *Dunaliella acidophila*—life at pH 1.0. In *Dunaliella: Physiology, Biochemistry, and Biotechnology*, edited by A Ben-Amotz and M Avron, CRC Press, Boca Raton, FL, pp 99–134.

Glaeser J and Overmann J (1999) Selective enrichment and characterization of *Roseospiillum parvum*, gen. nov. and sp. nov., a new purple nonsulfur bacterium with unusual light absorption properties. *Arch Microbiol* 171:405–416.

Gmitro JJ and Vermeulen T (1964) Vapor-liquid equilibria for aqueous sulfuric acid. *AIChE Journal* 10:740–746.

Govindjee R (1974) The absorption of light in photosynthesis. *Sci Am* 231:68–87.

Grant R and Slusser J (2005) Estimation of ultraviolet-A irradiance from measurements of 368-nm spectral irradiance. *J Atmos Ocean Technol* 22:1853–1863.

Grimspoon D and Bullock M (2007) Astrobiology and Venus exploration. In *Exploring Venus as a Terrestrial Planet*, edited by LW Esposito, ER Stofan, TE and Cravens, American Geophysical Union, Washington, DC, pp 191–206.

Hallsworth JE, Koop T, Dallas TD, *et al.* (2021) Water activity in Venus's uninhabitable clouds and other planetary atmospheres. *Nat Astron* 5:665–675.

Hamilton TL, Bennett AC, Murugapiran SK, *et al.* (2019) Anoxygenic phototrophs span geochemical gradients and diverse morphologies in terrestrial geothermal springs. *mSystems* 4, doi:10.1128/mSystems.00498-19.

Hammett LP and Deyrup AJ (1932) A series of simple basic indicators. I. The acidity functions of mixtures of sulfuric and perchloric acids with water. *J Am Chem Soc* 54:2721–2739.

Hänni N, Gasc S, Etter A, *et al.* (2019) Ammonium salts as a source of small molecules observed with high-resolution

- electron-impact ionization mass spectrometry. *J Phys Chem A* 123:5805–5814.
- Hansen JE and Hovenier J (1974) Interpretation of the polarization of Venus. *J Atmos Sci* 31:1137–1160.
- Haynes WM (2016) *CRC Handbook of Chemistry and Physics*. CRC Press, Boca Raton, FL.
- Hoffman JH, Hodges RR, Donahue TM, *et al.* (1980) Composition of the Venus lower atmosphere from the Pioneer Venus mass spectrometer. *J Geophys Res Space Phys* 85:7882–7890.
- Holtrop T, Huisman J, Stomp M, *et al.* (2021) Vibrational modes of water predict spectral niches for photosynthesis in lakes and oceans. *Nat Ecol Evol* 5:55–66.
- Hummel JR, Shettle EP, and Longtin DR (1988) *A New Background Stratospheric Aerosol Model for Use in Atmospheric Radiation Models*, AFGL-TR-88-0166, Scientific Report No.8, Hanscom Air Force Base, Massachusetts. Available online at <https://apps.dtic.mil/sti/pdfs/ADA210110.pdf>
- Ignatiev N, Moroz V, Moshkin B, *et al.* (1997) Water vapour in the lower atmosphere of Venus: a new analysis of optical spectra measured by entry probes. *Planet Space Sci* 45:427–438.
- Imamura T, Ando H, Tellmann S, *et al.* (2017) Initial performance of the radio occultation experiment in the Venus orbiter mission Akatsuki. *Earth Planets Space* 69:1–11.
- Imhoff JF (2017) Anoxygenic phototrophic bacteria from extreme environments. In *Modern Topics in the Phototrophic Prokaryotes*, edited by P Hallenbeck, Springer, Cham, Switzerland, pp 427–480.
- Imhoff JF and Trüper HG (1977) *Ectothiorhodospira halochloris* sp. nov., a new extremely halophilic phototrophic bacterium containing bacteriochlorophyll *b*. *Arch Microbiol* 114:115–121.
- Izenberg NR, Gentry DM, Smith DJ, *et al.* (2020) The Venus life equation. *Astrobiology* 21:1305–1315.
- Jenkins JM, Steffes PG, Hinson DP, *et al.* (1994) Radio occultation studies of the Venus atmosphere with the Magellan spacecraft: 2. Results from the October 1991 experiments. *Icarus* 110:79–94.
- Johnson CD, Katritzky A, and Shapiro S (1969) Temperature variation of the H_0 acidity function in aqueous sulfuric acid solution. *J Am Chem Soc* 91:6654–6662.
- Johnson NM and de Oliveira MR (2019) Venus atmospheric composition *in situ* data: a compilation. *Earth Space Sci* 6: 1299–1318.
- Kawabata K, Coffeen DL, Hansen JE, *et al.* (1980) Cloud and haze properties from Pioneer Venus polarimetry. *J Geophys Res* 85:8129–8140.
- Kiang NY, Segura A, Tinetti G, *et al.* (2007a) Spectral signatures of photosynthesis. II. Coevolution with other stars and the atmosphere on extrasolar worlds. *Astrobiology* 7:252–274.
- Kiang NY, Siefert J, and Blankenship RE (2007b) Spectral signatures of photosynthesis. I. Review of Earth organisms. *Astrobiology* 7:222–251.
- Kliore AJ, Elachi C, Patel IR, *et al.* (1979) Liquid content of the lower clouds of Venus as determined from Mariner 10 radio occultation. *Icarus* 37:51–72.
- Knollenberg R, Travis L, Tomasko M, *et al.* (1980) The clouds of Venus: a synthesis report. *J Geophys Res Space Phys* 85: 8059–8081.
- Kolodner MA, and Steffes PG (1998) The microwave absorption and abundance of sulfuric acid vapor in the Venus atmosphere based on new laboratory measurements. *Icarus* 132:151–169.
- Krasnopolsky V (1989) Vega mission results and chemical composition of venusian clouds. *Icarus* 80:202–210.
- Krasnopolsky VA (2013) S_3 and S_4 abundances and improved chemical kinetic model for the lower atmosphere of Venus. *Icarus* 225:570–580.
- Krasnopolsky VA (2015) Vertical profiles of H_2O , H_2SO_4 , and sulfuric acid concentration at 45–75 km on Venus. *Icarus* 252:327–333.
- Krasnopolsky VA (2017) On the iron chloride aerosol in the clouds of Venus. *Icarus* 286:134–137.
- Krembs C and Deming JW (2008) The role of exopolymers in microbial adaptation to sea ice. In *Psychrophiles: From Biodiversity to Biotechnology*, Springer, Cham, Switzerland, pp 247–264.
- Lazaridis M (2019) Bacteria as cloud condensation nuclei (CCN) in the atmosphere. *Atmosphere* 10, doi:10.3390/atmos10120786.
- Lee YJ, Sagawa H, Haus R, *et al.* (2016) Sensitivity of net thermal flux to the abundance of trace gases in the lower atmosphere of Venus. *J Geophys Res Planets* 121:1737–1752.
- Lee YJ, Jessup K-L, Perez-Hoyos S, *et al.* (2019) Long-term variations of Venus's 365 nm albedo observed by Venus Express, Akatsuki, MESSENGER, and the Hubble Space Telescope. *Astron J* 158, doi:10.3847/1538-3881/ab3120.
- Lehmer OR, Catling DC, Parenteau MN, *et al.* (2018) The productivity of oxygenic photosynthesis around cool, M dwarf stars. *Astrophys J* 859, doi:10.3847/1538-4357/aac104.
- Limaye SS, Mogul R, Smith DJ, *et al.* (2018) Venus' spectral signatures and the potential for life in the clouds. *Astrobiology* 18:1181–1198.
- Limaye SS, Mogul R, Baines KH, *et al.* (2021) Venus, an astrobiology target. *Astrobiology* 21:1163–1185.
- Longobardo A, Palomba E, Zinzi A, *et al.* (2012) Limb darkening study using Venus nightside infrared spectra from VIRTIS-Venus Express data. *Planet Space Sci* 69: 62–75.
- Lovelace B, Frazer J, and Sease V (1921) The lowering of the vapor pressure of water at 20° produced by dissolved potassium chloride. *J Am Chem Soc* 43:102–110.
- Madigan MT (2003) Anoxygenic phototrophic bacteria from extreme environments. *Photosynth Res* 76:157–171.
- Manske AK, Glaeser J, Kuypers MMM, *et al.* (2005) Physiology and phylogeny of green sulfur bacteria forming a monospecific phototrophic assemblage at a depth of 100 meters in the Black Sea. *Appl Environ Microbiol* 71:8049–8060.
- Marschall E, Jogler M, Henßge U, *et al.* (2010) Large-scale distribution and activity patterns of an extremely low-light-adapted population of green sulfur bacteria in the Black Sea. *Environ Microbiol* 12:1348–1362.
- Marti JJ, Jefferson A, Cai XP, *et al.* (1997) H_2SO_4 vapor pressure of sulfuric acid and ammonium sulfate solutions. *J Geophys Res Atmos* 102:3725–3735.
- Massie S (1994) Indices of refraction for the HITRAN compilation. *J Quant Spectrosc Radiat Transf* 52:501–513.
- Meadows VS and Crisp D (1996) Ground-based near-infrared observations of the Venus nightside: the thermal structure and water abundance near the surface. *J Geophys Res Planets* 101:4595–4622.
- Mogul R, Limaye SS, Way M, *et al.* (2021) Venus' mass spectra show signs of disequilibria in the middle clouds. *Geophys Res Lett* 48, doi:10.1029/2020GL091327.

- Möhlmann D (2012) Widen the belt of habitability! *Orig Life Evol Biosph* 42:93–100.
- Moroz V (1983) Summary of preliminary results of the Venera 13 and Venera 14 missions. In *Venus*, edited by DM Hunten, University of Arizona Press, Tucson, AZ, pp 45–68.
- Moroz V, Moshkin B, Ekonomov A, *et al.* (1982) The Venera 13 and Venera 14 spectrophotometry experiments. *Pisma v Astronomicheskii Zhurnal* 8:404–410.
- Moroz V, Ekonomov A, Moshkin B, *et al.* (1985) Solar and thermal radiation in the Venus atmosphere. *Adv Space Res* 5: 197–232.
- Mueller N, Helbert J, Hashimoto G, *et al.* (2008) Venus surface thermal emission at 1 μm in VIRTIS imaging observations: evidence for variation of crust and mantle differentiation conditions. *J Geophys Res Planets* 113, doi:10.1029/2008JE003118.
- Nieto J and Olcina PV (1999) Study of electrochemical properties of the ternary system water-dimethylsulfoxide-copper sulphate. *Portugalia* 17:139–147.
- Nishio J (2000) Why are higher plants green? Evolution of the higher plant photosynthetic pigment complement. *Plant Cell Environ* 23:539–548.
- Orellana G, Gómez-Silva B, Urrutia M, *et al.* (2020) UV-A irradiation increases scytonemin biosynthesis in cyanobacteria inhabiting halites at Salar Grande, Atacama Desert. *Microorganisms* 8, doi:10.3390/microorganisms8111690.
- Oschlisniok J, Häusler B, Pätzold M, *et al.* (2012) Microwave absorptivity by sulfuric acid in the Venus atmosphere: first results from the Venus Express Radio Science experiment VeRa. *Icarus* 221:940–948.
- Oschlisniok J, Häusler B, Pätzold M, *et al.* (2021) Sulfuric acid vapor and sulfur dioxide in the atmosphere of Venus as observed by the Venus Express radio science experiment VeRa. *Icarus* 362, doi:10.1016/j.icarus.2021.114405.
- Overmann J and Pfennig N (1989) *Pelodictyon phaeocla-thratiforme* sp. nov., a new brown-colored member of the Chlorobiaceae forming net-like colonies. *Arch Microbiol* 152:401–406.
- Overmann J, Cypionka H, and Pfennig N (1992) An extremely low-light adapted phototrophic sulfur bacterium from the Black Sea. *Limnol Oceanogr* 37:150–155.
- Oyama VI, Carle GC, Woeller F, *et al.* (1980) Pioneer Venus gas chromatography of the lower atmosphere of Venus. *J Geophys Res* 85:7891–7902.
- Palmer KF and Williams D (1975) Optical constants of sulfuric acid; application to the clouds of Venus? *Appl Opt* 14:208–219.
- Paul M and Long F (1957) H_0 and related indicator acidity function. *Chem Rev* 57:1–45.
- Pérez-Hoyos S, Sánchez-Lavega A, García-Muñoz A, *et al.* (2018) Venus upper clouds and the UV absorber from MESSENGER/MASCS observations. *J Geophys Res Atmos* 123, doi:10.1002/2017JE005406.
- Piacentini RD, Cede A, and Bárcena H (2003) Extreme solar total and UV irradiances due to cloud effect measured near the summer solstice at the high-altitude desertic plateau Puna of Atacama (Argentina). *J Atmos Sol Terr Phys* 65:727–731.
- Ragent B, Esposito LW, Tomasko MG, *et al.* (1985) Particulate matter in the Venus atmosphere. *Adv Space Res* 5:85–115.
- Rard JA, Habenschuss A, and Spedding FH (1976) A review of the osmotic coefficients of aqueous sulfuric acid at 25°C. *J Chem Eng Data* 21:374–379.
- Raymond JA, Christner BC, and Schuster SC (2008) A bacterial ice-binding protein from the Vostok ice core. *Extremophiles* 12:713–717.
- Ren Z, Guo M, Cheng Y, *et al.* (2020) Design of a compact time-of-flight mass spectrometer for space application. *J Am Soc Mass Spectrom* 31:434–440.
- Rimmer PB, Jordan S, Constantinou T, *et al.* (2021) Hydroxide salts in the clouds of Venus: their effect on the sulfur cycle and cloud droplet pH. *Planet Sci J* 2, doi:10.3847/PSJ/ac0156.
- Rummel JD, Beatty DW, Jones MA, *et al.* (2014) A new analysis of Mars “special regions”: findings of the second MEPAG Special Regions Science Analysis Group (SR-SAG2). *Astrobiology* 14:887–968.
- Sato M, Travis LD, and Kawabata K (1996) Photopolarimetry analysis of the Venus atmosphere in polar regions. *Icarus* 124:569–585.
- Schaefer L and Fegley B Jr (2004) Heavy metal frost on Venus. *Icarus* 168:215–219.
- Schleper C, Puehler G, Holz I, *et al.* (1995) *Picrophilus* gen. nov., fam. nov.: a novel aerobic, heterotrophic, thermo-acidophilic genus and family comprising archaea capable of growth around pH 0. *J Bacteriol* 177:7050–7059.
- Scotter AJ, Marshall CB, Graham LA, *et al.* (2006) The basis for hyperactivity of antifreeze proteins. *Cryobiol* 53: 229–239.
- Seager S, Petkowski JJ, Gao P, *et al.* (2020) The venusian lower atmosphere haze as a depot for desiccated microbial life: a proposed life cycle for persistence of the venusian aerial biosphere. *Astrobiology* 21:1206–1223.
- Staples BR (1981) Activity and osmotic coefficients of aqueous sulfuric acid at 298.15 K. *J Phys Chem Ref Data* 10:779–798.
- Stomp M, Huisman J, Stal LJ, *et al.* (2007) Colorful niches of phototrophic microorganisms shaped by vibrations of the water molecule. *ISME J* 1:271–282.
- Surkov YA, Andreichikov B, and Kalinkina O (1974) Ammonia content in the atmosphere of Venus according to data from the Venus-8 probe. *Soviet Physics Doklady* 18:679.
- Surkov IA, Ivanova V, Pudov A, *et al.* (1983) Water vapour content in the Venus atmosphere from Venera 13 and Venera 14 data. *Kosmicheskie Issledovaniia* 21:231–235.
- Surkov IA, Shcheglov O, Ryvkin M, *et al.* (1987) Water vapour distribution in the middle and lower Venus atmosphere. *Kosmicheskie Issledovaniia* 25:678–690.
- Takaichi S, Maoka T, Hanada S, *et al.* (2001) Dihydroxyglycopenone diglucoside diesters: a novel class of carotenoids from the phototrophic purple sulfur bacteria *Halorhodospira abdelmalekii* and *Halorhodospira halochloris*. *Arch Microbiol* 175:161–167.
- Tang I and Munkelwitz H (1994) Water activities, densities, and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance. *J Geophys Res Atmos* 99: 18801–18808.
- Thornber J, Cogdell R, Seftor R, *et al.* (1980) Further studies on the composition and spectral properties of the photochemical reaction centers of bacteriochlorophyll b-containing bacteria. *Biochim Biophys Acta Bioenergetics* 593:60–75.
- Titov DV, Ignatiev NI, McGouldrick K, *et al.* (2018) Clouds and hazes of Venus. *Space Sci Rev* 214, doi:10.1007/s11214-018-0552-z.
- Tomasko M, Doose L, Smith PH, *et al.* (1980) Measurements of the flux of sunlight in the atmosphere of Venus. *J Geophys Res Space Phys* 85:8167–8186.

- Von Zahn U, Kumar S, Niemann H, *et al.* (1983) Composition of the Venus atmosphere. In *Venus*, edited by DM Hunten, University of Arizona Press, Tucson, AZ, pp 299–430.
- Walrafen G, Yang W-H, Chu Y, *et al.* (2000) Structures of concentrated sulfuric acid determined from density, conductivity, viscosity, and Raman spectroscopic data. *J Solution Chem* 29:905–936.
- Warren-Rhodes KA, Rhodes KL, Pointing SB, *et al.* (2006) Hypolithic cyanobacteria, dry limit of photosynthesis, and microbial ecology in the hyperarid Atacama Desert. *Microb Ecol* 52:389–398.
- Wolstencroft R and Raven JA (2002) Photosynthesis: likelihood of occurrence and possibility of detection on Earth-like planets. *Icarus* 157:535–548.
- Wurz P, Abplanalp D, Tulej M, *et al.* (2012) A neutral gas mass spectrometer for the investigation of lunar volatiles. *Planet Space Sci* 74:264–269.
- Yacobi YZ, Köhler J, Leunert F, *et al.* (2015) Phycocyanin-specific absorption coefficient: eliminating the effect of chlorophylls absorption. *Limnol Oceanogr Methods* 13:e10015.
- Young T, Maranville L, and Smith H (1959) Chapter 4. In *Structure of Electrolytic Solutions*, edited by WJ Harner, Wiley, New York, pp 35–63.

Address correspondence to:

Rakesh Mogul
Chemistry & Biochemistry Department
California State Polytechnic University Pomona
3801 W. Temple Ave.
Pomona, CA 91768
USA

E-mail: rmogul@cpp.edu

Submitted 6 March 2021

Accepted 27 August 2021

Associate Editor: Sara Seager

Abbreviations Used

H_0 = Hammett acidity factor
PPFD = photosynthetic photon flux density
RI = refractive index
SZA = solar zenith angle
TOA = top of the atmosphere