SSL4EO-S12: A Large-Scale Multi-Modal, Multi-Temporal Dataset for Self-Supervised Learning in Earth Observation

Yi Wang, Student Member, IEEE, Nassim Ait Ali Braham, Zhitong Xiong, Member, IEEE, Chenying Liu, Conrad M Albrecht, Member, IEEE, Xiao Xiang Zhu, Fellow, IEEE

Abstract—Self-supervised pre-training bears potential to generate expressive representations from large-scale Earth observation (EO) data without human annotation. However, most existing pre-training in the field is based on ImageNet or medium-size, labeled remote sensing (RS) datasets. In this paper, we share an unlabeled dataset SSL4EO-S12:

Self-Supervised Learning for Earth Observation - Sentinel-1/2

to assemble a large-scale, global, multimodal, and multi-seasonal corpus of satellite imagery. We demonstrate SSL4EO-S12 to succeed in self-supervised pre-training for a set of representative methods: MoCo-v2, DINO, MAE and data2vec, and multiple downstream applications including scene classification, semantic segmentation and change detection. Our benchmark results prove the effectiveness of SSL4EO-S12 compared to existing datasets. The dataset, related source code, and pre-trained models are available at https://github.com/zhu-xlab/SSL4EO-S12.

Index Terms-Self-supervised learning, dataset, benchmark.

I. INTRODUCTION

S ELF-SUPERVISED learning (SSL) has attracted wide attention in the remote sensing (RS) community with the ability to learn generic representations from unlabeled data. Numerous studies in the literature have proven the potential of SSL in Earth observation (EO) beyond natural images [1]. Despite the focus SSL for EO receives, only limited effort is dedicated to providing large-scale datasets and benchmarks for pre-training. On the one hand, relying on computer vision datasets like ImageNet [2] is not a preferred option due to the domain gap. On the other hand, while RS datasets like SEN12MS [3] or SeCo [4] exist, they are limited by geospatial overlap, sparse geographical distribution, or lack diversity in seasonal or multimodal information. Therefore, big EOspecific datasets for unsupervised pre-training are necessary to be developed.

In this work, we introduce a large-scale, globally distributed, multi-temporal and multi-sensor dataset *SSL4EO-S12: Self-Supervised Learning for Earth Observation - Sentinel-1/2*. The dataset samples 250K locations around the globe, each providing Sentinel-2 L1C, Sentinel-2 L2A, and Sentinel-1 GRD images with four snapshots from different seasons (in

Y. Wang, N. A. A. Braham, C. Liu are with the Chair of Data Science in Earth Observation, Technical University of Munich (TUM), and the Remote Sensing Technology Institute, German Aerospace Center (DLR). Z. Xiong, X. X. Zhu are with the Chair of Data Science in Earth Observation, Technical University of Munich (TUM). C. M. Albrecht is with the Remote Sensing Technology Institute, German Aerospace Center (DLR). total 3 million $2640m \times 2640m$ patches). Additionally, we guarantee optimal geospatial coverage by avoiding the overlap of the randomly sampled locations. This renders SSL4EO-S12 the largest and most generic multi-spectral/SAR dataset in the RS literature [5].

We demonstrate the potential of SSL4EO-S12 dataset through a series of extensive experiments. Specifically, we evaluate four representative SSL algorithms—namely: MoCo [6], DINO [7], MAE [8], and data2vec [9]—on three different downstream tasks: scene classification, semantic segmentation and change detection. Our results indicate that pre-training on SSL4EO-S12 improves the downstream performance compared to existing datasets. Moreover, our ablation studies prove the benefits of RS-specific data augmentations including multisensor, multi-temporal and atmospheric correction.

II. RELATED WORK

Self-supervised learning Over the past years, self-supervised learning (SSL) has reached important milestones in computer vision, especially through contrastive methods with joint-embedding architectures. These methods get trained to promote similarity between augmented views of the same input, thereby enforcing invariance to data augmentation. Several families of such methods emerge: 1) contrasting negative samples for which the representations are encouraged to be dissimilar [6]; 2) Knowledge distillation between an asymmetric teacher-student network [7]; 3) redundancy reduction among the embedding dimensions; 4) clustering latent features to common prototypes from different views [10]. Meanwhile, recent developments in masked image modeling (MIM) reveal promising results in generative methods, which reconstruct the masked input at pixel-[8] or feature-[9] level.

We benchmark four representative methods MoCo [6], DINO [7], MAE [8], and data2vec [9] on the proposed dataset. This way, we cover a reasonably diverse set of representative methods from different categories: MoCo contrasts negative samples, DINO represents a distillation method, MAE is based on masked reconstruction, and data2Vec combines the masking mechanism with a joint-embedding architecture.

Pre-training datasets Pre-trained models on ImageNet are widely used for various computer vision tasks. However, this is less appropriate in the context of RS: 1) RS images are not object-centric; 2) there exist various types of sensors in RS; 3) temporal effects yield variations on the ground surface.

including reprinting/republishing this material for advertising or promotional purposes, creating new collective works, for resale or redistribution to servers or lists, or reuse of any copyrighted component of this work in other works.

^{© 2023} IEEE. Personal use of this material is permitted. Permission from IEEE must be obtained for all other uses, in any current or future media,



Fig. 1. Sample images of SSL4EO-S12 dataset assembled.

 TABLE I

 Summary of popular medium-resolution pre-training datasets in remote sensing. LC: land cover.

dataset	spatial cover	temporal cover	modality	overlap	patch size	# of locations	# of patches
BigEarthNet [13]	Europe	1 timestamp	SAR/optical	no	120×120	600K	1.2M
SEN12MS [3]	global	1 timestamp	SAR/optical/LC	yes	256×256	180K	540K
SeCo [4]	global	5 timestamps	optical	yes	264×264	200K	1M
SSL4EO-S12	global	4 timestamps	SAR/optical*2	\sim no	264×264	250K	3M

Therefore, EO-specific datasets are needed to provide the above in-domain knowledge. The literature has proven the benefits of pre-training on existing labeled RS datasets [11, 12], yet there are limitations such as class bias, and temporal and geographical coverage.

Consequently, there is a need for large-scale pre-training datasets in RS. Two datasets closely related to our efforts are SEN12MS [3] and SeCo [4]. However, SEN12MS is limited by temporal coverage, SeCo has only optical data, and both datasets contain strongly overlapping patches which limit the geospatial coverage. With the above in mind, our proposed SSL4EO-S12 dataset provides an improved spatio-temporal coverage by sampling more locations and removing overlapping patches, enclosing multiple seasons, and including Sentinel-1 as well as two Sentinel-2 products (Table I).

III. SSL4EO-S12 DATASET

A. Data curation & assembly

The SSL4EO-S12 dataset (Figure 1) exploits openly available SAR/optical satellite data collected by the European Space Agency's Sentinel mission. Following a well-organized baseline provided by SeCo [4], we utilize the Google Earth Engine [14] to download and process the data. We filter image patches to retrieve from the 10,000 most populated cities¹ in the world (top-10k) to guarantee reasonable global coverage. To obtain diverse land cover, we sample 251,079 locations close by the cities following a Gaussian distribution peaking at the city center and standard deviation of 50km—assuming most of the variability cast to the downtown and suburbs of cities [4]. At each location, we download 4 images drawn from four annual seasons to capture seasonal variation. We search for Sentinel-2 tiles with a cloud coverage lower than 10%. We also filter out most overlapping patches with an efficient grid search strategy. In total, we obtain about one million S1-GRD/S2-L1C/S2-L2A image triplets.

Data identification. The collection of SSL4EO-S12 differs from SeCo mainly by introducing overlap filtering and multiple sensors (**bold** below). The workflow is shown as follows:

- 1) Uniformly sample one city from top-10k populated cities;
- Sample one location from a Gaussian distribution with a standard deviation of 50km around the city center;
- 3) Check if a $2640m \times 2640m$ image patch centered around that location has significant overlap with previous patches. If not, continue to 4, otherwise return to 1;
- 4) For a 30-day interval around four reference dates (Mar 20, Jun 21, Sep 22, Dec 21) in 2021 (additionally look for 2020 as a buffer), check if there exist Sentinel-2 tiles with less than 10% of cloud coverage (for both L1C and L2A) and corresponding Sentinel-1 GRD tiles;
- 5) If there exist valid Sentinel-1/2 tiles close to all the four dates, process and download them into curated image patches, otherwise return to 1.

Overlap filtering. A simple way to check significant overlap between two patches is to calculate the distance between the two centers. If the distance is smaller than 3/4 the width of a patch, there is a non-negligible overlap ($i_225\%$). Naively, we need to execute this computation for every new patch relative to all existing patches. However, this becomes inefficient when the number of patches grows large, 250k+ for us. Therefore,



Fig. 2. Geographical distribution of SSL4EO-S12 dataset.



Fig. 3. Image patches without (left) and with (right) overlap filtering in Tokyo metropolitan area. We plot red circles of radius 1.32km (132 pixels) for better visualization.

we employ a grid search strategy to perform efficient overlap filtering. Instead of calculating the distance to all previous patches, we distribute the patch center coordinates into 360x180 geographical longitude-latitude, one-by-one-degree grids. For each new patch, we convert the center coordinates into integer grid coordinates. Subsequently, we search for existing patches within this grid cell and exclusively calculate distances to those local patches. Assuming potential overlap of sampled patches from distinct grid cells is statistically negligible, we significantly reduce computing time compared to a global overlap search. Indeed, for SSL4EO-S12 we record an overlap for approx. 3% tiles of densely populated Tokyo, 1.5% in Chicago, and below 1% for locations such as Bejing, Munich, Kampala, and Brasilia.

B. Data characteristics & volume

The presented SSL4EO-S12 dataset contains 251,079 globally distributed Sentinel-1 dual-pol SAR, Sentinel-2 top-ofatmosphere multispectral, and Sentinel-2 surface reflectance multispectral triplets over four seasonal timestamps. As of summer 2022, SSL4EO-S12 constitutes the biggest geospatial-temporal, multimodal dataset in terms of medium-resolution PolSAR and multi-spectral imagery serving more than 3 million images. The total data volume equates to an uncompressed size of $251,079 \times 4 \times [2 \cdot 4B + (13 + 12) \cdot 2B] \times 264^2 \approx 3.7TB$.

Figure 2 depicts the geospatial distribution of the SSL4EO-S12 dataset, highlighting the dense coverage across the globe. Figure 3 depicts the effect of overlap filtering around Tokyo area.

IV. EXPERIMENTAL SETUP

We evaluate SSL4EO-S12 dataset by self-supervised pretraining and transfer learning on RS downstream tasks. Specific implementation details are provided in the appendix.

A. Self-supervised pre-training

We perform pre-training using four representative SSL methods: *MoCo-v2/v3* [15, 16], *DINO* [7], *MAE* [8], and *data2vec* [9]. We pre-train ResNet [17] backbones with MoCo(-v2) and DINO, and Vision Transformer (ViT) [18] backbones for all four SSL methods listed above. Unless explicitly noted, Sentinel-2 L1C is used for pre-training. To utilize multi-temporal information, we use RandomSeasonContrast as a data augmentation strategy, i.e., for MoCo and DINO, the input views are randomly picked from two seasons. For MAE and data2vec, one random season is assigned for each patch.

Pre-training one ResNet/ViT model for 100 epochs takes 7–25 hours on 4 NVIDIA A100 GPUs, as shown in Table II.

 TABLE II

 100 epoch training time of the studied SSL methods.

	MoCo	DINO	MAE	data2vec
ResNet50	18h	25h	-	-
ViT-S/16	24h	25h	7h	14h

B. Transfer learning

The pre-trained models are transferred to various down-stream tasks. For

- *scene classification*, we evaluate EuroSAT [19] (singlelabel land cover classification), BigEarthNet [13] (multilabel land cover classification), and So2Sat-LCZ42 [20] (local climate zone classification, culture-10 version).
- *semantic segmentation*, we include DFC2020 [21] (land cover segmentation) and OSCD [22] (change detection).

We perform commonly used linear probing (freezing the pretrained encoder) and fine-tuning for the downstream tasks. The results are reported in percentage scores.

V. BENCHMARK RESULTS

A. Classification

1) Comparison of SSL methods: We first benchmark different SSL methods through linear probing on EuroSAT, BigEarthNet, and So2Sat-LCZ42. As detailed in Table III, all methods outperform random initialization (rand.init.) by a substantial margin. As expected, linear probing on BigEarth-Net with all labels performs worse than fully *supervised* training. Promisingly, the gap stays below 5%. On small datasets like BigEarthNet with 10% labels or EuroSAT, linear probing provides results comparable to supervised training within approx. $\pm 1\%$. The trends are slightly different for So2Sat-LCZ42, where the training and testing sets are built upon different cities with a challenging geographical split. Because of this significant domain shift, adding labeled training data does not necessarily improve the testing performance. In fact, fitting the training data distribution does not guarantee out-ofdistribution generalization. Nevertheless, the best pre-trained models with linear probing beat the supervised baseline by at least 1% up to about 4%.

Furthermore, we benchmark fine-tuning results in Table IV. All self-supervised methods outperform supervised learning

TABLE III

LINEAR PROBING RESULTS FOR EUROSAT, BIGEARTHNET (BE) AND SO2SAT-LCZ42 (10% AND 100% LABELS). WE REPORT OVERALL ACCURACY FOR EUROSAT AND SO2SAT-LCZ42, AND MEAN AVERAGE PRECISION (MICRO) FOR BIGEARTHNET. TWO BACKBONE NETWORKS GET TRAINED: RESNET-50 (RN50) AND A SMALL(-S) EMBEDDING DIMENSION VISION TRANSFORMER(VIT) SUBDIVIDING INPUT PATCHES INTO 16x16 TILES(/16). BOLD VALUES INDICATE BEST PER-COLUMN PERFORMANCE.

downstream dataset	Eu	roSAT	BE	E-10%	BE	-100%	So2S	Sat-10%	So2S	at-100%
model \setminus backbone	RN50	ViT-S/16	RN50	ViT-S/16	RN50	ViT-S/16	RN50	ViT-S/16	RN50	ViT-S/16
rand.init.	82.0	81.3	61.0	60.0	60.0	60.0	48.8	49.3	49.0	50.2
supervised	98.0	96.7	83.4	81.3	88.7	87.4	57.5	59.7	57.5	59.3
MoCo	- <u>98.</u> 0 -	97.7	-82.1	82.3	84.2	83.1	61.3	59.6	61.8	62.2
DINO	97.2	97.7	82.0	81.7	83.9	83.4	55.5	60.9	57.0	62.5
MAE	-	94.1	-	77.5	-	78.2		59.5	-	60.0
data2vec	-	96.9	-	77.3	-	79.4	-	58.2	-	59.7
TABLE IV										

FINE-TUNING RESULTS FOR EUROSAT, BIGEARTHNET, AND SO2SAT-LCZ42. ALL BEAT supervised training, cf. Table III.

downstream dataset	Eu	roSAT	BI	E-10%	BE	-100%	So28	Sat-10%	So2S	at-100%
model \setminus backbone	RN50	ViT-S/16								
MoCo	99.1	98.6	86.2	86.1	91.8	89.9	60.4	61.2	60.9	61.6
DINO	99.1	99.0	87.1	86.9	90.7	90.5	63.2	61.5	63.6	62.2
MAE		98.7	-	84.8	-	88.9	-	60.8	-	63.9
data2vec	-	99.1	-	85.6	-	90.3	-	63.2	-	64.8

with a margin from 1% to 6%. Top SSL-models score 99.1% on EuroSAT (MoCo/DINO) and over 90% on BigEarthNet (MoCo/DINO). Comparing linear probing and fine-tuning results, one interesting phenomenon shows up: in linear probing contrastive methods (MoCo and DINO) consistently score better than their image-masking (MAE and data2vec) counterparts.

2) Comparison of pre-training datasets: To compare SSL4EO-S12 with other RS pre-training datasets, we report corresponding linear probing results pre-trained with MoCo-v2 (ResNet50 backbone) in Table V. Similar to SSL4EO-S12, RandomSeasonContrast is used to pick one timestamp image for each geospatial patch in SeCo dataset. In the first set of comparison, we use RGB bands only. SSL4EO-S12 significantly outperforms ImageNet by about 10%, SeCo by about 6%, and SEN12MS by 1.7% to 3.5%.

 TABLE V

 Comparison of different pre-training datasets under linear

 probing evaluation. *italic* means cited from the literature.

dataset	EuroSAT	BE-10%	BE-100%
ImageNet (RGB) [4]	86.4	70.5	71.8
SeCo (RGB) [4]	89.5	74.5	76.3
SEN12MS (RGB)	94.9	76.6	79.6
SSL4EO-S12 (RGB)	96.6	80.1	82.3
SeCo ² (all bands)	89.2	73.7	76.6
SEN12MS (all bands)	95.5	79.6	82.1
BigEarthNet (all bands)	94.4	80.6	83.9
SSL4EO-S12 (all bands)	98.0	82.1	84.2

In a second set of experiments we evaluate all multispectral bands. Results indicate consistent performance gain as in RGB setting comparing SSL4EO-S12 with SEN12MS and SeCo. In addition, pre-training on SSL4EO-S12 outperforms BigEarth-Net on itself and EuroSAT (both are EU only). This proves SSL4EO-S12's benefits to improve model transferability by learning valuable knowledge from a larger scale and wider geographical coverage. 3) Comparison of different amounts of labels: Figure 4 visualizes performance results of transfer learning on BigEarth-Net with a varying fraction of labeled samples. Compared to the supervised baseline, self-supervised pre-training on SSL4EO-S12 provides significant benefits when the amount of labeled samples is limited. In fact, fine-tuning on 10% of the labels outperforms 50%-labels supervised training; and with ViT-S/16, fine-tuning on 50% of the labels outperforms 100%labels supervised training.

B. Segmentation

1) Land cover segmentation: We use DFC2020 [21] dataset to evaluate land cover semantic segmentation. We pre-train ResNet50 with MoCo-v2 on SSL4EO-S12 L1C products, and fine-tune a DeepLabv3+ [23] for segmentation. Table VI lists results with notable improvements when compared to SeCo pre-training. However, SSL4EO-S12 performs worse than SEN12MS in average accuracy (AA) and mean intersection over union (mIoU). This can be expected, since DFC2020 was built with direct reference to SEN12MS and they have similar data distribution. Nevertheless, the results are still comparable, proving again the transferability of the proposed dataset.

TABLE VI DFC2020 land cover segmentation results.

dataset	OA	AA	mIoU
rand.init.	81.97	56.46	42.11
SeCo	87.31	57.05	49.68
SEN12MS	88.64	67.69	54.83
SSL4EO-S12	89.58	64.01	54.68

2) Change detection: We evaluate the pre-trained models for change detection on the OSCD [22] dataset. We pretrain ResNet50 with MoCo-v2 on SSL4EO-S12 L1C products, freeze the backbone, and fine-tune a U-Net [24] for segmentation. The differences in feature maps between two timestamps are input to the network. As Table VII indicates, pre-training on SSL4EO-S12 yields superior performance in recall and F1-score when referenced to SeCo and SEN12MS. While SSL4EO-S12 performs worse in precision, this is due

²The available SeCo data at https://github.com/ServiceNow/ seasonal-contrast has only about 160k geographical patches (instead of 200k in the paper), which may affect our reproduced performance.



Fig. 4. BigEarthNet (BE) performance depending on amount of labels available to train downstream task. We report linear probing and fine-tuning results with ResNet50 and ViT-S/16 encoders pre-trained using MoCo-v2.

TABLE VII OSCD CHANGE DETECTION RESULTS.

dataset	precision	recall	F1
rand.init.	72.31	13.75	23.10
SeCo	74.85	17.47	28.33
SEN12MS	74.67	19.26	30.62
SSL4EO-S12	70.23	23.38	35.08

to the significant class unbalance that predicting all pixels as unchanged would result in a good precision score.

VI. ADDITIONAL STUDIES

We complete our benchmark by reporting a set of additional results to document key characteristics of the SSL4EO-S12 dataset, namely: multi-temporal, multimodal, multi-product-level, and data scale. For all studies, we pre-train ResNet50 with MoCo-v2 as a common setting.

A. Ablation studies

1) Benefits of multimodality: While Section V employs only optical data for fair comparison to existing literature, we highlight the benefits of multimodal pre-training in this section. We integrate SAR data by early fusion, and use RandomSensorDrop [12] as an additional data augmentation strategy. During training, the model gets fed random combinations of SAR/optical patches, thus learning both inner- and inter-modality representations. Then, the pre-trained model gets transferred to different scenarios where either both modalities or a single one is available. We compare multimodal pretraining (MM) to uni-modal pre-training (S1/2) on BigEarth-Net. Table VIII presents results with notable improvement of 1%-3% for 100% and 1% label splits. While singlemodality pre-training already works well for both Sentinel-2 and Sentinel-1 data, pre-training exploiting both modalities further improves performance.

TABLE VIII LINEAR PROBING RESULTS OF MULTIMODAL SSL. MOCO-S1/2 REPRESENTS PRE-TRAINING WITH ONE SINGLE MODALITY, AND MOCO-MM REPRESENTS PRE-TRAINING WITH BOTH MODALITIES.

downstr./		BE-19	6		BE-100	%
model	S 1	S2	S1+S2	S1	S2	S1+S2
MoCo-S1/2	71.1	75.9	_	75.9	84.2	_
MoCo-MM	73.3	76.7	76.8	79.5	85.1	85.2
supervised	66.7	75.7	76.4	77.2	88.7	88.9

2) Ablation of seasonal information: We evaluate the effectiveness of multi-temporal information by replacing seasonal *augment*ation (cf. Section IV) by *random* season: the same randomly selected season for the two positive views; and *fixed* season: the same season for each patch during training. We pre-train on a 50k subset of SSL4EO-S12, and evaluate on BigEarthNet-10% and EuroSAT. Table IX clearly proves the benefits of seasonal augmentation.

 TABLE IX

 Linear probing results of multi-temporal ablation study.

downstr./season	BE-10%	EuroSAT
fixed	75.1	93.1
random	76.7	94.0
augment	77.6	96.2

3) Atmospheric correction as data augmentation: The motivation to include Sentinel-2 L1C and L2A products in SSL4EO-S12 is to match corresponding downstream tasks. However, these product levels with or without atmospheric correction can also be considered natural data augmentation for SSL. Accordingly, we conduct an ablation study on a 50k SSL4EO-S12 subset utilizing Sentinel-2 L1C, L2A or both (L1C+L2A). Table X summarizes our findings: 1) models pretrained on the same product level as the downstream task have a slight edge ($\sim 1\%$) over models trained on the other product level, and 2) pre-training on both modalities generates a notable improvement of up to 4% compared to pre-training on single modality.

TABLE X LINEAR PROBING RESULTS OF DIFFERENT PRODUCT LEVELS OF Sentinel-2.

product	BE-10% (L2A)	EuroSAT (L1C)
L1C	74.0	93.1
L2A	75.1	92.0
L1C+L2A	78.0	93.8

4) Impact of pre-training scale: An aspect relevant to largescale data mining in Earth observation is scaling of results with training data volume: why don't we add more images to SSL4EO-S12? One reason concerns computational costs. We believe the current dataset (1M patches for each Sentinel product) is comparable to the scale of ImageNet, and can serve as a good baseline in remote sensing for further development. Moreover, as observed by [25], saturating downstream performance kicks in beyond 500k pre-training images on ImageNet, with 250k images yielding acceptable results with as little as 1-2% accuracy loss. We observe such a trend in our dataset, too. As demonstrated by Table XI, we pre-train on various amounts of data to report linear probing results for BigEarthNet-10%. While 50% (500K) or less pre-training data yields significant performance drops, there's little diminishing gaps from 75% (750K) on. Note this saturation effect depends also on the model size.

TABLE XI LINEAR PROBING RESULTS ON BIGEARTHNET-10% FOR VARIOUS SENTINEL-2 L1C PRE-TRAINING DATA SIZES.

data size	100K	250K	500K	750K	1M
accuracy (%)	64	73	78	81	82

B. Representation visualization

We qualitatively evaluate the data representations learned from self-supervised pre-training by visualizing the latent distributions with t-SNE (Figure 5). We pre-train a ResNet50 with MoCo-v2 on SSL4EO-S12, and transfer the frozen encoder to EuroSAT to calculate one 128d representation vector for each image. We then visualize all the vectors with t-SNE, and compare the distribution with a randomly initialized encoder.



Fig. 5. t-SNE visualization of EuroSAT image representations. One color represents one class. Left: random-encoded features; right: SSL-encoded features. SSL-encoded features are well clustered even without label information.

VII. CONCLUSION

In this work, we present SSL4EO-S12—a large-scale multimodal, multi-temporal unlabeled dataset for self-supervised learning (SSL) in Earth observation. An extensive benchmark on various SSL methods and remote sensing applications proves the promising benefits of the proposed dataset.

SSL4EO-S12 has some limitations: 1) there's little coverage of polar regions; 2) geographical bias exists due to cloud filtering; 3) it is not strictly free of geospatial overlap; 4) mediumresolution radar and multispectral images are a limited subset of Earth observation data. Despite these, we believe SSL4EO-S12 renders a valuable basis to advance self-supervised pretraining and large-scale data mining in remote sensing.

ACKNOWLEDGMENTS

This work is jointly supported by the Helmholtz Association through the Framework of Helmholtz AI (grant number: ZT-I-PF-5-01) - Local Unit "Munich Unit @Aeronautics, Space and Transport (MASTr)" and Helmholtz Excellent Professorship "Data Science in Earth Observation - Big Data Fusion for Urban Research"(grant number: W2-W3-100), by the German Federal Ministry of Education and Research (BMBF) in the framework of the international future AI lab "AI4EO – Artificial Intelligence for Earth Observation: Reasoning, Uncertainties, Ethics and Beyond" (grant number: 01DD20001) and by German Federal Ministry for Economic Affairs and Climate Action in the framework of the "national center of excellence ML4Earth" (grant number: 50EE2201C). The computing resources were supported by the Helmholtz Association's Initiative and Networking Fund on the HAICORE@FZJ partition.

REFERENCES

- [1] Yi Wang et al. "Self-Supervised Learning in Remote Sensing: A Review". In: *IEEE Geoscience and Remote Sensing Magazine* (2022).
- [2] Jia Deng et al. "Imagenet: A large-scale hierarchical image database". In: *IEEE conference on computer vision and pattern recognition*. 2009.
- [3] Michael Schmitt et al. "SEN12MS a curated dataset of georeferenced multi-spectral Sentinel-1/2 imagery for deep learning and data fusion". In: *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences*. 2019.
- [4] Oscar Mañas et al. "Seasonal contrast: Unsupervised pretraining from uncurated remote sensing data". In: Proceedings of the IEEE/CVF International Conference on Computer Vision. 2021.
- [5] Zhitong Xiong et al. "EarthNets: Empowering AI in Earth Observation". In: *arXiv:2210.04936* (2022).
- [6] Kaiming He et al. "Momentum contrast for unsupervised visual representation learning". In: Proceedings of the IEEE/CVF conference on computer vision and pattern recognition. 2020.
- [7] Mathilde Caron et al. "Emerging properties in self-supervised vision transformers". In: Proceedings of the IEEE/CVF International Conference on Computer Vision. 2021.
- [8] Kaiming He et al. "Masked autoencoders are scalable vision learners". In: arXiv:2111.06377 (2021).
- [9] Alexei Baevski et al. "Data2vec: A general framework for self-supervised learning in speech, vision and language". In: *arXiv:2202.03555* (2022).
- [10] Mathilde Caron et al. "Unsupervised learning of visual features by contrasting cluster assignments". In: Advances in Neural Information Processing Systems (2020).
- [11] Maxim Neumann et al. "Training general representations for remote sensing using in-domain knowledge". In: *IEEE International Geoscience and Remote Sensing Symposium*. 2020.
- [12] Yi Wang, Conrad M Albrecht, and Xiao Xiang Zhu. "Selfsupervised Vision Transformers for Joint SAR-optical Representation Learning". In: arXiv:2204.05381 (2022).
- [13] Gencer Sumbul et al. "Bigearthnet: A large-scale benchmark archive for remote sensing image understanding". In: *IEEE International Geoscience and Remote Sensing Symposium*. 2019.
- [14] Noel Gorelick et al. "Google Earth Engine: Planetary-scale geospatial analysis for everyone". In: *Remote sensing of Environment* (2017).
- [15] Xinlei Chen et al. "Improved baselines with momentum contrastive learning". In: *arXiv:2003.04297* (2020).
- [16] Xinlei Chen, Saining Xie, and Kaiming He. "An empirical study of training self-supervised vision transformers". In: *Proceedings of the IEEE/CVF International Conference on Computer Vision*. 2021.
- [17] Kaiming He et al. "Deep residual learning for image recognition". In: *Proceedings of the IEEE conference on computer vision and pattern recognition.* 2016.
- [18] Alexey Dosovitskiy et al. "An image is worth 16x16 words: Transformers for image recognition at scale". In: arXiv:2010.11929 (2020).

- [19] Patrick Helber et al. "Eurosat: A novel dataset and deep learning benchmark for land use and land cover classification". In: *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* (2019).
- [20] Xiao Xiang Zhu et al. "So2Sat LCZ42: A benchmark data set for the classification of global local climate zones". In: *IEEE Geoscience and Remote Sensing Magazine* (2020).
- [21] Michael Schmitt et al. IEEE GRSS Data Fusion Contest. 2020.
- [22] Rodrigo Caye Daudt et al. "Urban change detection for multispectral earth observation using convolutional neural networks". In: *IEEE International Geoscience and Remote Sensing Symposium*. 2018.
- [23] Liang-Chieh Chen et al. "Encoder-decoder with atrous separable convolution for semantic image segmentation". In: *Proceedings of the European conference on computer vision* (ECCV). 2018, pp. 801–818.
- [24] Olaf Ronneberger, Philipp Fischer, and Thomas Brox. "U-net: Convolutional networks for biomedical image segmentation". In: International Conference on Medical image computing and computer-assisted intervention. 2015.
- [25] Elijah Cole et al. "When does contrastive visual representation learning work?" In: *Proceedings of the IEEE/CVF Conference* on Computer Vision and Pattern Recognition. 2022.

APPENDIX-1: ADDITIONAL DATASET INFORMATION

A. Sentinel-1/2

The proposed SSL4EO-S12 dataset exploits freely available SAR/optical satellite images from European Space Agency's Sentinel mission (under the CC-BY license).

The Sentinel-1 mission [26] consists of two polar-orbiting satellites, equipped with C-band SAR sensors, which enables them to acquire imagery regardless of the weather. For the Sentinel-1 images in the SSL4EO-S12 dataset, ground-range-detected (GRD) products with both VH and VV polarization acquired in the interferometric wide swath (IW) mode were used. These images contain the σ^0 backscatter coefficient in dB scale. The image resolution is 10m.

The Sentinel-2 mission [27] comprises two polar-orbiting satellites in the same orbit, equipped with multi-spectral imaging sensors. For Sentinel-2 images in the SSL4EO-S12 dataset, both level-1C top-of-atmosphere reflectance (13 bands) and level-2A atmospherically corrected surface reflectance (12 bands) were included. The image resolution ranges between 10m (visible and NIR), 20m (red edge and SWIR) and 60m (aerosols).

B. Dataset statistics

Table XII and XIII present the mean and standard deviation of each band for each product of the proposed SSL4EO-S12 dataset.

TABLE XII	
STATISTICS OF SENTINEL-1 IMAGES IN TH	HE SSL4EO-S12 DATASET.

	VV	VH
mean	-12.59	-20.26
std	5.26	5.91

 TABLE XIII

 Statistics of Sentinel-2 images in the SSL4EO-S12 dataset.

		B1	B2 (B)	B3 (G)	B4 (R)	B5	B6	B7	B8	B8A	B9	B10	B11	B12
-2-	mean	1612.9	1397.6	1322.3	1373.1	1561.0	2108.4	2390.7	2318.7	2581.0	837.7	22.0	2195.2	1537.4
s2c	std	791.0	854.3	878.7	1144.9	1127.5	1164.2	1276.0	1249.5	1345.9	577.5	47.5	1340.0	1142.9
- ? -	mean	756.4	889.6	1151.7	1307.6	1637.6	2212.6	2442.0	2538.9	2602.9	2666.8	-	2388.8	1821.5
82a	std	1111.4	1159.1	1188.1	1375.2	1376.6	1358.6	1418.4	1476.4	1439.9	1582.1	-	1460.7	1352.2

C. Data storage

The SSL4EO-S12 dataset is stored in GeoTiff format for each band of each patch. The file structure is shown in Figure 6, where s1/s2a/s2c represents Sentinel-1 / Sentinel-2 level-2A / Sentinel-2 level-1C, and t1 - t4 represent 4 seasons. Raw files (extracted GeoTiff) occupy about 500GB/800GB/800GB disk storage for S1/S2A/S2C, and compressed tar.gz files occupy about 450GB/500GB/500GB correspondingly. If converting to uint8 and encoding with jpeg, a lossy dataset occupies less than 50 GB for each product. We later show this won't affect much the downstream performance.

D. Metadata

Each patch comes with a metadata file that collects the image properties of this patch. See Table XX and XXI for details.

E. Example visualization

Figure 7 visualizes an example geospatial tile of SSL4EO-S12.



Fig. 6. SSL4EO-S12 file structure.



Fig. 7. Sample visualization of one tile from SSL4EO-S12 dataset. The rows top-down present grayscale and false-color imagery based on the Sentinel-1 GRD product, Sentinel-2 level-1C, and Sentinel-2 level-2A multispectral data with corresponding columns representing the four seasons spring, summer, fall, and winter from left to right.

APPENDIX-2: IMPLEMENTATION DETAILS

A. Pre-training

We use Sentinel-2 level-1C images for the main pre-training experiments, which are pre-processed by converting to uint8 for efficiency (divided by 10000 and multiplied by 255, see Section VII-B5). We use 4 NVIDIA A100 GPUs with a total batch size of 256 for all the pre-training experiments. For the main experiments, we pre-train ResNet50 or Vit-S/16 for 100 epochs. Training time varies between different methods from 7 (MAE) to 25 (DINO) hours. The total experiments (including parameter tuning) take about 70k core hours (1400 GPU hours).

1) MoCo: We pre-train the MoCo-v2/v3 models using their default settings following the publicly available repository (https://github.com/facebookresearch/moco and https://github.com/facebookresearch/moco-v3). We use RandomResizedCrop, RandomBrightness/Contrast (to have a partial color jittering for multiple bands), RandomGrayscale, RandomGaussianBlur, RandomHorizontalFlip and RandomSeasonContrast as data augmentations. For MoCo-v2 (ResNet50), we use SGD optimizer and cosine learning rate schedule with a learning rate 0.03. For MoCo-v3 (ViT-S/16), we use AdamW optimizer and cosine schedule with a learning rate 1.5e-4.

2) *DINO:* We pre-train the DINO models using its default settings following the publicly available repository (https: //github.com/facebookresearch/dino). The data augmentations include those of MoCo, as well as additional Multi-Crop and Solarization. For ResNet50, we use SGD optimizer and cosine learning rate schedule with a learning rate 0.03. For ViT-S/16, we use AdamW optimizer and cosine learning rate schedule with a learning rate 1.5e-4.

3) MAE: We pre-train the MAE models using its default settings following the publicly available repository (https: //github.com/facebookresearch/mae). The mask ratio is set to 0.7. The data augmentations include RandomResizedCrop, RandomHorizontalFlip and RandomSeason. We use AdamW optimizer and cosine learning rate schedule with a learning rate 1.5e-4.

4) data2vec: We pre-train the data2vec models using its default settings following the publicly available repository (https://github.com/facebookresearch/fairseq/tree/main/examples/data2vec). The data augmentations include Resize/CenterCrop, RandomHorizontalFlip and RandomSeason. We use AdamW optimizer and cosine learning rate schedule with a learning rate 1e-3.

B. Downstream tasks

Below are additional implementation details for the downstream tasks.

1) EuroSAT: We split EuroSAT into 21600 training and 5400 testing images for evaluation. The data augmentations are RandomResizedCrop/RandomHorizontalFlip for training, and Resize/CenterCrop for testing. We resize the images to 224x224 for better performance (see Section VII-B5). The batch size is 256. We use CrossEntropyLoss and SGD optimizer with step decay learning rate (divided by 10 at epoch 60 and 80) for 100 epochs. We use simple grid search strategy to find suitable learning rates for linear probing and fine-tuning.

2) *BigEarthNet:* We use 311667 training and 103944 testing images from BigEarthNet for evaluation. Different settings of the amount of labels affect only the training split. The data augmentations are RandomResizedCrop/RandomHorizontalFlip for training, and Resize/CenterCrop for testing. We use a cropping scale of 0.8 to avoid strong occlusions (BigEarthNet is a multi-label dataset). We resize the images to 224x224 for better performance. The batch size is 256. We use MultiLabelSoftMarginLoss and SGD optimizer with step decay learning rate (divided by 10 at epochs 60 and 80) for 100 epochs. We use a simple grid search strategy to find suitable learning rates for linear probing and fine-tuning.

3) So2Sat-LCZ42: We use 352366 training and 24119 testing Sentinel-2 images from So2Sat-LCZ42 for evaluation. Different settings of the amount of labels affect only the training split. We use the *culture-10* version of So2Sat-LCZ42: the training data and the testing data are from different cities. The data augmentations are RandomResizedCrop/RandomHorizontalFlip for training, and Resize/CenterCrop for testing. We resize the images to 224x224 for better performance. The batch size is 256. We use CrossEntropyLoss and SGD optimizer with step decay learning rate (divided by 10 at epochs 60 and 80) for 100 epochs. We use a simple grid search strategy to find suitable learning rates for linear probing and fine-tuning.

4) *DFC2020:* We use 5128 training and 986 testing Sentinel-2 images from DFC2020 dataset for evaluation. The batch size is set to 8 and we train the models for 50 epochs. We use CrossEntropyLoss and SGD optimizer with momentum 0.9 and weight decay 5e-4. The initial learning rate is 1e-3, which is decayed by a factor of 0.9 in every epoch until 1e-4.

5) OSCD: This dataset is composed of 24 pairs of multispectral images from Sentinel-2 in total. Following [22], we use 14 of them for training, and the rest for testing. In fine-tuning stage, we adopt settings similar to those in [4]. That is, the original images are split into non-overlapping patches of 96×96 pixels as inputs, which leads to 827 and 285 patches for training and testing, respectively. The batch size is set to 32, and we in total train 50 epochs. We use Adam optimizer with a weight decay of 1e-4. The initial learning rate is 1e-3, and decreases exponentially with a multiplicative factor of 0.95 for every epoch. The resulting models are evaluated on the whole test set for overall precision, recall and F1 score (with a default threshold 0.5).

APPENDIX-3: ADDITIONAL EXPERIMENTAL RESULTS

A. Effect of data pre-processing

We show the influence of data pre-processing (for pre-training) in Table XIV, where int16 means 16 bits raw input, uint8 means compressed 8 bits input (divided by 10000 and multiplied by 255), uint8-n means normalization by mean and standard deviation, and L2A/L1C means with/without atmospheric correction. The results show similar performance between 16 bit and 8 bit, supporting compressed input as it saves a lot of storage space and computing time. The results also show comparable performance for L1C and L2A, as well as for the use of normalization. Therefore, in our main experiments, we use level-1C, uint8, unnormalized data for pre-training.

 TABLE XIV

 Effects of pre-processing for pretraining. We pre-train a ResNet50 with MoCo-v2 on a 50k subset for efficiency and report linear probing results.

	BE-10% (L2A)			Eu	roSAT (1	L1C)
	int16	uint8	uint8-n	int16	uint8	uint8-n
L2A	73.9	73.9	75.6	85.2	86.6	-
L1C	73.8	74	-	86.2	87.7	85.1

B. Effect of input image size

We analyze the impact of image resolution on pre-training and transfer learning in Table XV. We clearly observe the advantage of upsampling the input image size. Therefore, in our main experiments, we upscale the downstream input images to 224x224 for better performance.

TABLE XV EFFECTS OF INPUT SIZE OF DOWNSTREAM TASKS. WE PRE-TRAIN RESNET50 WITH MOCO-V2 ON THE 50K SUBSET FOR EFFICIENCY AND REPORT LINEAR PROBING RESULTS.

		BE-10%)	EuroSAT			
	224	112	56	224	112	56	
pretrain 224	75.1	71.1	-	93.1	89.4	89.4	
pretrain 112	75.9	73.9	-	92.7	91.1	86.6	
pretrain 56	73.1	70.5	70.5	89.7	89	86.4	

C. Effect of MAE masking ratio

Table XVI shows the influence of masking ratios in MAE during pre-training. We find 70% to be the best masking ratio, which is similar to natural images as reported in MAE paper, where 75% is the best. It is also promising to see that the model still learns good representations even with 90% pixels masked.

 TABLE XVI

 Effects of different masking ratios. We pre-train ViT-S/16 with MAE on the 50k subset for efficiency and report linear probing results on BigEarthNet-10%.

Mask ratio	90%	80%	75%	70%	60%	50%
BE-10%	72	73.5	73.8	74	73.6	72.9

D. Effect of different pre-training protocols

1) Different ImageNet pre-training protocols: Table XVII shows a comparison of different ImageNet pre-training protocols. We pre-train ResNet50 with MoCo-v2 for self-supervised pre-training, and report fine-tuning results on BigEarthNet. The table shows that ImageNet pre-training provides good representations that can be generalized well to remote sensing images with RGB but not all bands. It can also be seen that when using RGB, self-supervised pre-training on ImageNet can further improve the downstream performance in remote sensing compared to supervised pre-training.

2) Supervised pre-training on RS datasets: Table XVIII shows a comparison of supervised and unsupervised pre-training on remote sensing datasets. We do supervised pre-training on BigEarthNet and self-supervised pre-training (MoCo-v2, ResNet50) on both BigEarthNet and SSL4EO-S12. We evaluate the pre-trained models on EuroSAT. The results show that self-supervised pre-training outperforms supervised pre-training on remote sensing data.

 TABLE XVII

 Summary of different ImageNet pre-training protocols.

downstr./	All opti	cal bands	RGB		
pre-train	BE-10%	BE-100%	BE-10%	BE-100%	
Supervised	83.4	88.7	69.5	79.0	
ImageNet (Sup)	82.5	89.5	80.0	86.7	
ImageNet (SSL)	82.5	89.4	81.9	90.8	
SSL4EO-S12 (SSL)	86.2	91.8	82.7	90.9	

TABLE XVIII

A COMPARISON OF DIFFERENT PRE-TRAINING PROTOCOLS ON REMOTE SENSING DATASETS AND EVALUATED ON EUROSAT.

downstr./	Linear	probing	Fine-	tuning
pre-train	10%	100%	10%	100%
BigEarthNet (Sup)	80.3	89.3	94.2	98.7
BigEarthNet (SSL)	90.7	94.4	96.3	98.9
SSL4EO-S12 (SSL)	92.7	98.0	96.9	99.1

E. Additional dataset comparison results

Table XIX reports fine-tuning results pre-trained on different datasets in complement to the main paper. A difference most notable is ImageNet's catch-up in performance compared to the other geospatial pre-training datasets. However, a $\sim 5\%$ margin when compared to SSL4EO-S12 persists. The observed is a characteristic feature of the data *domain gap*: while pre-training on ImageNet learns good representations, the weights' distribution is shifted towards natural images, which can be further adjusted to remote sensing data with fine-tuning. We note that fine-tuning is more computationally expensive compared to linear probing, and Table XIX demonstrate: pre-training on SSL4EO-S12 outperforms all other datasets for downstream classification.

 TABLE XIX

 COMPARISON OF DIFFERENT PRE-TRAINING DATASETS UNDER FINE-TUNING EVALUATION.

dataset	EuroSAT	BE-10%	BE-100%	So2Sat-10%	So2Sat-100%
ImageNet (RGB)	96.5	80.0	86.7	-	-
SeCo (RGB)	-	80.2	86.1	-	-
SSL4EO-S12 (RGB)	98.0	82.7	90.9	-	-
BigEarthNet (MS)	98.9	85.5	89.3	53.0	53.0
SSL4EO-S12 (MS)	99.1	86.2	91.8	60.4	60.9

APPENDIX-4: METADATA

TABLE XX

METADATA SENTINEL-1 GRD[14].

Name	Туре	Description
GRD_Post_Processing_facility_country	STRING	Name of the country where the facility is located. This element is configurable within the IPF.
GRD_Post_Processing_facility_name	STRING	Name of the facility where the processing step was performed. This element is configurable within the
CRD Date Description for illiter service time	CTDDIC	IPF.
GRD_Post_Processing_facility_organisation	STRING	Name of the organisation responsible for the facility. This element is configurable within the IPF.
GRD_Post_Processing_facility_site	STRING	Geographical location of the facility. This element is configurable within the IPF.
GRD_Post_Processing_software_name	STRING	Name of the software.
GRD_Post_Processing_software_version	STRING	Software version identification.
GRD_Post_Processing_start	DOUBLE	Processing start time.
GRD_Post_Processing_stop	DOUBLE	Processing stop time.
SLC_Processing_facility_country	STRING	Name of the country where the facility is located. This element is configurable within the IPF.
SLC_Processing_facility_name	STRING	Name of the facility where the processing step was performed. This element is configurable within the IPF.
SLC_Processing_facility_organisation	STRING	Name of the organisation responsible for the facility. This element is configurable within the IPF.
SLC_Processing_facility_site	STRING	Geographical location of the facility. This element is configurable within the IPF.
SLC_Processing_software_name	STRING	Name of the software.
SLC Processing software version	STRING	Software version identification.
SLC Processing start	DOUBLE	Processing start time.
SLC Processing stop	DOUBLE	Processing stop time.
S1TBX Calibration Operator version	STRING	Sentinel-1 Toolbox calibration tool version.
S1TBX SAR Processing version	STRING	Sentinel-1 Toolbox SAB processing tool version.
SNAP Graph Processing Framework GPE version	STRING	Sentinel Application Platform (SNAP) version
startTimeANX	DOUBLE	Sensing start time of the input data relative to the ascending node crossing. This is a count of the time
start finite fi tx	DOUDLL	elansed since the orbit accending node crossing [ms]
stonTimeANY	DOUBLE	Sensing stop time of the input data relative to the according node crossing. This is a count of the time
stopTimeAtVX	DOUDLL	sensing stop time of the input data relative to the ascending hole crossing. This is a count of the time
needaldantifiar	STRING	Lapsed since the other according house crossing [mis].
lissueidentinei	STRING	Information (MDC SL) available hore
familyNama	STRINC	The full mission name E.e. "ENTIMEL 1"
	STRING	The full mission name, E.g. SENTINEL-1
platform_number	STRING	The appanument identifier of the platform within the mission.
instrument	STRING	Information related to the instrument on the platform to which acquired the data.
instrumentMode	STRING	IW (Interferometric Wide Swath), EW (Extra Wide Swath) or SM (Strip Map)
instrumentSwath	STRING	List of the swaths contained within a product. Most products will contain only one swath, except for
		TOPS SLC products which include 3 or 5 swaths.
orbitNumber_start	DOUBLE	Absolute orbit number of the oldest line within the image data.
orbitNumber_stop	DOUBLE	Absolute orbit number of the most recent line within the image data.
relativeOrbitNumber_start	DOUBLE	Relative orbit number of the oldest line within the image data.
relativeOrbitNumber_stop	DOUBLE	Relative orbit number of the most recent line within the image data.
cycleNumber	DOUBLE	Absolute sequence number of the mission cycle to which the oldest image data applies.
phaseIdentifier	DOUBLE	Id of the mission phase to which the oldest image data applies.
orbitProperties_pass	STRING	Direction of the orbit ('ASCENDING' or 'DESCENDING') for the oldest image data in the product (the
		start of the product).
orbitProperties_ascendingNodeTime	DOUBLE	UTC time of the ascending node of the orbit. This element is present for all products except ASAR L2
		OCN products which are generated from an ASAR L1 input.
resolution	STRING	H for high or M for medium.
resolution meters	DOUBLE	Resolution in meters.
instrumentConfigurationID	DOUBLE	The instrument configuration ID (Radar database ID) for this data.
missionDataTakeID	DOUBLE	Unique ID of the datatake within the mission
transmitterReceiverPolarisation	DOUBLE	Transmit/Receive polarisation for the data. There is one element for each Tx/Rx combination: ["VV"]
datistitueriveeriveri olarisation	DOUDLL	This indicate the polarisation for the data. There is one element for each Twick combination. [$\uparrow \uparrow \downarrow$], ["HIP"] ["V(V", "VIP"] or ["HIP" HV"]
productClass	STRING	Output product class "A for Annotation or "S" for Standard
productClassDescription	STRING	Taytial description of the output product class
productComposition	STRING	The composition type of this product "Individual" "Slice" or "Assambled"
productCompOsition	STRING	The composition type of this product. Individual, Site, of Assembled.
product Type	STRING	The product type (concettion level) of this product.
product finefinessCategory	STRING	bescribes the required unmenness of the processing. One or: NK1-10m, NK1-1h, NK1-3h, Fast-24h, Off-
l' - De le dEle e	CTDINC	me, or reprocessing
sicerroductriag	STRING	rue if this is a since from a larger product or faise if this is a complete product.
segmentStartTime	DOUBLE	Sensing start time of the segment to which this slice belongs. This field is only present if sliceProductFlag
	DOUDI -	= true
sliceNumber	DOUBLE	Absolute slice number of this slice starting at 1. This field is only present if sliceProductFlag = true.
totalSlices	DOUBLE	Total number of slices in the complete data take. This field is only present if sliceProductFlag = true.

TABLE XXI

METADATA SENTINEL-2 L1C[14].

Name	Type	Description
AOT RETRIEVAL ACCURACY	DOUBLE	Accuracy of Aerosol Ontical thickness model
CLOUDY PIXEL PERCENTAGE	DOUBLE	Granule-specific cloudy pixel bercentage taken from the original metadata
CLOUD_COVERAGE_ASSESSMENT	DOUBLE	Cloudy pixel percentage for the whole archive that contains this granule. Taken from the original metadata
CLOUDY_SHADOW_PERCENTAGE	DOUBLE	Percentage of pixels classified as cloud shadow
DARK_FEATURES_PERCENTAGE	DOUBLE	Percentage of pixels classified as dark features or shadows
DATASTRIP_ID	STRING	Unique identifier of the datastrip Product Data Item (PDI)
DATATAKE_IDENTIFIEK	STRING	Uniquely identities a given Datatake. The ID contains the Sentiner-2 saterine, start date and time, absolute orbit number, and processing baseline.
DATATAKE TYPE	STRING	MSI operation mode
DEGRADED_MSI_DATA_PERCENTAGE	DOUBLE	Percentage of degraded MSI and ancillary data
FORMAT_CORRECTNESS	STRING	Synthesis of the On-Line Quality Control (OLQC) checks performed at granule (Product_Syntax) and datastrip (Product
	6770 D.10	Syntax and DS_Consistency) levels
GENERAL_QUALITY GENERATION TIME	STRING	Synthesis of the OLQC checks performed at the datastrip level (Relative_Orbit_Number)
GEOMETRIC QUALITY	STRING	Product generation time Synthesis of the OLOC checks performed at the datastrin level (Attitude Quality Indicator)
GRANULE ID	STRING	Unique identifier of the granule PDI (PDI ID)
HIGH_PROBA_CLOUDS_PERCENTAGE	DOUBLE	Percentage of pixels classified as high probability clouds
MEAN_INCIDENCE_AZIMUTH_ANGLE_B1	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B1 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B2	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B2 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B3 MEAN_INCIDENCE_AZIMUTH_ANGLE_B4	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B3 and for all detectors
MEAN INCIDENCE AZIMUTH ANGLE B5	DOUBLE	Mean value containing viewing incidence azimuti angle average for band B5 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B6	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B6 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B7	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B7 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B8	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B8 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B8A	DOUBLE DOUBLE	Mean value containing viewing incidence azimuth angle average for band B8a and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B9	DOUBLE	Mean value containing viewing incidence azimuti angle average for band B0 and for all detectors.
MEAN INCIDENCE AZIMUTH ANGLE B11	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B11 and for all detectors
MEAN_INCIDENCE_AZIMUTH_ANGLE_B12	DOUBLE	Mean value containing viewing incidence azimuth angle average for band B12 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B1	DOUBLE	Mean value containing viewing incidence zenith angle average for band B1 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B2	DOUBLE	Mean value containing viewing incidence zenith angle average for band B2 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B3 MEAN_INCIDENCE_ZENITH_ANGLE_B4	DOUBLE	Mean value containing viewing incidence zenith angle average for band B4 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B4	DOUBLE	Mean value containing viewing incidence zenith angle average for band B5 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B6	DOUBLE	Mean value containing viewing incidence zenith angle average for band B6 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B7	DOUBLE	Mean value containing viewing incidence zenith angle average for band B7 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B8	DOUBLE	Mean value containing viewing incidence zenith angle average for band B8 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B8A	DOUBLE	Mean value containing viewing incidence zenith angle average for band B8a and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B9 MEAN_INCIDENCE_ZENITH_ANGLE_B10	DOUBLE	Mean value containing viewing incidence zenith angle average for band B9 and for all detectors
MEAN INCIDENCE ZENITH ANGLE B11	DOUBLE	Mean value containing viewing incidence zenith angle average for band B11 and for all detectors
MEAN_INCIDENCE_ZENITH_ANGLE_B12	DOUBLE	Mean value containing viewing incidence zenith angle average for band B12 and for all detectors
MEAN_SOLAR_AZIMUTH_ANGLE	DOUBLE	Mean value containing sun azimuth angle average for all bands and detectors
MEAN_SOLAR_ZENITH_ANGLE	DOUBLE	Mean value containing sun zenith angle average for all bands and detectors
MEDIUM_PROBA_CLOUDS_PERCENTAGE	DOUBLE	Percentage of pixels classified as medium probability clouds
NODATA PIXEL PERCENTAGE	DOUBLE	Descentary of the Reference System (MOKS) the Percentage of No Data nivels
NOT_VEGETATED_PERCENTAGE	DOUBLE	Percentage of pixels classified as non-vegetated
PROCESSING_BASELINE	STRING	Configuration baseline used at the time of the product generation in terms of processor software version and major Ground
		Image Processing Parameters (GIPP) version
PRODUCT_ID	STRING	The full id of the original Sentinel-2 product
RADIATIVE_TRANSFER_ACCURACT	STRING	Accuracy of radiative transfer model Based on the OLOC reports contained in the Datastrins/OLDATA with PADIOMETRIC OUALITY checklist name
REFLECTANCE CONVERSION CORRECTION	DOUBLE	Earth-Sun distance correction factor
SATURATED_DEFECTIVE_PIXEL_PERCENTAGE	DOUBLE	Percentage of saturated or defective pixels
SENSING_ORBIT_DIRECTION	STRING	Imaging orbit direction
SENSING_ORBIT_NUMBER	DOUBLE	Imaging orbit number
SENSUK_QUALITY	STRING	Synthesis of the OLQU checks performed at granule (Missing_Lines, Corrupted_ISP, and Sensing_Time) and datastrip (Degraded SAD and Datation Model) levels
SOLAR IRRADIANCE B1	DOUBLE	Mean solar exotino solariti irradiance for band B1
SOLAR_IRRADIANCE_B2	DOUBLE	Mean solar exoatmospheric irradiance for band B2
SOLAR_IRRADIANCE_B3	DOUBLE	Mean solar exoatmospheric irradiance for band B3
SOLAR_IRRADIANCE_B4	DOUBLE	Mean solar exoatmospheric irradiance for band B4
SULAK_IKKADIANCE_B5	DOUBLE	Mean solar exoatmospheric irradiance for band B5 Mean solar exoatmospheric irradiance for band B6
SOLAR_IRRADIANCE_B0	DOUBLE	Mean solar exolatiospheric irradiance for band B7
SOLAR_IRRADIANCE_B8	DOUBLE	Mean solar exoatmospheric irradiance for band B8
SOLAR_IRRADIANCE_B8A	DOUBLE	Mean solar exoatmospheric irradiance for band B8a
SOLAR_IRRADIANCE_B9	DOUBLE	Mean solar exoatmospheric irradiance for band B9
SOLAR_IRRADIANCE_B10	DOUBLE	Mean solar exoatmospheric irradiance for band B10
SOLAR IRRADIANCE B12	DOUBLE DOUBLE	Mean solar exoatmospheric irradiance for band B11
SNOW_ICE_PERCENTAGE	DOUBLE	Percentage of pixels classified as snow or ice
SPACECRAFT_NAME	STRING	Sentinel-2 spacecraft name: Sentinel-2A, Sentinel-2B
THIN_CIRRUS_PERCENTAGE	DOUBLE	Percentage of pixels classified as thin cirrus clouds
UNCLASSIFIED_PERCENTAGE	DOUBLE	Percentage of unclassified pixels
VEGETATION_PERCENTAGE	DOUBLE DOUBLE	Percentage of pixels classified as wegetation Percentage of pixels classified as weter
WATER VAPOUR RETRIEVAL ACCURACY	DOUBLE	Declared accuracy of the Water Vapor model
	DOODLD	

DATASHEETS FOR DATASETS

Here we answer the questions outlined in the datasheets for datasets paper by Gebru et al. [28].

A. Motivation

For what purpose was the dataset created? The dataset was created for unsupervised pre-training in Earth observation. By integrating global coverage, multiple modalities and multiple timestamps, the dataset is intended to serve for diverse applications in remote sensing. The dataset fills the gap between multiple existing pre-training datasets, e.g. domain gap of ImageNet, regional coverage of BigEarthNet [13], single modality of SeCo [4], single timestamp of SEN12MS [3], and patch overlap of both SEN12MS and SeCo.

Who created the dataset (e.g., which team, research group) and on behalf of which entity (e.g., company, institution, organization)? The dataset was created by the lab "Data Science in Earth Observation" at Technical University of Munich and German Aerospace Center.

Who funded the creation of the dataset? The creation of the dataset was funded by the Helmholtz Association through the Framework of Helmholtz AI (grant number: ZT-I-PF-5-01) - Local Unit "Munich Unit @Aeronautics, Space and Transport (MASTr)". The computing resources for benchmark experiments were supported by the Helmholtz Association's Initiative and Networking Fund on the HAICORE@FZJ partition.

B. Composition

What do the instances that comprise the dataset represent (e.g., documents, photos, people, countries)? This dataset only contains satellite images. In addition we provide meta-data for these images, which contain information about data acquisition.

How many instances are there in total (of each type, if appropriate)? The dataset contains 251079 geographical patches, each patch including 3 product types and 4 seasons. In total there are 1M patches each for Sentinel-1 GRD, Sentinel-2 L1C and Sentinel-2 L2A, resulting in 1.5TB as three tar.gz files.

Does the dataset contain all possible instances or is it a sample (not necessarily random) of instances from a larger set? The dataset is a sample of all Sentinel-1/2 satellite images. While the dataset can still be extended, we make it as representative as possible by ensuring global coverage, multiple modalities and multiple timestamps.

What data does each instance consist of? Sentinel-1/2 images along with meta-data captured from the space.

Is there a label or target associated with each instance? No, our dataset is unlabeled. However, each patch is bound with geographical location and acquisition time, thus a match to other labeled maps is possible.

Is any information missing from individual instances? No.

Are relationships between individual instances made explicit (e.g., users' movie ratings, social network links)? Not applicable, though geographic location / acquisition time / product type / other properties can be extracted if needed.

Are there recommended data splits (e.g., training, development/validation, testing)? The dataset is intended for unsupervised pre-training. Users are free to use either the full split or a subset (either a subset of modalities or a subset of geographical patches) based on their targeted applications.

Are there any errors, sources of noise, or redundancies in the dataset? Yes, as mentioned in the data collection section, there are two kinds of noise/redundancies: first, potential overlap around grid cell boundaries; second, potential noise of clouds from inaccurate cloud filtering.

Is the dataset self-contained, or does it link to or otherwise rely on external resources (e.g., websites, tweets, other datasets)? The dataset is self-contained.

Does the dataset contain data that might be considered confidential (e.g., data that is protected by legal privilege or by doctor-patient confidentiality, data that includes the content of individuals' non-public communications)? No.

Does the dataset contain data that, if viewed directly, might be offensive, insulting, threatening, or might otherwise cause anxiety? No.

Does the dataset identify any subpopulations (e.g., by age, gender)? No.

Is it possible to identify individuals (i.e., one or more natural persons), either directly or indirectly (i.e., in combination with other data) from the dataset? No.

Does the dataset contain data that might be considered sensitive in any way (e.g., data that reveals race or ethnic origins, sexual orientations, religious beliefs, political opinions or union memberships, or locations; financial or health data; biometric or genetic data; forms of government identification, such as social security numbers; criminal history)? No.

C. Collection process

How was the data associated with each instance acquired? The data was collected from the publicly available Sentinel-1/2 database.

What mechanisms or procedures were used to collect the data (e.g., hardware apparatus or sensor, manual human curation, software program, software API)? Google Earth Engine with Python were used to collect the data.

If the dataset is a sample from a larger set, what was the sampling strategy (e.g., deterministic, probabilistic with specific sampling probabilities)? The patch locations are Gaussian sampled around a city center (50km) which is uniformly sampled from top-10k populated cities across the globe. The timestamps are sampled from four seasons (dates around Mar 20th, Jun 21st, Sep 22nd and Dec 21st) in the year 2020/2021.

Who was involved in the data collection process (e.g., students, crowdworkers, contractors) and how were they compensated (e.g., how much were crowdworkers paid)? The data was automatically collected and verified by the authors.

Over what timeframe was the data collected? The data was collected by the authors between February and March 2022. The images within the dataset were captured in the year 2020/2021.

Were any ethical review processes conducted (e.g., by an institutional review board)? No.

Did you collect the data from the individuals in question directly, or obtain it via third parties or other sources (e.g., websites)? The data was collected from open sources.

Were the individuals in question notified about the data collection? N/A.

Did the individuals in question consent to the collection and use of their data? N/A.

If consent was obtained, were the consenting individuals provided with a mechanism to revoke their consent in the future or for certain uses? N/A.

Has an analysis of the potential impact of the dataset and its use on data subjects (e.g., a data protection impact analysis) been conducted? N/A.

D. Preprocessing/cleaning/labeling

Was any preprocessing/cleaning/labeling of the data done (e.g., discretization or bucketing, tokenization, part-of-speech tagging, SIFT feature extraction, removal of instances, processing of missing values)? The data was pre-processed online during the collection/downloading process: filtering out cloudy patches and overlapping patches. No further pre-processing was done.

Was the "raw" data saved in addition to the preprocessed/cleaned/labeled data (e.g., to support unanticipated future uses)? No. The cloudy and overlapping patches were removed before downloading.

Is the software used to preprocess/clean/label the instances available? Yes, we use Google Earth Engine with Python which is freely available.

E. Uses

Has the dataset been used for any tasks already? In this paper we use the dataset to benchmark several self-supervised methods on several downstream tasks.

Is there a repository that links to any or all papers or systems that use the dataset? Yes we will organize and maintain all related information at https://github.com/zhu-xlab/SSL4EO-S12.

What (other) tasks could the dataset be used for? The main function of this dataset is to provide a pre-training dataset for both the study of self-supervised learning, and specific downstream applications. The dataset can also be used as a baseline for further pre-training datasets in Earth observation. In addition, the dataset can be used directly for applications like image retrieval, domain adaptation and style transfer.

Is there anything about the composition of the dataset or the way it was collected and preprocessed/cleaned/labeled that might impact future uses? We do not unify the orbiting (ascending/descending) of Sentinel-1 data, which should be taken into consideration for SAR related applications. However, the orbiting information can be found in the meta-data and the dataset can be further processed for targeting applications.

Are there tasks for which the dataset should not be used? The authors are not aware of any specific task that should be avoided.

F. Distribution

Will the dataset be distributed to third parties outside of the entity (e.g., company, institution, organization) on behalf of which the dataset was created? Yes, the dataset is publicly available.

How will the dataset will be distributed (e.g., tarball on website, API, GitHub)? The dataset is distributed as tarball on mediaTUM. Access to the dataset can be found at https://github.com/zhu-xlab/SSL4EO-S12.

When will the dataset be distributed? Starting from June 2022.

Will the dataset be distributed under a copyright or other intellectual property (IP) license, and/or under applicable terms of use (ToU)? CC-BY.

Have any third parties imposed IP-based or other restrictions on the data associated with the instances? No. Do any export controls or other regulatory restrictions apply to the dataset or to individual instances? No.

G. Maintenance

Who is supporting/hosting/maintaining the dataset? The dataset is hosted by mediaTUM and supported/maintained by the authors.

How can the owner/curator/manager of the dataset be contacted (e.g., email address)? The authors can be reached at their email addresses: {yi.wang, nassim.aitalibraham, conrad.albrecht, chenying.liu}@dlr.de, and {zhitong.xiong, xiaoxiang.zhu}@tum.de.

Is there an erratum? If errors are found an erratum will be added.

Will the dataset be updated (e.g., to correct labeling errors, add new instances, delete instances)? Any updates will be posted and the dataset will be versioned.

If the dataset relates to people, are there applicable limits on the retention of the data associated with the instances (e.g., were individuals in question told that their data would be retained for a fixed period of time and then deleted)? N/A.

Will older versions of the dataset continue to be supported/hosted/maintained? Depending on the updates (if there are), we will either continue hosting the older versions or make a clear update log that older versions can be generated from the newest version.

If others want to extend/augment/build on/contribute to the dataset, is there a mechanism for them to do so? Yes, please feel free to reach out to us.

H. Author statement of responsibility

The authors confirm all responsibility in case of violation of rights and confirm the licence associated with the dataset.

REFERENCES

- [3] Michael Schmitt et al. "SEN12MS a curated dataset of georeferenced multi-spectral Sentinel-1/2 imagery for deep learning and data fusion". In: ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences. 2019.
- [4] Oscar Mañas et al. "Seasonal contrast: Unsupervised pre-training from uncurated remote sensing data". In: *Proceedings of the IEEE/CVF International Conference on Computer Vision*. 2021.
- [13] Gencer Sumbul et al. "Bigearthnet: A large-scale benchmark archive for remote sensing image understanding". In: *IEEE International Geoscience and Remote Sensing Symposium*. 2019.
- [14] Noel Gorelick et al. "Google Earth Engine: Planetary-scale geospatial analysis for everyone". In: *Remote sensing of Environment* (2017).
- [22] Rodrigo Caye Daudt et al. "Urban change detection for multispectral earth observation using convolutional neural networks". In: *IEEE International Geoscience and Remote Sensing Symposium*. 2018.
- [26] Ramon Torres et al. "GMES Sentinel-1 mission". In: Remote sensing of environment (2012).
- [27] Matthias Drusch et al. "Sentinel-2: ESA's optical high-resolution mission for GMES operational services". In: *Remote sensing of Environment* (2012).
- [28] Timnit Gebru et al. "Datasheets for datasets". In: Communications of the ACM (2021).