


# OBSERVATION AND REAL-TIME ANALYSIS OF SPACE WEATHER EFFECTS ON GNSS THROUGH GNSS MEASUREMENTS





*Jens Berdermann*

*German Aerospace Center*

*Institute for Solar-Terrestrial Physics*



# Space Weather is challenging for GNSS based navigation



## Climatological variations

- Radiation
- Solar Wind

## Space Weather Events

- Solar Flares
- Radio Bursts
- CMEs
- SEPs

- Solar cycle, Solar rotation
- Day-Night
- Coupling Processes
- Seasons
- Region
- Forcing from below (Gravity Waves)

Ionospheric plasma causes

- Delay of the radio signals
- Signal strength fluctuations
- Defocussing of the signal

➔ **Excess of Distance**



➔ **Loss of signal**




# GNSS based Space Weather observation

**Ground based GNSS** observation providing TEC measurements are recently one of the most important data source for Space Weather research and services.

- Global coverage
  - Multi-frequency, multi-GNSS
  - Good horizontal resolution
  - High temporal resolution
  - (Near) Real time
- 
- Bad coverage over ocean and mountain regions
  - No vertical resolution




# GNSS based Space Weather observation




**Space based GNSS measurements on board LEO Satellites** play an increasing role in ionospheric monitoring

Radio Occultation




Topside



- Global coverage
- Good vertical resolution





- Near real time capability depends on data download
- High spatial resolution depends on number of satellites






# Space Weather Impact on GNSS – Solar Flare


Space Weather Event 06/09/2017 (Solar Flare X9.3, Ranking 14)



Maximal Rate of TEC index - 1 min update 2017-09-06T11:45:00



Maximal Rate of TEC index - 1 min update 2017



DLR Real Time ROTI in one minute  
Resolution at [impc.dlr.de](http://impc.dlr.de).

# Space Weather Impact on GNSS – Solar Flare




**Source:** X9.3 Solar Flare on 6<sup>th</sup> September 2017


**Region:** Earth Day Side

**Duration:** Minutes


**Impact:** Solar flares with a strong EUV component around 30 nm can seriously affect GNSS positioning services used in e.g. aviation, maritime navigation. All the GNSS satellite systems in view were affected in a similar way, including GPS, GLONASS and Galileo.




## Loss of Lock



## EGNOS




## Precise Point Positioning





Berdermann, J., Kriegel, M., Banys, D., Heymann, F., Hoque, M. M., Wilken, V., et al. (2018). **Ionospheric response to the X9.3 Flare on 6 September 2017 and its implication for navigation services over Europe.** *Space Weather*, 16. <https://doi.org/10.1029/2018SW001933>



# Space Weather Impact on GNSS – Radio Burst



Solar radio observation on the 6 September 2017



Ondrejov solar radio spectrum in the 1.0- to 2.0-GHz range (top) and flux intensity near the GPS frequencies (bottom).

Signal to noise ratio for different GPS frequencies at high, mid and low latitudes.

**Source:** Solar Radio Burst on 6<sup>th</sup> September 2017



**Region:** Earth Day Side

**Duration:** Minutes

**Impact:** The solar radio pulsation caused larger SNR reduction for GPS L2/L5 and GALILEO L5 frequencies. All the GNSS satellite systems in view were affected in a similar way, including GPS, GLONASS and Galileo.

Sato, H. , Jakowski, N. , Berdermann, J. , Jiricka, K., Heßelbarth, A. , Banyś, D. , Wilken V. (2018), *Solar Radio Burst Events on 6 September 2017 and Its Impact on GNSS Signal Frequencies*. Space Weather, 16. <https://doi.org/10.1029/2018SW001933>

# Space Weather Impact on GNSS – Ionospheric Storm



**Source:** Ionospheric storm on 27.02.2014


**Region:** North America

**Duration:** ca. 3h

**Impact:** Outages of SBAS due to storm induced Ionospheric Disturbances

No LPV availability of WAAS over Alaska on 27th February 2014.

(**L**ocalizer **P**erformance with **V**ertical **G**uidance)






# Characterization and Prediction of Ionospheric Disturbances




Disturbance Ionosphere Index Spatial Gradient (DIXSG) as a measure for ionospheric storm impact




Ionospheric storm on the 17th March 2015 „St. Patrick Day Storm“

Important to develop methods and indices providing direct information on the performance of precise and safety-critical GNSS applications.



Wilken, V., Kriegel, M., Jakowski, N., Berdermann, J., (2018), **An ionospheric index suitable for estimating the degree of ionospheric perturbations**, Space Weather Space Clim. 8 A19 (2018) DOI: 10.1051/swsc/2018008

# Space Weather Impact on GNSS – Small scale irregularities



## Polar Region:

**Source:** Geomagnetic Storms (Polar Region)

**Impact:** GNSS Signal is disturbed by gradients and may be lost in severe case.

## Equatorial Region:

**Source:** Flow inversion of the equatorial plasma during evening hours

**Impact:** GNSS Signal is disturbed by ionospheric irregularities (plasma bubble) and may be lost in severe case.


Amplitude scintillation can cause stripes on L-band SAR images. Scintillation cause loss of spatial resolution.



# Space Weather Impact on GNSS – Small scale irregularities




Plasma Bubble degrades availability of GNSS Precision Approach




Two divergent Information can cause hazardous misleading situations.

GNSS Signal is disturbed by ionospheric irregularities (plasma bubble) and may be lost in severe case.





Scintillation effects on UHF satellite communication systems





Risk of communication outage due to scintillation.

# PECASUS for ICAO

The PECASUS consortium is one of the four global centers providing space weather advisories according to ICAO regulations.




Continuous NRT delivery of GNSS related space weather data and scientific support in case of events



PECASUS DASHBOARD									
PECASUS DUTY STATUS: On Duty Centre									
GNSS	Moderate	Severe	Time UTC	Values	Status	Alert	Max-3h values	Max-3h status	
Amplitude Scintillation	0.5	0.8	2021-01-07 09:20	0.28	QUIET		0.35	QUIET	
Phase Scintillation	0.4	0.7	2021-01-07 09:20	0.10	QUIET		0.18	QUIET	
Vertical TEC	125	175	2021-01-07 09:25	39.77	QUIET		39.77	QUIET	
RADIATION	Moderate	Severe	Time UTC	Flags	Status	Alert	Max-3h flags	Max-3h status	
Effective Dose FL5400	30	80	2021-01-07 09:30	0	QUIET		0	QUIET	
Effective Dose FL > 460	/	80	2021-01-07 09:30	0	QUIET		0	QUIET	
HF COM	Moderate	Severe	Time UTC	Values/Flags	Status	Alert	Max-3h values	Max-3h status	
Auroral Absorption (AA)	8	9	2021-01-07 09:27	1.0	QUIET		2.0	QUIET	
Polar Cap Absorption (PCA)	2	5	2021-01-07 09:30	0.00	QUIET		0.00	QUIET	
Shortwave Fadeout (SWF)	x1.0	x10.0	2021-01-07 09:30	< M.5-flare	QUIET		< M.5-flare	QUIET	
Post-Storm Depression (PSD)	30%	50%	2021-01-07 09:30	0	QUIET		0	QUIET	

Sound alarm is triggered when MOD or SEV thresholds are exceeded.  
Alarm will NOT ring for OUTDATED data status.

ICAO Space Weather Advisories



Data Analysis, Event detection, Decision Making, Alerting



FMI  
Finnish Meteorological Institute



Met Office  
Met Office UK



DLR  
The German Aerospace Center



KNMI  
Royal Netherlands Meteorological Institute



STCE  
The Solar-Terrestrial Centre of Excellence in Belgium



FU  
Frederick University (Cyprus)



SRC  
Space Research Center of the Polish Academy of Sciences



SL  
The Seibersdorf Laboratories in Austria



INGV  
Istituto Nazionale di Geofisica e Vulcanologia (Italy)




SANS  
South African National Space Agency



# ADS-Messages



- **A**utomated **D**ependent **S**urveillance is a system to monitor and control flight routes
  - **A**ddressed (sent on request)
  - **B**roadcast (sent continuously)
  - **C**ontract (transmitted as part of a data link)
- Possible space weather impacts:
  - Data gaps due to signal loss (UHF, SATCOM)
  - Position errors (GNSS)



## ADS-B

- Aircraft information
- Position information
- Speed information

Research  
perspective:  
High-rate and -  
resolution,  
global data set





## ADS-C

- Continuous data link
- Aircraft information
- Position information
- Navigation information

# Space weather effects on ADS-B: Flare 1<sup>st</sup> May 2023

(Automatic Dependent Surveillance – Broadcast)




- Approx. 14000 aircraft recorded during M-class flare 1<sup>st</sup> May 2023 (13:02-13:09)
- Expected impacts: data gaps, position errors






# Examples – data appropriate for analysis?




On route




Takeoff and landing




Type



Glider





Helicopter



Drone

# Space weather effects on ADS-B: Flare 1<sup>st</sup> May 2023

(Automatic Dependent Surveillance – Broadcast)



Red dots: ADS-B messages during flare (13:02-13:09)


Arrow: Flight direction

1 source: <https://www.jetphotos.com/photo/11081148>


2 source: [https://commons.wikimedia.org/wiki/File:Piper\\_PA-28R-200\\_Cherokee\\_Arrow\\_%28D-EATT%29\\_02.jpg](https://commons.wikimedia.org/wiki/File:Piper_PA-28R-200_Cherokee_Arrow_%28D-EATT%29_02.jpg)



# Discussion ADS-B



- Position errors:
  - How to differentiate from other impacts?
  - How to detect smaller position errors successful during flares?
- Other Problems:
  - No messages over certain countries/regions
  - Different rates depending on type/source
- Open challenges:
  - Definition of position error
  - Definition of data gaps
  - Selection of impacted aircraft



# ADS-C messages: Events






ADS-C data link is established between an air traffic service (ATS) unit and an aircraft in order to exchange standard information as well as contract-specific information.



Analysis of ADS-C downlink reports for September 2017  
(approx. 54000 messages)



Regularly established contracts: **Periodic**  
**Waypoint**

Irregularly established contracts: **Lateral deviation event**  
**Altitude range change event**  
**Vertical rate change event**







# ADS-C-Messages: Context space weather?



# ADS-C messages on different solar activity.







We thank the Airways Corporation of New Zealand and the FAA William J. Hughes Technical Center for making the ADS-C records available. We thank Klaus Sievers (Vereinigung Cockpit) for forwarding the datasets and providing feedback.



# Recent Steps:

- To analyze the impact of weather and space weather
- Analyze more data sets if available



# SWIGPAD (Space Weather Impact on GNSS Performance Application Development)



- A software application for evaluating the effects of space weather on GNSS positioning addressing 6 different user groups
- Based on data products of the ESA Ionospheric Weather Expert Center (ESA SW Portal)

**Advanced Options**

**Accuracy & Deviation**

☒ Use presets based on the selected User Type

Confidence level constraints

1  $\sigma$

Coordinate error constraints

North East Vertical

1 m 1 m 4 m

**Constellations**

☒ GPS

☒ GLONASS

☒ Galileo

☒ BEIDOU

☒ QZSS

**Timeseries**

☐ Enable Timeseries

Date & Time - Start (UTC)

Date & Time - End (UTC)

Number of intervals

**GNSS Performance Indicators** About Help

**Input**

Latitude Longitude

53,34 °N 13,05 °E

User Type

USR01 - Single frequency, average accuracy, no ii

Date & Time (UTC)

2023-05-05T10:46

**Advanced Options**

Reset Submit

**Output**


No results yet.

This web page forms part of the ESA Space Safety Programme's network of space weather service development activities, and is supported under ESA contract number 4000131051/20/D/CT. For further product-related information or enquiries contact helpdesk. E-mail: [helpdesk.swe@ssa.esa.int](mailto:helpdesk.swe@ssa.esa.int)

All publications and presentations using data obtained from this site should acknowledge the Ionosphere Monitoring and Prediction Center operated by the German Aerospace Center (DLR) and The ESA Space Safety Programme.

For further information about space weather in the ESA Space Safety Programme see: [www.esa.int/spaceweather](http://www.esa.int/spaceweather)

Access the S2P-SWE Portal here: [swe.ssa.esa.int](http://swe.ssa.esa.int)








Work has been performed in the frame of ESA Space Safety Programme's network of Space Weather Service development and pre-operational activities, supported under ESA Contract 4000131051/20/D/CT.




# Current Space Weather situation is optimal for impact studies!

Multiple space weather events with moderate and severe geomagnetic storms impacting satellite navigation are expected on the upcoming path to solar maximum



Ionospheric storm based gradients and scintillation occurrence on February 27<sup>th</sup>.

# Thank you!



## Contact:

Jens Berdermann  
Institute for Solar-Terrestrial Physics

[Jens.Berdermann@dlr.de](mailto:Jens.Berdermann@dlr.de)  
+49 3981 480 106