ELECTRON SCATTERING BY ELECTROSTATIC ELECTRON CYCLOTRON HARMONIC WAVES DURING STORM-TIME

Katja Stoll^{1,2}, Leonie Pick¹, Dedong Wang³, Bernhard Haas^{2,3}, Xing Cao⁴, Binbin Ni⁴, Yuri Shprits^{2,3,5}

¹ Institute for Solar-Terrestrial Physics, German Aerospace Centre (DLR), Neustrelitz, Germany ² Institute of Physics and Astronomy, University of Potsdam, Rotsdam, Germany GAZ German Research Centre for Geosciences, Potsdam, Germany ⁴ Department of Space Physics, School of Electronic Information, Wuhan University, Wuhan, China ⁵ Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles, CA, USA

Katja Stoll, IMC IV, June 04, 202

Motivation



- Haas et al. (2023): missing loss process in the electron ring current
- Are ECH waves responsible for (part of) the missing loss?



Electrostatic electron cyclotron harmonic waves





ECH wave-induced diffusion coefficients L = 2.4 L = 3.0 L = 3.4



VERB-4D Simulations





Katja Stoll, IMC IV, June 04, 2024

BACKUP SLIDES

DLR

Magnetic field and plasma density





Fits to phase space density





Wave amplitude





Lifetimes



Lifetimes in VERB-4D





(a) ECH lifetimes at ${\it E}=1\,{\rm keV}$, ${\it \alpha}=50\,^{\circ}$

(b) ECH lifetimes at E = 5 keV, $\alpha = 50^{\circ}$