

<https://doi.org/10.1038/s41526-026-00644-7>

A computational model of altered neuronal activity in altered gravity

Check for updates

Camille Gontier^{1,2}, Laura Drouvé³, Johannes Striebel⁴, Maximilian Sturm³, Zoe Meerholz³, Sarah Schunk^{3,4}, Yannick Lichterfeld³ & Christian Liemersdorf³

Electrophysiological experiments have shown that neuronal activity changes upon exposure to altered gravity. More specifically, neurons' firing rates increase during microgravity and decrease during centrifugal-induced hypergravity. Different biophysical explanations have been proposed for this phenomenon; however, they have not been backed by quantitative analyses or simulations. More generally, classical computational models of neurons and networks do not account for the effect of altered gravity, which limits the possibility to perform in-silico experiments and simulations. Here, we propose computational implementations for different effects of altered gravity on cellular functions, and modify existing models to account for the effect of micro- and hyper-gravity. Firstly, in line with previous experiments, we suggest that microgravity could be modeled as an increase in the voltage-dependent channel transition rates, which is assumed to be the result of a higher membrane fluidity and can be readily implemented into the Hodgkin–Huxley model. Using in silico simulations of single neurons, we show that this model of the influence of gravity on neuronal activity allows for reproducing the observed increased firing and burst rates. Secondly, we explore the role of mechano-gated (MG) ion channels on population activity. We show that recordings can be fitted by a network of connected excitatory neurons, whose activity is balanced by firing rate adaptation. Adding a small depolarizing current to account for the activation of MG channels also reproduces the observed increased firing and burst rates. Overall, our results fill an important gap in the literature, by providing a computational link between altered gravity and neuronal activity. Starting from historical observations of the effects of gravity on cellular functions, we derived gravity-sensitive models of neurons and networks, whose predictions could be refined using future experiments.

During spaceflight, humans experience a variety of physiological changes due to deviations from familiar Earth conditions. Specifically, the lack of gravity is responsible for a variety of effects observed in astronauts, including a loss of muscle mass and bone density as a result of musculoskeletal deconditioning^{1–3} and modifications of the vestibular and cardiovascular systems immediately upon microgravity exposure⁴. These impairments also include structural, as well as functional changes of the brain: an elevated intracranial pressure possibly leading to neuro-ophthalmic anatomical changes (spaceflight-associated neuro-ocular syndrome, or SANS)^{5–7}, an increased total ventricular volume^{7,8}, and altered neurotransmitter functions and neurotrophic factors⁹.

Given the rising interest and need to conduct experiments in microgravity conditions and the cost and complexity of accessing the International Space Station (ISS), different gravity research platforms have been developed on Earth to study the impact of gravity. Different types of 2D clinostats are mostly used to simulate microgravity in the lab^{10,11}. Real microgravity can be achieved by larger gravity research platforms, including drop towers¹², sounding rockets^{13,14}, and parabolic flights with large aircraft^{15–17}, single-engine aircraft¹⁸, and gliders¹⁹. Similarly, the effect of hypergravity can be studied using centrifuges^{20,21} or an aircraft performing high-bank turns²².

Several of these platforms have already been employed to study how neuronal activity is impacted under altered gravity. Previous

¹LIDE Space, Louvain-la-Neuve, Belgium. ²INRIA Center at University of Lorraine, Strasbourg, France. ³Department of Applied Aerospace Biology, Institute of Aerospace Medicine, German Aerospace Center, Cologne, Germany. ⁴Department of Ophthalmology, Medical Faculty, University of Bonn, Bonn, Germany.

e-mail: Christian.Liemersdorf@dlr.de

electrophysiological experiments have revealed different effects, including a higher firing rate^{23–25}, as well as a higher intracellular calcium concentration¹¹, following exposure to microgravity. Similarly, experiments using multi-electrode arrays (MEA) reported an increase in the firing rates of neurons during microgravity phases and opposing effects upon exposure to hypergravity²¹. These initial changes in spontaneous action potential spiking frequency were compensated within a minute time range, hinting at an adaptation mechanism.

Different phenomena have been proposed to explain the increased firing rate during microgravity and the decreased firing rate during hypergravity. First, a possible explanation is that microgravity increases the membrane fluidity of neuronal cells, which modifies properties of the ion channels. More specifically, a previously proposed model²⁴ hypothesizes that an increase in membrane fluidity modifies the open-state probabilities of ion channels, hence leading to a depolarization of the cell and to an increase in the spontaneous firing rate. Although intuitive, this model is purely qualitative and has not been backed by analytical computations or simulations. In computational models, a neuron is often described by a set of parameters, e.g., its membrane time constant τ_m or its ion channel conductance g . A proper description of how these parameters might be impacted by altered gravity conditions is currently missing.

A second explanation for the increased firing and burst rates in microgravity is based on the role of mechanosensitive ion channels, which operate as gravireceptors in eukaryotic cells²⁶. They have been shown to be widely present in the nervous system²⁷, including hippocampal²⁸ and pyramidal neurons²⁹. These mechano-gated (MG) ion channels have been observed to generate inward currents and to elicit spiking activity in neuronal cells²⁹. The modified neuronal activity during periods of altered gravity has thus been attributed to the activity of MG ion channels, whose stretching during acceleration and deceleration phases might impact neuronal activity²¹. But the final effect of an inward current on firing rates is going to strongly depend on the properties of the studied neuron population. Specifically, these MG channels would have different effects depending on whether the observed neurons are isolated or show signs of synchronized activity; whether their recurrent connections are mostly excitatory or inhibitory; or whether they are impacted by non-linear dynamical effects, such as short-term depression or firing rate adaptation. To date, these features have not been characterized in the neuron cell types used in altered gravity experiments.

First, in section “A proposed model of the effect of gravity on ion channels’ gating rates”, we propose that the effect of altered gravity on neuronal dynamics could be represented as an acceleration of the rate at which voltage-dependent ion channels switch from open to closed configurations, i.e., as a decrease of the time constants $\tau_n(V)$, $\tau_m(V)$, and $\tau_h(V)$. This is in line with a set of previous observations reporting a modification of the fluidity of the cell membrane in altered gravity, and a direct link between membrane lipids and ion channels. We verify that decreasing these time constants during in-silico simulations allows us to reproduce the increased firing and burst rates observed during altered gravity experiments (section “In-silico simulations of single cell activity under altered gravity”).

Second, in section “Observed synchronized bursts can be explained by spike frequency adaptation”, inspired by ref. 30, we propose a computational model to explain the activity of networks of neurons in microgravity. Human stem cell-derived neuronal cells, which have already been used in microgravity experiments, have two remarkable features: they are mostly excitatory, and show slow oscillations of coordinated activity³¹. We show that these features can be accounted for by a network of connected excitatory leaky integrate-and-fire (LIF) neurons whose recurrent connections are balanced by firing rate adaptation. We verify that injecting an external depolarizing current into the network (to model the activation of MG channels) also allows us to reproduce the increased firing and burst rates observed during altered gravity experiments (section “In-silico simulations of single cell activity under altered gravity”).

Results

A proposed model of the effect of gravity on ion channels’ gating rates

A classically used model of the time evolution of the membrane potential of a neuron, and especially of how action potentials are generated, is called the Hodgkin–Huxley model^{32,33}. It consists of a set of differential equations, describing the evolution of the cell voltage as a function of its ion channels conductances; the evolution of these conductances as a function of the open probabilities of the channels; and the evolution of these probabilities as a function of their voltage-dependent gating rates.

The standard equation describing the time evolution of the voltage V of a single-compartment model of a neuron (see³³ for a detailed derivation) is

$$C_m \frac{dV}{dt} = -i_m \quad (1)$$

where C_m is the specific membrane capacitance (in $F \cdot m^{-2}$) and i_m is the total membrane current. The latter can be expressed as the sum of a delayed-rectified potassium current, a transient sodium current, and a leakage current:

$$i_m = \bar{g}_K n^4 (V - E_K) + \bar{g}_{Na} m^3 h (V - E_{Na}) + \bar{g}_L (V - E_L) \quad (2)$$

where E_L , E_{Na} , and E_K are the reversal potentials, and $\bar{g}_L$, $\bar{g}_{Na}$, and $\bar{g}_K$ are the maximal surface conductances (in $S \cdot m^{-2}$). n corresponds to the probability that a subunit in a potassium channel is activated and hence contributes to the opening of the K^+ channel. Similarly, m and h express the activation and inactivation probabilities of the Na^+ conductances. The time evolution of these gating variables can also be described by a set of differential equations:

$$\frac{dn}{dt} = \alpha_n(V)(1 - n) - \beta_n(V)n \quad (3)$$

$$\frac{dm}{dt} = \alpha_m(V)(1 - m) - \beta_m(V)m \quad (4)$$

$$\frac{dh}{dt} = \alpha_h(V)(1 - h) - \beta_h(V)h \quad (5)$$

where $\alpha_x(V)$ and $\beta_x(V)$ are the voltage-dependent opening and closing rates of the subunit gates, which are usually obtained by fitting on experimental observations (see ref. 33 for a detailed discussion). Eq. (3) can also be written as

$$\tau_n(V) \frac{dn}{dt} = n_\infty(V) - n \quad (6)$$

with

$$\tau_n(V) = \frac{1}{\alpha_n(V) + \beta_n(V)} \quad (7)$$

and

$$n_\infty = \frac{\alpha_n(V)}{\alpha_n(V) + \beta_n(V)} \quad (8)$$

(similar equalities can be derived for the other gating variables m and h). The time constants $\tau_n(V)$ (Eq. 6), $\tau_m(V)$, and $\tau_h(V)$ thus govern the dynamics of channel subunits, i.e., the rate at which they open and close.

Altered levels of gravity have been observed to act on the activity of single ion channels^{34–36}, but more importantly have been shown to act on the fluidity of the cell membrane itself. Studies using random positioning machines (RPM) reported a dramatic increase in the fluidity of cell membrane models under simulated microgravity conditions³⁷. The increased

fluidity under microgravity (and reciprocally the reduced fluidity during hypergravity) has been further observed in different membranes³⁸ and under different altered gravity modalities^{23,25}.

In a previously proposed model^{23,24}, a link between gravity-induced changes in fluidity and firing rates was suggested: it hypothesized that an increase in membrane fluidity modifies the properties of ion channels, hence leading to a depolarization of the cell and to an increase in the spontaneous firing rate. Although heuristic, this model is purely qualitative and has not been backed by quantitative analyses or simulations.

However, several independent studies reported that ion channel functions are regulated by the cell membrane's lipid microdomains³⁹, and that lipid binding sites exist on ion channels, yielding fluidity-dependent interactions between them⁴⁰. More specifically, arachidonic acid has been shown to increase the rate of voltage-gated calcium channel activation kinetics⁴¹, while lipid gating influences the activity of potassium channels⁴². Additionally, a modulation of acetylcholine receptors by membrane lipids was highlighted⁴³ (although the lipid composition is more important than bulk membrane fluidity for ion channel function⁴⁴), as well as a link between medium temperature, membrane fluidity, and channels closing rate⁴⁵.

We propose to represent the effect of microgravity on computational models of neurons by a unitless multiplicative constant λ , which is applied to the opening and closing rates $\alpha_x(V)$ and $\beta_x(V)$, and hence to the time constants $\tau_n(V)$, $\tau_m(V)$, and $\tau_h(V)$ (Fig. S1). The classical gating equations (Eqs. 3–5) thus become

$$\frac{dn}{dt} = \lambda(\alpha_n(V)(1 - n) - \beta_n(V)n) \quad (9)$$

$$\frac{dm}{dt} = \lambda(\alpha_m(V)(1 - m) - \beta_m(V)m) \quad (10)$$

$$\frac{dh}{dt} = \lambda(\alpha_h(V)(1 - h) - \beta_h(V)h) \quad (11)$$

with $\lambda = 1$ in 1g, $\lambda < 1$ in hypergravity, and $\lambda > 1$ in microgravity. Figure S1 shows that, for a given voltage, increasing the value of λ decreases the gating time constants (left panel) without modifying the gating steady-state levels (right panel). Our approach allows us to reconcile two independent observations: the fact that microgravity impacts the cell membrane fluidity^{23,25,37,38}, and the fact that membrane lipids impact the channels' functions^{39–45}.

In silico simulations of single cell activity under altered gravity

Next, we sought to assess the effect of λ on simulated neuronal activity, and especially to verify that a higher value of λ (to represent the effect of microgravity) would reproduce the different features observed during experiments, namely a higher firing rate and burst rate—and that a value of λ set below 1 would reproduce the reduced activity observed following the onset of hypergravity in centrifuge experiments. Simulations were performed using the Brian neuron simulator⁴⁶ and the numerical values described in Methods.

Figure 1A shows the firing rate of a neuron simulated by the Hodgkin–Huxley model as a function of the input current I added to Eq. (2) and of the value of λ in Eqs. (9–11). For a given input current, the firing rate increases with λ , which is in line with experimental observations. We verified that this result is robust to the values used in the simulation by randomly varying the values of the neuron's parameters and checking that a higher value for λ robustly leads to a higher firing rate (Fig. S2). Figure 1B shows the time evolution of the voltage of the same neuron for an input current $I = 0.35$ nA. A shorter time constant for the gating variables leads to a faster voltage increase, an earlier time to first spike, and a shorter inter-spike interval.

Figure 1C shows the effect of λ on the open probabilities of the potassium and sodium channels, i.e., $P_K = n^4$ (left column) and $P_{Na} = m^3h$ (right column). The upper row shows their time evolution in the same

condition as in Fig. 1B, while the bottom row shows the average value of P_K and P_{Na} during 2 s of simulation (normalized by the number of action potentials). The end results in the lower row are independent of I , since the firing rate is nearly linear with I for sufficiently large values (see Fig. 1A). These results show that, under our modeling assumptions, an increase in λ does not increase the opening probabilities of the ion channels, but rather accelerates the time at which the Na^+ conductances activate. Previous studies had reported seemingly contradictory results on the effect of altered gravity on ion channel opening probabilities^{34–36}. Additionally, previous qualitative models of the increased firing rates observed in microgravity were based on the assumption that microgravity exposure would reduce the ion channels' opening probabilities, ultimately yielding an increased firing rate²⁴. However, ion channels are dynamical systems, for which the opening probability cannot be reduced to a single time-invariant value. Furthermore, they are not isolated, and their dynamics are voltage-dependent and correlated to the activity of other channels in the cell. Here, we suggest a more nuanced approach to the effect of gravity on ion channels, in which altered gravity does not simply modify the baseline opening probabilities of channels but rather acts on their dynamics.

Finally, we also verified that varying the value of the parameter λ allows us to reproduce experimental observations not only in isolated neurons, but also in more realistic networks of recurrently connected neurons. To do so, we simulated the activity of a population of $N = 4000$ connected Hodgkin–Huxley neurons (whose parameters are described in Table 1), and verified that increasing the value of λ leads to an increase in both the population firing rate (Fig. 1D, left) and burst rate (Fig. 1D, right). In line with previous studies²¹, we define a burst as a group of consecutive spikes separated by less than 20 ms. These results are akin to what has been previously observed during altered gravity experiments²¹.

Observed synchronized bursts can be explained by spike frequency adaptation

In addition to the role of cell membrane fluidity, another possible explanation for the altered neuronal activity involves MG ion channels, whose activation in altered gravity phases would depolarize cells²¹. However, although this explanation makes sense for single isolated neurons, the effect of an inward current on recurrently connected neurons would depend on the properties of the network. More specifically, it would depend on its excitation-to-inhibition ratio, and on the presence of non-linear dynamics, such as short-term plasticity or firing rate adaptation. Such features have not been characterized in the cell lines classically used in altered gravity experiments.

Such cell lines usually include human stem cell-derived neuronal cells (i.e., iNGN cells) (see ref. 31 and ref. 21 for a detailed discussion), which have two main remarkable features. Firstly, their population activity shows synchronized bursts and action potentials (Fig. 2A, and Figure 3C in ref. 31) at a low frequency (i.e., <1 Hz), which is a telltale sign of recurrent connections among neurons. Secondly, the vast majority of these connections are supposed to be excitatory: synchronized activity in iNGN neuronal cell lines is abolished by glutamate receptor blockers, and although inhibitory GABAergic synapses appear in the last stages of cell development, inhibitory neurons only make up approximately 2.3% of the whole population after 30 days of cell growth⁴⁷. This is significantly below the level of inhibition which is classically assumed to be necessary for stable self-sustained population activity⁴⁸.

Similar low-frequency oscillations had been previously observed in neuronal populations, especially during sleep, anesthesia, or in-vitro when using a mock cerebrospinal fluid (CSF)^{49,50}. Moreover, different computational models have been proposed to reproduce and explain them. In⁴⁹, the authors augment the classical Hodgkin–Huxley model with a slow sodium-dependent potassium current, which mediates the transition between up and down states (respectively characterized by higher and lower levels of spontaneous activity). However, this model relies on an important population of inhibitory neurons to stabilize the strong recurrent excitation during the up state. In⁵¹, the authors propose a simpler model (based on LIF




Fig. 1 | In silico simulations of single cell activity under altered gravity.

A Frequency-current curve of a simulated Hodgkin–Huxley neuron (whose parameters are described in Table 1) for different values of λ . Inset: firing rate as a function of λ for $I = 0.7$ nA. **B** Time evolution of the membrane voltage for the same cell as in **A**. Inset: zoom on the first ms of the simulation, showing the earlier depolarization induced by a higher value for λ . **C** Evolution of potassium P_K (left

column) and sodium P_{Na} (right column) channel opening probabilities for the same cell as in **A**. Upper row: time evolution of these probabilities for $I = 0.35$ nA. Lower row: average (over $T = 2$ s) value of these probabilities (normalized by the number of spikes during the simulation). **D** Mean firing and burst rates (averaged across all neurons) in a network of connected Hodgkin–Huxley neurons as a function of λ . Solid line: mean value over 30 repetitions. Shaded area: s.e.m.

neurons) to reproduce these slow oscillations, but here again the recurrent excitation needs to be balanced by inhibitory neurons. This does not coincide with the very low proportion of inhibitory neurons in the observed population in ref. 21.

However, other phenomena, such as firing rate adaptation⁵² and short-term depression⁵³, have been proposed as ways to stabilize the coordinated activity of connected excitatory neurons without inhibitory feedback. In line with⁵⁴, we model the evolution of the voltage potential V in the presence of an injected current I using the adaptive Exponential Integrate-and-Fire model (AdEx), which can be seen as a simplification of Eq. (2) in that it neglects the individual ion-specific dynamics:

$$C_m \frac{dV}{dt} = -g_L(V - E_L) + g_L \Delta_T \exp\left(\frac{V - V_T}{\Delta_T}\right) + I - w \quad (12)$$

where C_m is the cell total membrane capacitance, g_L is the total leak conductance, E_L is the effective rest potential, Δ_T is the threshold slope factor, and V_T is the effective threshold potential. The last term w (which is not present in Eq. 2) tends to hyperpolarize the cell following a spike, hence reducing the probability of a subsequent spike and leading to spike-rate adaptation. It is increased by a quantity b following each spike, and otherwise decays with a time constant τ_w :

$$\tau_w \frac{dw}{dt} = a(V - E_L) - w \quad (13)$$

We verify that the stable self-sustained slow oscillations observed in neuronal cell lines (Fig. 2A) can be reproduced by a model consisting of a

population of LIF neurons recurrently connected by excitatory synapses, whose activity is balanced by firing rate adaptation, without the need for an inhibitory population (Fig. 2B). Across all simulated neurons, the average firing rate was 10.85 ± 1.39 Hz and the average burst rate was 0.44 ± 0.05 Hz, which is in agreement with the experimental values reported in Fig. 2D, E in ref. 21. Note that using adaptation to induce regular bursting in in-silico networks had already been implemented in a different study, although with different models of neurons and synapses and still assuming a small subpopulation (5%) of inhibitory neurons³⁰.

The model reproduces features of network activity observed in altered gravity

Finally, after having identified the main features of the neuron networks used in altered gravity experiments (i.e., excitatory recurrent connections and population activity stabilized by firing rate adaptation), we verified that injecting an additional current onto this network (to represent the activity of MG channels) allows us to reproduce the increased firing and burst rates observed during the dynamic phases of altered gravity experiments. This acceleration-induced current is represented by an extra term I_λ in the right-hand side of Eq. (12):

$$C_m \frac{dV}{dt} = -g_L(V - E_L) + g_L \Delta_T \exp\left(\frac{V - V_T}{\Delta_T}\right) + I - w + I_\lambda \quad (14)$$

Figure 3A shows the raster plot for the same network as in Fig. 2B, for $I_\lambda = 0$ pA (top) and $I_\lambda = 10$ pA (bottom). Figure 3B shows the main firing rate (left) and burst rate (right) of the network (averaged over all neurons) as a function of I_λ . Both the firing and burst rates increase with I_λ , which is in line

with experimental observations. More specifically, hypergravity experiments using the same hiPSC-derived neurons on the DLR human centrifuge in Cologne, Germany have reported decreased neuronal activity compared to baseline recordings during exposure to 6g hypergravity (as detailed in section In-silico simulations of single cell activity under altered gravity), but also increased activity during both the ramp-up and ramp-down phases of the centrifuge (Figure 2D and E in ref. 21). These ramp-up and ramp-down phases correspond to a dynamic modification of the perceived acceleration, during which MG channels are likely to be activated and to contribute to the depolarization of the cells.

In addition to its potential effect on ion channel dynamics (which was previously suggested in section “A proposed model of the effect of gravity on

ion channels’ gating rates”), the gravity-induced modification of cellular membrane fluidity^{23,25,37,38} could also impact the activity of cell networks by modifying synaptic transmission. Indeed, both exocytosis and endocytosis, which are critical for synaptic transmission, are dependent on the fluidity of the cell membrane: a high tension will increase the exocytosis rate^{55,56} while also decreasing the endocytosis rate^{57,58}. These phenomena are still observed in the active zones of nerve terminals⁵⁹ and happen over a fast time scale, making them potentially relevant for the rapid modification of neuronal activity following the onset of altered gravity. Since both endocytosis and exocytosis are critical for synaptic transmission, whether the decrease of the former and the increase of the latter (which are both induced by an increased membrane fluidity) will compensate each other, or whether one effect will dominate, remains an open question. However, it was also suggested that centrifugation hypergravity stress alters both neurotransmitter reuptake and exocytotic release of glutamate⁶⁰: the effect of hypergravity (resp. microgravity) could thus be computationally represented as a decrease (resp. increase) of the connection weights in the simulated network. We added an offset w_e^1 to the base post-synaptic potential amplitude w_e . Interestingly, increasing the connection weights in the network increases its overall firing rate (Fig. 4A, left), as observed in microgravity, but also reduces the bursting rate (Fig. 4A, right): the inter-burst interval grows with w_e^1 , but so does the length of each burst (Fig. 4B), ultimately yielding an increased firing rate. This negative correlation between firing rate and burst rate in bursting networks had already been identified and discussed in previous theoretical studies³⁰.

Discussion

Neuronal activity has been observed to be gravity-dependent. More specifically, MEA recordings have shown that the firing and burst rates of cultured stem cells increase during drop tower-induced microgravity and decrease during centrifuge-induced hypergravity, with important variability between neuron subtypes²¹. This is in line with calcium activity recordings of hippocampal neurons obtained during parabolic flights, which showed that more neurons are active during altered gravity than during the 1 g baseline period, and that inter-spike intervals are shorter during microgravity phases⁶¹.

However, literature on how to model this phenomenon is sparse. On the one hand, a qualitative model has been proposed, based on the influence of gravity on ion channels^{23,24}; but it neither accounts for correlated dynamics of ion channels nor for possible connectivity between neurons. On the other hand, classical computational models, such as the Hodgkin–Huxley model, or recurrent networks of in silico neurons, do not incorporate a gravity parameter that enables the representation of altered gravity effects. Here, our approach proposes computational

Table 1 | Parameter values used in Fig. 1 (similar values as in ref. 66)

Name	Value	Description
A	$2 \times 10^{-4} \text{ cm}^2$	Membrane area
C_m	$1 \mu\text{F} \cdot \text{cm}^{-2} \cdot \text{A}$	Membrane capacitance
$\bar{g}_L$	$5 \times 10^{-5} \text{ S} \cdot \text{cm}^{-2} \cdot \text{A}$	Membrane conductance
$\bar{g}_K$	$30 \times 10^{-3} \text{ S} \cdot \text{cm}^{-2} \cdot \text{A}$	Potassium conductance
$\bar{g}_{Na}$	$100 \times 10^{-3} \text{ S} \cdot \text{cm}^{-2} \cdot \text{A}$	Sodium conductance
E_L	-60 mV	Leakage reversal potential
E_K	-90 mV	Potassium reversal potential
E_{Na}	50 mV	Sodium reversal potential
V_{th}	-63 mV	Spike threshold potential
τ_e	$5 \times 10^{-3} \text{ s}$	Decay time constant (excitatory synapses)
τ_i	$10 \times 10^{-3} \text{ s}$	Decay time constant (inhibitory synapses)
w_e	6 nS	Conductance quantum (excitatory synapses)
w_i	67 nS	Conductance quantum (inhibitory synapses)
E_e	0 mV	Reversal potential (excitatory synapses)
E_i	-80 mV	Reversal potential (inhibitory synapses)

The detailed equations for the Hodgkin–Huxley model and the COBA synapses used in Fig. 1 are also described in ref. 66.




Fig. 2 | Observed synchronized bursts can be explained by spike frequency adaptation. **A** Raster plot of detected action potentials on 14 units observed from a human stem cell-derived neuronal network (same setting as in ref. 21). Spike sorting was performed using SpikeInterface⁷⁷ and the MountainSort 5 algorithm⁷⁸.

B Membrane voltage of 2 example cells from a simulated network of connected LIF neurons (without inhibition but with firing rate adaptation). Dashed dark line: action potential threshold. Numerical values used for the simulations are described in Table 2. Bursts happen regularly approx. every 5 s.




Fig. 3 | The model reproduces features of network activity observed in altered gravity. **A** Simulated raster plot of the same network as in Fig. 2B for different values of I in Eq. (14). λ Top: $I_\lambda = 0$ pA. Bottom: $I_\lambda = 10$ pA. **B** Firing rate (left) and bursting

rate (right) in the same network for different values of I_λ . Solid line: Value averaged over all neurons and over ten repetitions with random initial cell membranes. Shaded area: s.e.m.




Fig. 4 | Gravity-induced modifications of synaptic transmission. **A** Firing rate (left) and bursting rate (right) in the same network as in Fig. 2B for different values of w_e^λ added to w_e in Table 2. Solid line: value averaged over all neurons and over ten

repetitions with random initial cell membranes. Shaded area: s.e.m. **B** Voltage traces for two example neurons and for different values of w_e^λ , showing the increasing inter-burst interval with increasing values of w_e^λ .

implementations of how gravity impacts neuronal activity, which are coherent with both recorded neuronal activity under altered gravity and previous observations on how a variety of cellular functions are gravity-dependent.

Many different parameters could be modified *in silico* to yield a change in firing and burst rates. However, a mechanism through which gravity alters neuronal activity should satisfy two points: First, it needs to be fast since modified neuronal activity upon altered gravity exposure happens on very short (seconds) time scales⁶¹. Second, variations should have been observed to occur during altered gravity experiments beforehand. A first effect of gravity was repeatedly observed on the fluidity of the cellular membrane^{23,25,37,38}. Based on this effect, a seminal explanatory model states that exposure to altered gravity modifies the physical properties of the cells' membrane, ultimately altering their voltage dynamics^{23,24}. This finding aligns with previous theoretical studies which demonstrated how firing rates can be modulated by the dynamical properties of the cell membrane when exposed to the same stationary input current, thereby facilitating prospective decoding⁶². Since ion channel dynamics are gravity-dependent^{34–36} and influenced by the mechanical properties of the cell membrane^{39–45}, we represent this effect by a linear scaling of the channels' gating rate (Eqs. 9–11): we verify that it allows to reproduce altered firing and bursting rates, both in single neurons (Fig. 1A–C) and in connected networks (Fig. 1D).

Secondly, a previous study using centrifuge-induced hypergravity reported an increase in neuronal activity both during the ramp-up (i.e., the

transition from 1g to hypergravity) and ramp-down (i.e., the transition from hypergravity to 1g) phases. This effect was attributed to MG ion channels, which are present in neuronal cells²⁸ and are known to induce action potentials²⁹. Mechanical loads and deformation of cells during acceleration and deceleration phases might thus impact neuronal activity²¹. The effect of an additional excitatory current is easy to analyze for single cells, but may not be trivial in networks of connected neurons. We show that the main features of the cell lines used in altered gravity experiments (i.e., synchronized activity with low-frequency bursts) can be reproduced by a network of connected excitatory cells stabilized by firing rate adaptation (Fig. 2), and that its firing and bursting rates will increase upon addition of an excitatory current (Fig. 3).

Our overall contribution is several-fold. Based on previous studies, we identified mechanisms which may explain the fast neuronal responses to altered gravity (namely: the influence of membrane fluidity on ion channel dynamics and synaptic transmission, and the activation of depolarizing MG channels). We proposed simple computational implementations of these mechanisms, which showed results coherent with previous observations. Note that these identified phenomena do not mutually exclude each other, but potentially originate from additive effects which future experiments should aim to disentangle (see below). Indeed, our modeling assumptions suggest that these effects could be additive (see the increase in firing rates with both I and λ in Figs. 1A and S2). We address contradictory previous results on the effect of altered gravity on ion channel dynamics^{23,24,34–36} and

Table 2 | Parameter values used in Figs. 2–4

Name	Value	Description
C_m	200 pF	Total capacitance
g_L	10 nS	Total leak conductance
E_L	−60 mV	Effective rest potential
Δ_T	2 mV	Threshold slope factor
V_T	−50 mV	Effective threshold potential
a	2 nS	Adaptation conductance
τ_w	800 mV	Adaptation time constant
b	30 pA	Spike-triggered adaptation
V_r	−46 mV	Reset potential
I	102 pA	Injected current
p	0.02	Connection probability
w_e	0.5 mV	Post-synaptic potential amplitude

Values for C_m , g_L , E_L , V_T , Δ_T , a , and V_r are the same as those used to simulate regular bursting in ref. 54. Values for τ_w , b , and I were adjusted to fit the long inter-burst intervals observed in experimental data (Fig. 2A).

propose a unified explanation for the effect of altered gravity on membrane fluidity, ion channel dynamics, and neuronal firing rates. Finally, we fitted recordings with a network model including firing rate adaptation: this proposed computational model explains the activity of networks of neurons used in altered gravity experiments, and especially their coordinated, low-frequency burst patterns in the absence of inhibitory cells.

Our approach has evident limitations. Our single-neuron analysis is based on the Hodgkin–Huxley model, which is a middle point between the simplicity of LIF neurons (which do not model the dynamics of single channels) and the realism of biophysical neuron models (e.g., ref. 63). The Hodgkin–Huxley model is largely descriptive rather than mechanistic, and does not account for several cellular mechanisms. Future work should consider extending our analysis to more extensive neuron models that were designed to resemble neuronal networks *in vivo* more reliably (e.g., ref. 49).

Furthermore, the goal of our approach is not to provide a mechanism for the effect of gravity on neuronal activity. Rather, our approach is a bottom-up process: using the measurement results of prior experiments on the effect of altered gravity on neuronal functions, we proposed computational implementations which are coherent with the observed effect of altered gravity on neuronal firing rates. Nevertheless, our results could greatly benefit from ad-hoc intracellular recordings under altered gravity conditions. Previous data on neuronal activity under altered gravity were obtained using either extracellular recordings²¹ or calcium imaging⁶¹, as these are suitable measurement approaches for the extreme environmental conditions at which the experiment needs to be performed. A remote-controlled device for patch-clamping has been developed⁶⁴ and implemented on a drop tower and parabolic flights to study action potential propagation of a simple model system without too complex neuronal networks⁶⁵. However, intracellular recordings under altered gravity remain rare due to the difficulty in carrying out such experiments and the sensitivity of the setup. Future experiments could be designed to isolate different effects, e.g., by pharmacological silencing or stimulating MG channels or modifying the experimental setup to follow potential changes in neuronal membrane fluidity. Patch clamp and sequencing experiments could help in further specifying the type and dynamics of the MG channels present, and to refine the assumptions used in our models.

The models we use in our simulations (and especially the Hodgkin–Huxley model) rely on a large number of free parameters (see Tables 1 and 2). However, these parameters can be easily constrained to small ranges of biologically plausible values. Seminal papers on conductance-based neurons⁶⁶ and firing rate adaptation⁵⁴ have proposed biologically plausible values for these parameters, which we based our simulations on. Some of

these parameters (namely the adaptation parameters τ_w and b , and the connectivity parameters I , p , and w_e in Table 2) were adjusted to match the specificities of the studied neurons (and especially to reproduce the synchronized, low-frequency bursting activity observed in Fig. 2A and in ref. 21). We verified that these values are in line with those previously used in seminal studies on modeling neuron networks⁴⁸. In Tables 1 and 2, the values for the capacitance C_m and leakage reversal potential E_L are in line with those reported for actual iNGN neuronal cells⁶⁷. Future experiments could focus on estimating the exact value of the other parameters (i.e., the ion-specific parameters in Table 1, or the connectivity parameters in Table 2) for iNGN neurons using dedicated intracellular recordings. Overall, we argue that our results are based on realistic values of the simulation parameters, and that they are robust to reasonable variations in these values (Fig. S2).

The altered response observed in cell cultures usually immediately follows the onset of microgravity. However, in longer-term altered gravity experiments (i.e., in sounding rocket experiments), neuronal activity recordings, utilizing the same experimental hardware as in the cited drop tower and centrifuge studies, suggest a complex and dynamic response characterized by a similar initial increase in firing rates followed by a transient decline (likely overcompensation) and partial recovery over several minutes. The computational models outlined here only account for the former (i.e., the short-term modification of neuronal activity following exposure to microgravity, for which quantitative analyses and numerical verification were lacking). To account for the latter (i.e., the compensation of neuronal activity trying to approach baseline levels following prolonged exposure to altered gravity), many different phenomena and models have been proposed to explain responses to activity perturbation, including individual neurons' spike frequency adaptation^{68–70} and synaptic homeostatic plasticity^{71–73}. Further exploration into the dynamics of neuronal activity during prolonged exposure to microgravity will be essential to determine the persistence of changes in firing rates. Such long-term investigations could reveal implications for brain function, potentially leading to acute and chronic issues. More generally, whether results obtained at the cellular levels will translate at the behavioral and cognitive levels remains an open question. Recent studies performed during parabolic flights have reported a modification of the EEG signal, but not of the participants' cognitive performances, under different micro- and hypergravity conditions^{16,74}: this hints at the existence of adaptation mechanisms, making cognition robust to altered gravity levels despite its effect on cultured cells.

Overall, our approach fills an important gap in the literature, by reviewing the mechanisms through which gravity levels could alter neuronal activity. In the absence of intracellular recordings obtained under various gravity levels, and following the principle of Occam's razor, our computational implementations of these phenomena remain simple: linear scaling of the gating time constants, and an additional uniform excitatory current. Future work will focus on refining these modeling assumptions by obtaining intracellular recordings at different gravity levels.

Methods

Simulation of single neurons

In Fig. 1, the voltage of simulated neurons is governed by Eqs. (1), (2), (9–11), with parameter values listed in Table 1. In Fig. 1D, we further implemented synaptic connections between simulated neurons. Following⁶⁶, the voltage of neuron k is given by

$$C_m \frac{dV_k}{dt} = -i_m - \sum_j g_{jk}(t)(V_k - E_j) \quad (15)$$

where i_m is given by Eq. (2) and $g_{jk}(t)(V_k - E_j)$ represents the synaptic connection between neurons k and j . The value of the synaptic reversal potential E_j depends on whether the synapse is excitatory or inhibitory (see values in Table 1). The synaptic conductance g_{jk} decays exponentially with time constants τ_e (for excitatory synapses) and τ_i (for inhibitory synapses), and is incremented by a quantum w_e or w_i whenever an action potential is generated by neuron j .

Simulation of networks of neurons

In Figs. 2–4, the evolution of the voltage of each individual neuron is governed by Eqs. (12–14), with parameter values listed in Table 2. Furthermore, voltage-jump synapses are implemented as follows: a synaptic connection from neuron j to k is randomly created with probability p . If a synaptic connection is established, the voltage of neuron k is immediately increased by a quantum w_e whenever neuron j generates an action potential.

Experimental data

This study mostly refers to previously analyzed and published data. Only in Fig. 2A are original data from a recent rocket flight presented for illustration purposes.

Methods and procedures that were employed to collect the data were previously described in ref. 21. In brief, we used human induced pluripotent stem cell (hiPSC)-derived neurons (iNGN cells) to conduct the experiments. Upon induction via the TetOn system, Neurogenin-1 and Neurogenin-2 were expressed, and cells differentiated into post-mitotic, bipolar neurons⁷⁵. iNGN cells were cultured and reseeded on MEAs following a previously published protocol⁷⁶. Thawed iNGN cells were cultured on Matrigel-coated plates in mTeSR 1 medium (STEMCELL Technologies, Germany). After at least two passages, cells were induced with 0.5 μ g/ml doxycycline for three days. At 2 dpi, 5 μ M Ara-C was added to eliminate undifferentiated cells. Concurrently, MEAs were prepared with poly-D-lysine and laminin coatings. Induced cells were reseeded on MEA chips at 3 dpi. Cells were dissociated, centrifuged, and resuspended in BrainPhys[®] medium (STEMCELL Technologies, Germany), and 100,000 cells were seeded per MEA. BrainPhys[®] medium was supplemented with neurotrophic factors and ascorbic acid. Astrocyte-conditioned media were added for long-term culture. Media was partially refreshed weekly with a 1:1 mix of fresh and astrocyte-conditioned BrainPhys[®] medium.

The experimental setup included a MEA system housed in a pressure chamber to maintain physiological conditions. The MEA's integrated heaters kept cells at 37 °C, while the pressure-tight chamber ensured stable pressure and CO₂ levels. The system's compact design is compatible with various altered gravity research platforms (DLR SAHC, ZARM drop tower, parabolic flights, and sounding rockets). A rail system allows for easy removal of the experiment module, facilitating time-sensitive operations like those on sounding rockets.

After culturing MEA samples for several weeks, they were brought to respective experiment facilities using transport incubators (CellTrans 2018 and 4016, Labotect, Germany; 37 °C, 5% CO₂).

Launching rocket experiments were carried out at the ESA ESRANGE space center in Kiruna, Sweden. iNGN MEA cultures were transported to the experiment site at least 10 days before launch. During the hot count-down and right before late access, MEA chips were integrated into the BIODECODER module, and functionality of the system was checked. The entire module was kept under a constant temperature of 37 °C within a custom-designed transport incubator (DLR). Upon late access, which is a time window between 2.5 and 1 h before launch, the BIODECODER module containing the samples was loaded into the dedicated slot within the rocket. Recording was started 10 min prior to launch, and data were recorded continuously throughout the entire flight, including post-baseline recording back on the ground. Battery operation ensured constant heating of the samples after launch and landing. The module was retrieved by a helicopter and brought back to the launch site, where additional post-experiment recordings and microscopy were executed.

Data availability

Data shown in Fig. 2A are not publicly available due to them being currently used in yet unpublished studies but are available from the corresponding author on reasonable request.

Code availability

Analysis code is available from the following repository: https://github.com/camillegontier/neuron_microgravity.

Received: 2 September 2024; Accepted: 23 July 2026;

Published online: 07 August 2026

References

1. di Prampero, P. E. & Narici, M. V. Muscles in microgravity: from fibres to human motion. *J. Biomech.* **36**, 403–412 (2003).
2. Droppert, P. A review of muscle atrophy in microgravity and during prolonged bed rest. *J. Br. Interplanet. Soc.* **46**, 83–86 (1993).
3. Grimm, D. et al. The impact of microgravity on bone in humans. *Bone* **87**, 44–56 (2016).
4. Hargens, A. R. & Richardson, S. Cardiovascular adaptations, fluid shifts, and countermeasures related to space flight. *Respir. Physiol. Neurobiol.* **169**, S30–S33 (2009).
5. Lawley, J. S. et al. Effect of gravity and microgravity on intracranial pressure. *J. Physiol.* **595**, 2115–2127 (2017).
6. Lee, A. G., Mader, T. H., Gibson, C. R., Brunstetter, T. J. & Tarver, W. J. Space flight-associated neuro-ocular syndrome (sans). *Eye* **32**, 1164–1167 (2018).
7. Lee, A. G. et al. Spaceflight associated neuro-ocular syndrome (sans) and the neuro-ophthalmologic effects of microgravity: a review and an update. *npj Microgravity* **6**, 1–10 (2020).
8. Roberts, D. et al. Prolonged microgravity affects human brain structure and function. *Am. J. Neuroradiol.* **40**, 1878–1885 (2019).
9. Popova, N. K., Kulikov, A. V. & Naumenko, V. S. Spaceflight and brain plasticity: spaceflight effects on regional expression of neurotransmitter systems and neurotrophic factors encoding genes. *Neurosci. Biobehav. Rev.* **119**, 396–405 (2020).
10. Herranz, R. et al. Ground-based facilities for simulation of microgravity: organism-specific recommendations for their use, and recommended terminology. *Astrobiology* **13**, 1–17 (2013).
11. Hauslage, J. et al. Cytosolic calcium concentration changes in neuronal cells under clinorotation and in parabolic flight missions. *Microgravity Sci. Technol.* **28**, 633–638 (2016).
12. Callens, N. et al. Esa parabolic flights, drop tower and centrifuge opportunities for university students. *Microgravity Sci. Technol.* **23**, 181–189 (2011).
13. Liemersdorf, C., Lichterfeld, Y., Hemmersbach, R. & Hauslage, J. The mapheus module cellfix for studying the influence of altered gravity on the physiology of single cells. *Rev. Sci. Instrum.* **91**, 014101 (2020).
14. Hauslage, J. et al. Arabidomics—a new experimental platform for molecular analyses of plants in drop towers, on parabolic flights, and sounding rockets. *Rev. Sci. Instrum.* **91**, 034504 (2020).
15. Pletser, V. Short duration microgravity experiments in physical and life sciences during parabolic flights: the first 30 ESA campaigns. *Acta Astronaut.* **55**, 829–854 (2004).
16. Badali, C., Wollseiffen, P. & Schneider, S. Shades of gravity-effects of planetary gravity levels on electrocortical activity and neurocognitive performance. *Brain Struct. Funct.* **229**, 1–13 (2024).
17. Acharya, A. et al. Parabolic, flight-induced, acute hypergravity and microgravity effects on the beating rate of human cardiomyocytes. *Cells* **8**, 352 (2019).
18. Selig, H., Gierse, A. & König, G. Parabolic flight with light aircraft. In *67th International Astronautical Congress (IAC)*. *International Astronautical Federation (IAF)*, Guadalajara. (IAC-16-A2-5-6-x34402) (IAF, 2016).
19. Caprace, D.-G., Gontier, C., Iranmanesh, M., Scoubeau, M. & Pletser, V. Experimental characterization of weightlessness during glider parabolic flights. *Microgravity Sci. Technol.* **32**, 1121–1132 (2020).
20. Lichterfeld, Y. et al. Hypergravity attenuates reactivity in primary murine astrocytes. *Biomedicine* **10**, 1966 (2022).
21. Striebel, J. et al. Human neural network activity reacts to gravity changes in vitro. *Front. Neurosci.* **17**, 1085282 (2023).
22. Pletser, V., Frischauf, N., Laufer, R. & Cohen, D. Parabolic flights with gliders as an innovative low-cost platform for microgravity and hypergravity research. In *68th International Astronautical Congress*

- (IAC). *International Astronautical Federation (IAF), Adelaide*. (IAC-17-A2-5-6-x36752) 1–5 (IAC, 2017).
23. Kohn, F., Hauslage, J. & Hanke, W. Membrane fluidity changes, a basic mechanism of interaction of gravity with cells? *Microgravity Sci. Technol.* **29**, 337–342 (2017).
24. Kohn, F. P. & Ritzmann, R. Gravity and neuronal adaptation, in vitro and in vivo—from neuronal cells up to neuromuscular responses: a first model. *Eur. Biophys. J.* **47**, 97–107 (2018).
25. Kohn, F. P. & Hauslage, J. The gravity dependence of pharmacodynamics: the integration of lidocaine into membranes in microgravity. *npj Microgravity* **5**, 5 (2019).
26. Häder, D.-P., Braun, M., Grimm, D. & Hemmersbach, R. Gravireceptors in eukaryotes—a comparison of case studies on the cellular level. *npj Microgravity* **3**, 13 (2017).
27. Kirischuk, S. Mechanosensitive channels in neuronal and astroglial cells in the nervous system. In *Mechanosensitivity of the Nervous System: Forewords by Nektarios Tavernarakis and Pontus Persson* 3–22 (2009).
28. Falleroni, F. et al. Mechanotransduction in hippocampal neurons operates under localized low piconewton forces. *Isience* **25**, 103807 (Springer, 2022).
29. Nikolaev, Y. A., Dosen, P. J., Laver, D. R., Van Helden, D. F. & Hamill, O. P. Single mechanically-gated cation channel currents can trigger action potentials in neocortical and hippocampal pyramidal neurons. *Brain Res.* **1608**, 1–13 (2015).
30. Sukenik, N. et al. Neuronal circuits overcome imbalance in excitation and inhibition by adjusting connection numbers. *Proc. Natl. Acad. Sci. USA* **118**, e2018459118 (2021).
31. Habibey, R., Striebel, J., Schmieder, F., Czarske, J. & Busskamp, V. Long-term morphological and functional dynamics of human stem cell-derived neuronal networks on high-density micro-electrode arrays. *Front. Neurosci.* **16**, 951964 (2022).
32. Hodgkin, A. L. & Huxley, A. F. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J. Physiol.* **117**, 500 (1952).
33. Dayan, P. & Abbott, L. F. *Theoretical Neuroscience: Computational and Mathematical Modeling of Neural Systems* (MIT Press, 2005).
34. Klinke, N., Goldermann, M. & Hanke, W. The properties of alamethicin incorporated into planar lipid bilayers under the influence of microgravity. *Acta Astronaut.* **47**, 771–773 (2000).
35. Hanke, W. Studies of the interaction of gravity with biological membranes using alamethicin doped planar lipid bilayers as a model system. *Adv. Space Res.* **17**, 143–150 (1996).
36. Goldermann, M. & Hanke, W. Ion channel are sensitive to gravity changes. *Microgravity Sci. Technol.* **13**, 35–38 (2001).
37. Asuwin Prabu, R., Manohar, A., Narendran, S., Kabir, A. & Sudhakar, S. Effect of simulated microgravity on artificial single cell membrane mechanics. *Microgravity Sci. Technol.* **36**, 47 (2024).
38. Klymchuk, D. et al. Properties of plasma membrane from pea root seedlings under altered gravity. In *35th COSPAR Scientific Assembly*, Vol. 35, 1356 (COSPAR, 2004).
39. Dart, C. Symposium review: Lipid microdomains and the regulation of ion channel function. *J. Physiol.* **588**, 3169–3178 (2010).
40. Duncan, A. L., Song, W. & Sansom, M. S. Lipid-dependent regulation of ion channels and g protein-coupled receptors: insights from structures and simulations. *Annu. Rev. Pharmacol. Toxicol.* **60**, 31–50 (2020).
41. Roberts-Crowley, M. L., Mitra-Ganguli, T., Liu, L. & Rittenhouse, A. R. Regulation of voltage-gated Ca²⁺ channels by lipids. *Cell Calcium* **45**, 589–601 (2009).
42. Schmidpeter, P. A. et al. Membrane phospholipids control gating of the mechanosensitive potassium leak channel *trk1*. *Nat. Commun.* **14**, 1077 (2023).
43. Sunshine, C. & McNamee, M. G. Lipid modulation of nicotinic acetylcholine receptor function: the role of neutral and negatively charged lipids. *Biochim. Biophys. Acta* **1108**, 240–246 (1992).
44. Sunshine, C. & McNamee, M. G. Lipid modulation of nicotinic acetylcholine receptor function: the role of membrane lipid composition and fluidity. *Biochim. Biophys. Acta* **1191**, 59–64 (1994).
45. Zanello, L., Aztiria, E., Antolini, S. & Barrantes, F. Nicotinic acetylcholine receptor channels are influenced by the physical state of their membrane environment. *Biophys. J.* **70**, 2155–2164 (1996).
46. Stimberg, M., Brette, R. & Goodman, D. F. Brian 2, an intuitive and efficient neural simulator. *eLife* **8**, e47314 (2019).
47. Lu, C. et al. Overexpression of *neurog2* and *neurog1* in human embryonic stem cells produces a network of excitatory and inhibitory neurons. *FASEB J.* **33**, 5287 (2019).
48. Kriener, B. et al. Dynamics of self-sustained asynchronous-irregular activity in random networks of spiking neurons with strong synapses. *Front. Comput. Neurosci.* **8**, 136 (2014).
49. Compte, A., Sanchez-Vives, M. V., McCormick, D. A. & Wang, X.-J. Cellular and network mechanisms of slow oscillatory activity (<1 Hz) and wave propagations in a cortical network model. *J. Neurophysiol.* **89**, 2707–2725 (2003).
50. Capone, C., Pastorelli, E., Golosio, B. & Paolucci, P. S. Sleep-like slow oscillations improve visual classification through synaptic homeostasis and memory association in a thalamo-cortical model. *Sci. Rep.* **9**, 8990 (2019).
51. Torao-Angosto, M., Manasanch, A., Mattia, M. & Sanchez-Vives, M. V. Up and down states during slow oscillations in slow-wave sleep and different levels of anesthesia. *Front. Syst. Neurosci.* **15**, 609645 (2021).
52. Gutkin, B. & Zeldenrust, F. Spike frequency adaptation. *Scholarpedia* **9**, 30643 (2014).
53. Vreeswijk, C. V. & Hansel, D. Patterns of synchrony in neural networks with spike adaptation. *Neural Comput.* **13**, 959–992 (2001).
54. Naud, R., Marcille, N., Clopath, C. & Gerstner, W. Firing patterns in the adaptive exponential integrate-and-fire model. *Biol. Cybern.* **99**, 335–347 (2008).
55. Gauthier, N. C., Fardin, M. A., Roca-Cusachs, P. & Sheetz, M. P. Temporary increase in plasma membrane tension coordinates the activation of exocytosis and contraction during cell spreading. *Proc. Natl. Acad. Sci. USA* **108**, 14467–14472 (2011).
56. Masters, T. A., Pontes, B., Viasnoff, V., Li, Y. & Gauthier, N. C. Plasma membrane tension orchestrates membrane trafficking, cytoskeletal remodeling, and biochemical signaling during phagocytosis. *Proc. Natl. Acad. Sci. USA* **110**, 11875–11880 (2013).
57. Dai, J., Ting-Beall, H. P. & Sheetz, M. P. The secretion-coupled endocytosis correlates with membrane tension changes in rbl 2h3 cells. *J. Gen. Physiol.* **110**, 1–10 (1997).
58. Raucher, D. & Sheetz, M. P. Membrane expansion increases endocytosis rate during mitosis. *J. Cell Biol.* **144**, 497–506 (1999).
59. Wu, X.-S. et al. Membrane tension inhibits rapid and slow endocytosis in secretory cells. *Biophys. J.* **113**, 2406–2414 (2017).
60. Borisova, T., Krisanova, N. & Himmelreich, N. Exposure of animals to artificial gravity conditions leads to the alteration of the glutamate release from rat cerebral hemispheres nerve terminals. *Adv. Space Res.* **33**, 1362–1367 (2004).
61. Lecoq, P.-E. et al. Influence of microgravity on spontaneous calcium activity of primary hippocampal neurons grown in microfluidic chips. *npj Microgravity* **10**, 15 (2024).
62. Brandt, S., Petrovici, M. A., Senn, W., Wilmes, K. A. & Benitez, F. Prospective and retrospective coding in cortical neurons. Preprint at <https://arxiv.org/abs/2405.14810> (2024).
63. Deistler, M. et al. Differentiable simulation enables large-scale training of detailed biophysical models of neural dynamics. Preprint at <https://doi.org/10.1101/2024.08.21.608979> (2024).
64. Meissner, K. & Hanke, W. Patch-clamp experiments under microgravity. In *Life in Space for Life on Earth*, Vol. 501, 399–400 (Springer, 2002).
65. Meissner, K. & Hanke, W. Action potential properties are gravity dependent. *Microgravity Sci. Technol.* **17**, 38–43 (2005).

66. Brette, R. et al. Simulation of networks of spiking neurons: a review of tools and strategies. *J. Comput. Neurosci.* **23**, 349–398 (2007).
67. Lam, R. S., Töpfer, F. M., Wood, P. G., Busskamp, V. & Bamberg, E. Functional maturation of human stem cell-derived neurons in long-term cultures. *PLoS One* **12**, e0169506 (2017).
68. Benda, J. & Herz, A. V. A universal model for spike-frequency adaptation. *Neural Comput.* **15**, 2523–2564 (2003).
69. La Camera, G. et al. Multiple time scales of temporal response in pyramidal and fast spiking cortical neurons. *J. Neurophysiol.* **96**, 3448–3464 (2006).
70. Fuhrmann, G., Markram, H. & Tsodyks, M. Spike frequency adaptation and neocortical rhythms. *J. Neurophysiol.* **88**, 761–770 (2002).
71. Delvendahl, I. & Müller, M. Homeostatic plasticity—a presynaptic perspective. *Curr. Opin. Neurobiol.* **54**, 155–162 (2019).
72. Frank, C. A., James, T. D. & Müller, M. Homeostatic control of Drosophila neuromuscular junction function. *Synapse* **74**, e22133 (2020).
73. Davis, G. W. & Müller, M. Homeostatic control of presynaptic neurotransmitter release. *Annu. Rev. Physiol.* **77**, 251–270 (2015).
74. Badali, C., Wollseiffen, P. & Schneider, S. Under pressure—the influence of hypergravity on electrocortical activity and neurocognitive performance. *Exp. Brain Res.* **241**, 2249–2259 (2023).
75. Busskamp, V. et al. Rapid neurogenesis through transcriptional activation in human stem cells. *Mol. Syst. Biol.* **10**, 760 (2014).
76. Schmieder, F. et al. Tracking connectivity maps in human stem cell-derived neuronal networks by holographic optogenetics. *Life Sci. Alliance* **5**, e202101268 (2022).
77. Buccino, A. P. et al. Spikeinterface, a unified framework for spike sorting. *eLife* **9**, e61834 (2020).
78. Chung, J. E. et al. A fully automated approach to spike sorting. *Neuron* **95**, 1381–1394 (2017).

Acknowledgements

We thank Martin Müller, Nicolas Vitale, Jakob Jordan, Aitor Morales-Gregorio, Igor Delvendahl, Roberto Martins de Freitas, Denis-Gabriel Caprace, Mohammad Iran-manesh, Mehdi Scoubeau, Simon Brandt, Timo Gierlich, Ben von Hünerbein, and Jérémy Rabineau for the fruitful discussions. This study was funded by the German Aerospace Center. J.S. acknowledges support by the Joachim Herz Foundation. The funders played no role in study design, data collection, analysis and interpretation of data, or the writing of this manuscript.

Author contributions

C.G. performed the analysis and simulations and wrote the manuscript. L.D., J.S., M.S., Z.M., S.S., Y.L., and C.L. designed the experiments and acquired the data. All authors contributed to reviewing and editing the paper.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing interests

C.G. is involved with LIDE Space, a non-profit association that provides glider-based microgravity services. The other authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at

<https://doi.org/10.1038/s41526-026-00644-7>.

Correspondence and requests for materials should be addressed to Christian Liemersdorf.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026