

Second Harmonic Generation by BaTiO₃ microparticles in porous materials

S. Lisinski, D. Schaniel, Th. Woike, L. Ratke, M. Imlau

DLR, Institut für Materialphysik im Weltraum, Linder Höhe, 51147 Köln, Germany

Universität zu Köln, I. Physikalisches Institut., Zülpicher Str. 49b, 50937 Köln, Germany


Universität Osnabrück, Fachbereich Physik, 49069 Osnabrück, Germany

Motivation

Preparation of random medias with nonlinear optical properties

- cheap nonlinear optical (NLO) materials
- easy synthesis of NLO materials: Sol-Gel method
- novel applications
- use of disordered materials for NLO applications, e.g., random laser, frequency conversion

Aim: Embedding of NLO materials (powder) in aerogel/xerogel matrix





- Synthesis of silica and titana aerogels/xerogels via sol-gel method
- Embedding of NLO particles during liquid phase
- Homogeneous distribution of particles in aerogel/xerogel matrix
- Measuring of SHG

Def.: Aerogels are supercritical dried gels (90-99% pores) and xerogels are dried under ambient conditions (<50% pores)

Properties of aerogels and xerogels:

- 3D nano-network of colloidal particles (SiO_2 or TiO_2)
- Transparent in the range 280-2700 nm
- refractive index tunable in the range 1.2-2.1



Transparency range of dry gels:


SiO_2 aerogel: 330-2650nm

SiO_2 xerogel: 280-2460nm

TiO_2 xerogel: 390-1880nm


NaWG xerogel: 350-2680nm

NaWG=Sodium waterglass



Refractive index matching


Mixed SiO_2 - TiO_2 xerogels have an adjustable refractive index in the range 1.2-2.1 and can thus be matched to the index of the particles, such that $n_{\text{aerogel/xerogel}} = n_{\text{powder}}$




For **second harmonic generation** we used the 1064nm pulse of a Nd:YAG laser (FWH=6ns, Input energy=0-80mJ). The SHG-effect is measured for BaTiO₃ particles with an average particle size of 1μm in a silica xerogel based on sodium waterglass



Scanning electron microscope picture of the BaTiO₃ particles with an average particle size of 1μm




The generated green light scatters in all directions, because of the random particle polarisation



Linear increase of SH intensity with incident intensity

Nd-YAG pump laser:

- incident energy 0-80mJ
- full width at half-maximum 6ns
- incident wavelength 1064nm



Concentration dependence, i.e. there is an "optimal" number density n of BaTiO₃ particles in the xerogel matrix

$$I_{\text{SHG}} \sim n e^{-n\alpha} = n e^{-n\sigma d}$$

α : scattering coefficient
 σ : cross section
 d : thickness of sample

