

**CONTROL ID:** 1808412

**TITLE:** An Overview of Lightning NO<sub>x</sub> Production Research Associated with the Deep Convective Clouds and Chemistry (DC3) Experiment (*Invited*)

**AUTHORS (FIRST NAME, LAST NAME):** Kenneth E Pickering<sup>1, 3</sup>, Mary C Barth<sup>2</sup>, Kristin Cummings<sup>3</sup>, Megan M Bela<sup>4</sup>, Yunyao Li<sup>3</sup>, Dale J Allen<sup>3</sup>, Eric J Bucsela<sup>5</sup>, Eric C Bruning<sup>6</sup>, Steven A Rutledge<sup>7</sup>, Brody Fuchs<sup>7</sup>, Heidi Huntrieser<sup>8</sup>, Ilana B Pollack<sup>9</sup>, Thomas B Ryerson<sup>9</sup>

**INSTITUTIONS (ALL):** 1. NASA Goddard Space Flight Cent, Greenbelt, MD, United States.

2. National Center for Atmospheric Research, Boulder, CO, United States.

3. University of Maryland, College Park, MD, United States.

4. University of Colorado, Boulder, CO, United States.

5. SRI, International, Menlo Park, CA, United States.

6. Texas Tech University, Lubbock, TX, United States.

7. Colorado State University, Fort Collins, CO, United States.

8. Deutsches Zentrum für Luft-und Raumfahrt, Oberpfaffenhofen, Germany.

9. NOAA Earth System Research Laboratory, Boulder, CO, United States.

**ABSTRACT BODY:** Major goals of the DC3 experiment include determining the contribution of lightning to NO<sub>x</sub> in the anvils of observed thunderstorms, examining the relationship of lightning NO<sub>x</sub> (LNO<sub>x</sub>) production to flash rates and to other lightning characteristics, and estimating the relative production per flash for cloud-to-ground flashes and intracloud flashes. The talk will survey the observation types that were conducted during DC3 relevant to these goals and provide an overview of the analysis and modeling techniques which are being used to achieve them. NO<sub>x</sub> was observed on three research aircraft during DC3 (the NCAR G-V, the NASA DC-8, and the DLR Falcon) in flights through storm anvils in three study regions (NE Colorado, Central Oklahoma to West Texas, and northern Alabama) where lightning mapping arrays (LMAs) and radar coverage were available. Initial comparisons of the aircraft NO<sub>x</sub> observations in storm anvils relative to flash rates have been conducted. These analyses utilize estimates of the LNO<sub>x</sub> within the storm volume and the flux of LNO<sub>x</sub> through the anvils, which when combined with observed flash rates can be used to estimate storm-average LNO<sub>x</sub> production per flash. The WRF-Chem model is being run for cloud-resolved simulations of selected observed storms during DC3. Detailed lightning information from the LMAs (gridded flash rates as a function of time and information on the vertical distributions of flash activity) are compared with model-parameterized flash rates and assumed vertical distributions. Assumptions concerning NO<sub>x</sub> production per CG flash and per IC flash are tested through comparisons of the model output with the aircraft NO<sub>x</sub> data from anvil traverses. A specially designed retrieval method for lightning NO<sub>2</sub> column amounts from the OMI instrument on NASA's Aura satellite has been utilized to estimate NO<sub>2</sub> over the region affected by selected DC3 storms. Combined with NO<sub>x</sub> to NO<sub>2</sub> ratios from the aircraft data and observed flash rates, average NO<sub>x</sub> production per flash can be estimated.

**KEYWORDS:** 3324 ATMOSPHERIC PROCESSES Lightning, 0322 ATMOSPHERIC COMPOSITION AND STRUCTURE Constituent sources and sinks, 0320 ATMOSPHERIC COMPOSITION AND STRUCTURE Cloud physics and chemistry, 3314 ATMOSPHERIC PROCESSES Convective processes.

(No Image Selected)

(No Table Selected)

**Additional Details**

**Previously Presented Material:** 20% Fall 2012 AGU Meeting

**Contact Details**

**CONTACT (NAME ONLY):** Kenneth Pickering

**CONTACT (E-MAIL ONLY):** kenneth.e.pickering@nasa.gov

**TITLE OF TEAM:**

---